



Australian Government

**Department of Health
and Aged Care**



Schedule of Pharmaceutical Benefits

Summary of Changes

Effective 1 October 2023



Fees, Patient Contributions and Safety Net Thresholds

The following fees, patient contributions and safety net thresholds apply as at 1 October 2023 and are included, where applicable, in prices published in the Schedule —

Dispensing Fees:	Ready-prepared	\$8.37
	Dangerous drug fee	\$5.18
	Extemporaneously-prepared	\$10.41
	Allowable additional patient charge*	\$3.29
Additional Fees (for safety net prices):	Ready-prepared	\$1.40
	Extemporaneously-prepared	\$1.80
Patient Co-payments:	General	\$30.00
	Concessional	\$7.30
Safety Net Thresholds:	General	\$1563.50
	Concessional	\$262.80
Safety Net Card Issue Fee:		\$11.42

* The allowable additional patient charge is a discretionary charge to general patients if a pharmaceutical item has a dispensed price for maximum quantity less than the general patient co-payment. The pharmacist may charge general patients the allowable additional fee but the fee cannot take the cost of the prescription above the general patient co-payment for the medicine. This fee does not count towards the Safety Net threshold.

Summary of Changes

These changes to the Schedule of Pharmaceutical Benefits are effective from 1 October 2023. The Schedule is updated on the first day of each month and is available on the internet at www.pbs.gov.au.

General Pharmaceutical Benefits

Additions

Addition – Item

13644D	BIMEKIZUMAB , bimekizumab 160 mg/mL injection, 2 x 1 mL pen devices (<i>Bimzelx</i>)
13652M	BIMEKIZUMAB , bimekizumab 160 mg/mL injection, 2 x 1 mL pen devices (<i>Bimzelx</i>)
13643C	CEFUROXIME , cefuroxime 125 mg/5 mL powder for oral liquid, 70 mL (<i>Zinnat (UK)</i>)
13653N	CEFUROXIME , cefuroxime 125 mg/5 mL powder for oral liquid, 70 mL (<i>Zinnat (UK)</i>)
13649J	DEUCRAVACITINIB , deucravacitinib 6 mg tablet, 28 (<i>Sotyktu</i>)
13640X	FOSNETUPITANT + PALONOSETRON , fosnetupitant 235 mg + palonosetron 250 microgram injection, 1 vial (<i>Akynzeo IV</i>)
13651L	INSULIN ASPART , insulin aspart 100 units/mL fast acting injection, 5 x 3 mL cartridges (<i>Fiasp Penfill</i>)

Addition – Brand

1081X	<i>Blooms The Chemist Atenolol, BG</i> – ATENOLOL , atenolol 50 mg tablet, 30
13540P	<i>Blooms The Chemist Atenolol, BG</i> – ATENOLOL , atenolol 50 mg tablet, 30
13391T	<i>BTC Candesartan HCT, BG</i> – CANDESARTAN + HYDROCHLOROTHIAZIDE , candesartan cilexetil 16 mg + hydrochlorothiazide 12.5 mg tablet, 30
8504N	<i>BTC Candesartan HCT, BG</i> – CANDESARTAN + HYDROCHLOROTHIAZIDE , candesartan cilexetil 16 mg + hydrochlorothiazide 12.5 mg tablet, 30
13452B	<i>BTC Candesartan HCT, BG</i> – CANDESARTAN + HYDROCHLOROTHIAZIDE , candesartan cilexetil 32 mg + hydrochlorothiazide 12.5 mg tablet, 30
9314F	<i>BTC Candesartan HCT, BG</i> – CANDESARTAN + HYDROCHLOROTHIAZIDE , candesartan cilexetil 32 mg + hydrochlorothiazide 12.5 mg tablet, 30
13392W	<i>BTC Candesartan HCT, BG</i> – CANDESARTAN + HYDROCHLOROTHIAZIDE , candesartan cilexetil 32 mg + hydrochlorothiazide 25 mg tablet, 30
9315G	<i>BTC Candesartan HCT, BG</i> – CANDESARTAN + HYDROCHLOROTHIAZIDE , candesartan cilexetil 32 mg + hydrochlorothiazide 25 mg tablet, 30
10778G	<i>Blooms The Chemist Cefalexin, BG</i> – CEFALEXIN , cefalexin 500 mg capsule, 20
11934D	<i>Blooms The Chemist Cefalexin, BG</i> – CEFALEXIN , cefalexin 500 mg capsule, 20
3119E	<i>Blooms The Chemist Cefalexin, BG</i> – CEFALEXIN , cefalexin 500 mg capsule, 20
3318P	<i>Blooms The Chemist Cefalexin, BG</i> – CEFALEXIN , cefalexin 500 mg capsule, 20
10790X	<i>DICLOXACILLIN VIATRIS 500, MQ</i> – DICLOXACILLIN , dicloxacillin 500 mg capsule, 24
5097G	<i>DICLOXACILLIN VIATRIS 500, MQ</i> – DICLOXACILLIN , dicloxacillin 500 mg capsule, 24
8122L	<i>DICLOXACILLIN VIATRIS 500, MQ</i> – DICLOXACILLIN , dicloxacillin 500 mg capsule, 24
9432K	<i>Lexam 10, RW</i> – ESCITALOPRAM , escitalopram 10 mg tablet, 28
9433L	<i>Lexam 20, RW</i> – ESCITALOPRAM , escitalopram 20 mg tablet, 28

1434L	<i>Blooms the Chemist Fluoxetine, BG</i> – FLUOXETINE , fluoxetine 20 mg capsule, 28
2592K	<i>Pharmacor Isotretinoin, CR</i> – ISOTRETINOIN , isotretinoin 20 mg capsule, 60
12969N	<i>Thyrox, IX</i> – LEVOTHYROXINE , levothyroxine sodium 50 microgram tablet, 2 x 100
12960D	<i>Thyrox, IX</i> – LEVOTHYROXINE , levothyroxine sodium 75 microgram tablet, 2 x 100
12968M	<i>Thyrox, IX</i> – LEVOTHYROXINE , levothyroxine sodium 100 microgram tablet, 2 x 100
12970P	<i>Thyrox, IX</i> – LEVOTHYROXINE , levothyroxine sodium 200 microgram tablet, 2 x 100
1622J	<i>Chexate, OX</i> – METHOTREXATE , methotrexate 2.5 mg tablet, 30
1623K	<i>Chexate, OX</i> – METHOTREXATE , methotrexate 10 mg tablet, 50
10839L	<i>Tanilone (Fatty), AS</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
10840M	<i>Tanilone (Fatty), AS</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
10843Q	<i>Tanilone (Fatty), AS</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
10844R	<i>Tanilone (Fatty), AS</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
10848Y	<i>Tanilone (Fatty), AS</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
8128T	<i>Tanilone (Fatty), AS</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
11406H	<i>Pirfenidone Sandoz, SZ</i> – PIRFENIDONE , pirfenidone 267 mg tablet, 90
11410M	<i>Pirfenidone Sandoz, SZ</i> – PIRFENIDONE , pirfenidone 801 mg tablet, 90

Addition – Equivalence Indicator

1622J	<i>Methoblastin, PF</i> – METHOTREXATE , methotrexate 2.5 mg tablet, 30
1623K	<i>Methoblastin, PF</i> – METHOTREXATE , methotrexate 10 mg tablet, 50
10839L	<i>Advantan (Fatty), LO</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
10840M	<i>Advantan (Fatty), LO</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
10843Q	<i>Advantan (Fatty), LO</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
10844R	<i>Advantan (Fatty), LO</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
10848Y	<i>Advantan (Fatty), LO</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g
8128T	<i>Advantan (Fatty), LO</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g

Addition – Note

13615N	AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES , amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides with 2-fucosyllactose and lacto-N-neotetraose powder for oral liquid, 400 g (<i>Alfamino</i>)
13627F	AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES , amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides with 2-fucosyllactose and lacto-N-neotetraose powder for oral liquid, 400 g (<i>Alfamino</i>)
2900P	AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES , amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides powder for oral liquid, 400 g (<i>Alfamino</i>)
2928D	AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES , amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides powder for oral liquid, 400 g (<i>Alfamino</i>)

Deletions

Deletion – Item

11798Y	OCRIPLASMIN , ocriplasmin 375 microgram/0.3 mL intraocular injection, 0.3 mL vial (<i>Jetrea RTU</i>)
11136D	PIRFENIDONE , pirfenidone 267 mg capsule, 270 (<i>Esbriet</i>)

Deletion – Brand

9012H	ALENDRONATE PLUS D3 70mg/70ug APOTEX, GX – ALENDRONATE + COLECALCIFEROL , alendronate 70 mg + colecalciferol 70 microgram (2800 units) tablet, 4
-------	--

9183H	<i>ALENDRONATE PLUS D3 70mg/140ug APOTEX, GX</i> – ALENDRONATE + COLECALCIFEROL , alendronate 70 mg + colecalciferol 140 microgram (5600 units) tablet, 4
9022W	<i>Fenofibrate Mylan, AF</i> – FENOFIBRATE , fenofibrate 48 mg tablet, 60
9023X	<i>Fenofibrate Mylan, AF</i> – FENOFIBRATE , fenofibrate 145 mg tablet, 30
2592K	<i>Isotretinoin SCP 20, CR</i> – ISOTRETINOIN , isotretinoin 20 mg capsule, 60
5264C	<i>Methylprednisolone Alphapharm, AF</i> – METHYLPREDNISOLONE , methylprednisolone 1 g injection, 1 vial
13579Q	<i>Moxonidine MYL, AF</i> – MOXONIDINE , moxonidine 200 microgram tablet, 30
9019Q	<i>Moxonidine MYL, AF</i> – MOXONIDINE , moxonidine 200 microgram tablet, 30
11683X	<i>Omeprazole generichealth, GQ</i> – OMEPRAZOLE , omeprazole 20 mg enteric tablet, 30
12272X	<i>Omeprazole generichealth, GQ</i> – OMEPRAZOLE , omeprazole 20 mg enteric tablet, 30
8331L	<i>Omeprazole generichealth, GQ</i> – OMEPRAZOLE , omeprazole 20 mg enteric tablet, 30
8333N	<i>Omeprazole generichealth, GQ</i> – OMEPRAZOLE , omeprazole 20 mg enteric tablet, 30
11406H	<i>Esbriet, RO</i> – PIRFENIDONE , pirfenidone 267 mg tablet, 90
11410M	<i>Esbriet, RO</i> – PIRFENIDONE , pirfenidone 801 mg tablet, 90
1968N	<i>Qpril 5, AF</i> – QUINAPRIL , quinapril 5 mg tablet, 30
1969P	<i>Qpril 10, AF</i> – QUINAPRIL , quinapril 10 mg tablet, 30
1970Q	<i>Qpril 20, AF</i> – QUINAPRIL , quinapril 20 mg tablet, 30
13607E	<i>Telmisartan HCT GH 80/25, GQ</i> – TELMISARTAN + HYDROCHLOROTHIAZIDE , telmisartan 80 mg + hydrochlorothiazide 25 mg tablet, 28
9381R	<i>Telmisartan HCT GH 80/25, GQ</i> – TELMISARTAN + HYDROCHLOROTHIAZIDE , telmisartan 80 mg + hydrochlorothiazide 25 mg tablet, 28

Deletion – Note

11406H	PIRFENIDONE , pirfenidone 267 mg tablet, 90 (<i>Pirfenidone Sandoz</i>)
11410M	PIRFENIDONE , pirfenidone 801 mg tablet, 90 (<i>Pirfenidone Sandoz</i>)

Deletion – Restriction

11858D	RISANKIZUMAB , risankizumab 75 mg/0.83 mL injection, 2 x 0.83 mL syringes (<i>Skyrizi</i>)
--------	---

Alterations

Alteration – Note

12342N	ADALIMUMAB , adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL pen devices (<i>Humira, Yuflyma</i>)
12377K	ADALIMUMAB , adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL pen devices (<i>Humira, Yuflyma</i>)
12421R	ADALIMUMAB , adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL syringes (<i>Humira, Yuflyma</i>)
12422T	ADALIMUMAB , adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL syringes (<i>Humira, Yuflyma</i>)
13215M	ADALIMUMAB , adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL pen devices (<i>Yuflyma</i>)
13217P	ADALIMUMAB , adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL syringes (<i>Yuflyma</i>)
12366W	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Amgevita, Hadlima, Hyrimoz, Idacio</i>)
12403T	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Amgevita, Hadlima, Hyrimoz, Idacio</i>)
9425C	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Amgevita, Hadlima, Hyrimoz, Idacio</i>)
9426D	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Amgevita, Hadlima, Hyrimoz, Idacio</i>)
9427E	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Amgevita, Hadlima, Hyrimoz, Idacio</i>)
9428F	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Amgevita, Hadlima, Hyrimoz, Idacio</i>)
12394H	ADALIMUMAB , adalimumab 80 mg/0.8 mL injection, 0.8 mL syringe (<i>Humira</i>)
12447D	ADALIMUMAB , adalimumab 80 mg/0.8 mL injection, 0.8 mL pen device (<i>Humira</i>)
13624C	AZACITIDINE , azacitidine 300 mg tablet, 7 (<i>Onureg</i>)
11223Q	ETANERCEPT , etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
9037P	ETANERCEPT , etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)

9429G	ETANERCEPT , etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
11221N	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys</i>)
11222P	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys, Enbrel</i>)
9461Y	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys, Enbrel</i>)
9462B	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys, Enbrel</i>)
11224R	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys, Enbrel</i>)
11225T	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys</i>)
9091L	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys, Enbrel</i>)
9431J	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys, Enbrel</i>)
11614G	GUSELKUMAB , guselkumab 100 mg/mL injection, 1 x 1 mL syringe (<i>Tremfya</i>)
13050W	INFLIXIMAB , infliximab 120 mg/mL injection, 1 mL syringe (<i>Remsima SC</i>)
13062L	INFLIXIMAB , infliximab 120 mg/mL injection, 1 mL pen device (<i>Remsima SC</i>)
13067R	INFLIXIMAB , infliximab 120 mg/mL injection, 1 mL syringe (<i>Remsima SC</i>)
13076F	INFLIXIMAB , infliximab 120 mg/mL injection, 1 mL pen device (<i>Remsima SC</i>)
11032P	IXEKIZUMAB , ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices (<i>Taltz</i>)
11033Q	IXEKIZUMAB , ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices (<i>Taltz</i>)
12969N	LEVOTHYROXINE , levothyroxine sodium 50 microgram tablet, 2 x 100 (<i>LEVOXINE, Thyrox</i>)
2174K	LEVOTHYROXINE , levothyroxine sodium 50 microgram tablet, 200 (<i>APO-Levothyroxine, Eltroxin, Eutroxsig, Levothox, Levothyroxine Lup, Oroxine</i>)
12960D	LEVOTHYROXINE , levothyroxine sodium 75 microgram tablet, 2 x 100 (<i>LEVOXINE, Thyrox</i>)
9287T	LEVOTHYROXINE , levothyroxine sodium 75 microgram tablet, 200 (<i>APO-Levothyroxine, Eltroxin, Eutroxsig, Levothox, Levothyroxine Lup, Oroxine</i>)
12968M	LEVOTHYROXINE , levothyroxine sodium 100 microgram tablet, 2 x 100 (<i>LEVOXINE, Thyrox</i>)
2175L	LEVOTHYROXINE , levothyroxine sodium 100 microgram tablet, 200 (<i>APO-Levothyroxine, Eltroxin, Eutroxsig, Levothox, Levothyroxine Lup, Oroxine</i>)
12970P	LEVOTHYROXINE , levothyroxine sodium 200 microgram tablet, 2 x 100 (<i>LEVOXINE, Thyrox</i>)
2173J	LEVOTHYROXINE , levothyroxine sodium 200 microgram tablet, 200 (<i>APO-Levothyroxine, Eltroxin, Eutroxsig, Levothox, Levothyroxine Lup, Oroxine</i>)
10731T	NETUPITANT + PALONOSETRON , netupitant 300 mg + palonosetron 500 microgram capsule, 1 (<i>Akynzeo</i>)
11827L	RISANKIZUMAB , risankizumab 75 mg/0.83 mL injection, 2 x 0.83 mL syringes (<i>Skyrizi</i>)
11858D	RISANKIZUMAB , risankizumab 75 mg/0.83 mL injection, 2 x 0.83 mL syringes (<i>Skyrizi</i>)
10425Q	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
10494H	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
10910F	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
11613F	TILDRAKIZUMAB , tildrakizumab 100 mg/mL injection, 1 mL syringe (<i>Ilumya</i>)
11616J	TILDRAKIZUMAB , tildrakizumab 100 mg/mL injection, 1 mL syringe (<i>Ilumya</i>)
9304Q	USTEKINUMAB , ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial (<i>Stelara</i>)
9305R	USTEKINUMAB , ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial (<i>Stelara</i>)

Alteration – Restriction

12342N	ADALIMUMAB , adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL pen devices (<i>Humira, Yuflyma</i>)
12421R	ADALIMUMAB , adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL syringes (<i>Humira, Yuflyma</i>)
9425C	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Amgevita, Hadlima, Hyrimoz, Idacio</i>)
9426D	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Amgevita, Hadlima, Hyrimoz, Idacio</i>)
12394H	ADALIMUMAB , adalimumab 80 mg/0.8 mL injection, 0.8 mL syringe (<i>Humira</i>)
12447D	ADALIMUMAB , adalimumab 80 mg/0.8 mL injection, 0.8 mL pen device (<i>Humira</i>)

12218C	APREMILAST , apremilast 10 mg tablet [4] (&) apremilast 20 mg tablet [4] (&) apremilast 30 mg tablet [19], 27 (<i>Otezla</i>)
12223H	APREMILAST , apremilast 30 mg tablet, 56 (<i>Otezla</i>)
9037P	ETANERCEPT , etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
9461Y	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys, Enbrel</i>)
9091L	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys, Enbrel</i>)
11614G	GUSELKUMAB , guselkumab 100 mg/mL injection, 1 x 1 mL syringe (<i>Tremfya</i>)
11032P	IXEKIZUMAB , ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices (<i>Taltz</i>)
10731T	NETUPITANT + PALONOSETRON , netupitant 300 mg + palonosetron 500 microgram capsule, 1 (<i>Akynzeo</i>)
11827L	RISANKIZUMAB , risankizumab 75 mg/0.83 mL injection, 2 x 0.83 mL syringes (<i>Skyrizi</i>)
10910F	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
11616J	TILDRAKIZUMAB , tildrakizumab 100 mg/mL injection, 1 mL syringe (<i>Ilumya</i>)
9304Q	USTEKINUMAB , ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial (<i>Stelara</i>)

Alteration – Manufacturer Code

		From	To
12142C	<i>Apomine Solution for Infusion</i> – APOMORPHINE , apomorphine hydrochloride hemihydrate 100 mg/20 mL injection, 5 x 20 mL vials	PF	IT
12133N	<i>Apomine Intermittent</i> – APOMORPHINE , apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL cartridges	PF	IT
5296R	<i>Riamet 20mg/120mg Dispersible</i> – ARTEMETHER + LUMEFANTRINE , artemether 20 mg + lumefantrine 120 mg dispersible tablet, 18	SZ	NV
9498X	<i>Riamet</i> – ARTEMETHER + LUMEFANTRINE , artemether 20 mg + lumefantrine 120 mg tablet, 24	SZ	NV
13457G	<i>CALITROL</i> – CALCITRIOL , calcitriol 0.25 microgram capsule, 100	ED	XT
2502Q	<i>CALITROL</i> – CALCITRIOL , calcitriol 0.25 microgram capsule, 100	ED	XT
1394J	<i>Eleanor 150/30 ED</i> – LEVONORGESTREL + ETHINYLESTRADIOL , levonorgestrel 150 microgram + ethinylestradiol 30 microgram tablet [21] (&) inert substance tablet [7], 4 x 28	EA	XT
1900B	<i>Clobemix</i> – MOCLOBEMIDE , moclobemide 150 mg tablet, 60	ED	XT
8003F	<i>Clobemix</i> – MOCLOBEMIDE , moclobemide 300 mg tablet, 60	ED	XT

Advance Notices

1 November 2023

Deletion – Brand

2324H	<i>Optifresh Plus, PP</i> – CARMELLOSE SODIUM , carmellose sodium 1% eye drops, 30 x 0.4 mL unit doses
2338C	<i>Optifresh Tears, PP</i> – CARMELLOSE SODIUM , carmellose sodium 0.5% eye drops, 30 x 0.4 mL unit doses
5505R	<i>Optifresh Plus, PP</i> – CARMELLOSE SODIUM , carmellose sodium 1% eye drops, 30 x 0.4 mL unit doses
5506T	<i>Optifresh Tears, PP</i> – CARMELLOSE SODIUM , carmellose sodium 0.5% eye drops, 30 x 0.4 mL unit doses
8452W	<i>Amaryl, SW</i> – GLIMEPIRIDE , glimepiride 4 mg tablet, 30
8814X	<i>CHEMISTS' OWN OSTEO RELIEF PARACETAMOL, RF</i> – PARACETAMOL , paracetamol 665 mg modified release tablet, 96
2363J	<i>Pregabalin GH, GQ</i> – PREGABALIN , pregabalin 300 mg capsule, 56

1 December 2023

Deletion – Brand

8596K	<i>Amisulpride 400 Winthrop, WA</i> – AMISULPRIDE , amisulpride 400 mg tablet, 60
2418G	<i>Amitriptyline Alphapharm 25, MQ</i> – AMITRIPTYLINE , amitriptyline hydrochloride 25 mg tablet, 50
9020R	<i>Moxonidine MYL, AF</i> – MOXONIDINE , moxonidine 400 microgram tablet, 30
8228C	<i>Ikorel, SW</i> – NICORANDIL , nicorandil 10 mg tablet, 60
8229D	<i>Ikorel, SW</i> – NICORANDIL , nicorandil 20 mg tablet, 60

8411Q Zofran Zydys, AS – **ONDANSETRON**, ondansetron 8 mg wafer, 4

1 January 2024

Deletion – Brand

10062N Temodal, MK – **TEMOZOLOMIDE**, temozolomide 180 mg capsule, 5

2438H Temodal, MK – **TEMOZOLOMIDE**, temozolomide 180 mg capsule, 5

8381D Temodal, MK – **TEMOZOLOMIDE**, temozolomide 250 mg capsule, 5

Palliative Care

Advance Notices

1 November 2023

Deletion – Brand

5343F CHEMISTS' OWN OSTEO RELIEF PARACETAMOL, RF – **PARACETAMOL**, paracetamol 665 mg modified release tablet, 96

Highly Specialised Drugs Program (Private Hospital)

Additions

Addition – Item

13642B **LENALIDOMIDE**, lenalidomide 5 mg capsule, 21 (*Cipla Lenalidomide, Lenalide, Lenalidomide Dr.Reddy's, Lenalidomide Sandoz, Lenalidomide Viatris, Lenalidomide-Teva, Revlimid*)

13658W **LENALIDOMIDE**, lenalidomide 10 mg capsule, 21 (*Cipla Lenalidomide, Lenalide, Lenalidomide Dr.Reddy's, Lenalidomide Sandoz, Lenalidomide Viatris, Lenalidomide-Teva, Revlimid*)

13657T **LENALIDOMIDE**, lenalidomide 15 mg capsule, 21 (*Cipla Lenalidomide, Lenalide, Lenalidomide Dr.Reddy's, Lenalidomide Sandoz, Lenalidomide Viatris, Lenalidomide-Teva, Revlimid*)

13660Y **LENALIDOMIDE**, lenalidomide 25 mg capsule, 21 (*Cipla Lenalidomide, Lenalide, Lenalidomide Dr.Reddy's, Lenalidomide Sandoz, Lenalidomide Viatris, Lenalidomide-Teva, Revlimid*)

13632L **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)

13639W **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)

13646F **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)

13647G **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)

13659X **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)

Addition – Brand

12784W Azacitidine Dr.Reddy's, RI – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial

13033Y Azacitidine Dr.Reddy's, RI – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial

13038F Azacitidine Dr.Reddy's, RI – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial

13039G Azacitidine Dr.Reddy's, RI – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial

13040H Azacitidine Dr.Reddy's, RI – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial

6100C Azacitidine Dr.Reddy's, RI – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial

6138C Azacitidine Dr.Reddy's, RI – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial

Addition – Restriction

13064N **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)

13111C **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)

Deletions

Deletion – Item

11420C **BACLOFEN**, baclofen 10 mg/5 mL intrathecal injection, 5 x 5 mL ampoules (*Bacetheal*)

Deletion – Brand

6227R Octreotide MaxRx, GQ – **OCTREOTIDE**, octreotide 50 microgram/mL injection, 5 x 1 mL ampoules

Alterations

Alteration – Note

11128Q	BACLOFEN , baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules (<i>Sintetica Baclofen Intrathecal</i>)
6284R	BACLOFEN , baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule (<i>Bacthecal, Lioresal Intrathecal</i>)
11590B	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Remicade, Renflexis</i>)
11595G	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Renflexis</i>)
9617E	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Remicade, Renflexis</i>)
13045N	NUSINERSEN , nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (<i>Spinraza</i>)
13064N	NUSINERSEN , nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (<i>Spinraza</i>)
13091B	NUSINERSEN , nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (<i>Spinraza</i>)
13111C	NUSINERSEN , nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (<i>Spinraza</i>)
12609P	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)
12610Q	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)
12959C	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)

Alteration – Restriction

9617E	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Remicade, Renflexis</i>)
13045N	NUSINERSEN , nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (<i>Spinraza</i>)
13091B	NUSINERSEN , nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (<i>Spinraza</i>)
12609P	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)
12610Q	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)
12959C	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)

Alteration – Manufacturer Code

		From	To
11083H	<i>Apomine Solution for Infusion</i> – APOMORPHINE , apomorphine hydrochloride hemihydrate 100 mg/20 mL injection, 5 x 20 mL vials	PF	IT
11688E	<i>Apomine Intermittent</i> – APOMORPHINE , apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL cartridges	PF	IT

Advance Notices

1 November 2023

Deletion – Brand

6357N	<i>Valganciclovir Mylan, AF</i> – VALGANCICLOVIR , valganciclovir 450 mg tablet, 60
-------	--

1 December 2023

Deletion – Brand

6126K	<i>Neupogen, AN</i> – FILGRASTIM , filgrastim 300 microgram/mL injection, 10 x 1 mL vials
6127L	<i>Neupogen, AN</i> – FILGRASTIM , filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials
6291D	<i>Neupogen, AN</i> – FILGRASTIM , filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes
6292E	<i>Neupogen, AN</i> – FILGRASTIM , filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes
6229W	<i>Octreotide MaxRx, GQ</i> – OCTREOTIDE , octreotide 500 microgram/mL injection, 5 x 1 mL ampoules

Highly Specialised Drugs Program (Public Hospital)

Additions

Addition – Item

13636Q	LENALIDOMIDE , lenalidomide 5 mg capsule, 21 (<i>Cipla Lenalidomide, Lenalide, Lenalidomide Dr.Reddy's, Lenalidomide Sandoz, Lenalidomide Viatris, Lenalidomide-Teva, Revlimid</i>)
13661B	LENALIDOMIDE , lenalidomide 10 mg capsule, 21 (<i>Cipla Lenalidomide, Lenalide, Lenalidomide Dr.Reddy's, Lenalidomide Sandoz, Lenalidomide Viatris, Lenalidomide-Teva, Revlimid</i>)
13641Y	LENALIDOMIDE , lenalidomide 15 mg capsule, 21 (<i>Cipla Lenalidomide, Lenalide, Lenalidomide Dr.Reddy's, Lenalidomide Sandoz, Lenalidomide Viatris, Lenalidomide-Teva, Revlimid</i>)

13630J	LENALIDOMIDE , lenalidomide 25 mg capsule, 21 (<i>Cipla Lenalidomide, Lenalide, Lenalidomide Dr.Reddy's, Lenalidomide Sandoz, Lenalidomide Viatris, Lenalidomide-Teva, Revlimid</i>)
13662C	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [7 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13666G	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [8 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13670L	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [3 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13671M	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13672N	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [4 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13677W	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [6 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13680B	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13663D	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [8.3 mL vial], 1 pack (<i>Zolgensma</i>)
13668J	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [7 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13669K	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13674Q	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 2 x 8.3 mL vials (<i>Zolgensma</i>)
13679Y	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [6 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13681C	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13682D	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [3 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13683E	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [4 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
13678X	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 3 x 8.3 mL vials (<i>Zolgensma</i>)
13665F	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 4 x 8.3 mL vials (<i>Zolgensma</i>)
13667H	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 5 x 8.3 mL vials (<i>Zolgensma</i>)
13664E	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 6 x 8.3 mL vials (<i>Zolgensma</i>)
13675R	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 7 x 8.3 mL vials (<i>Zolgensma</i>)
13673P	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 8 x 8.3 mL vials (<i>Zolgensma</i>)
13676T	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 9 x 8.3 mL vials (<i>Zolgensma</i>)
13631K	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)
13633M	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)

13654P **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)
 13655Q **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)
 13656R **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)

Addition – Brand

12771E *Azacitidine Dr.Reddy's, RI* – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial
 13028Q *Azacitidine Dr.Reddy's, RI* – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial
 13036D *Azacitidine Dr.Reddy's, RI* – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial
 13042K *Azacitidine Dr.Reddy's, RI* – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial
 13044M *Azacitidine Dr.Reddy's, RI* – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial
 9597D *Azacitidine Dr.Reddy's, RI* – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial
 9598E *Azacitidine Dr.Reddy's, RI* – **AZACITIDINE**, azacitidine 100 mg injection, 1 vial

Addition – Restriction

13052Y **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)
 13083N **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)

Deletions

Deletion – Item

11421D **BACLOFEN**, baclofen 10 mg/5 mL intrathecal injection, 5 x 5 mL ampoules (*Bacethecal*)

Deletion – Brand

9508K *Octreotide MaxRx, GQ* – **OCTREOTIDE**, octreotide 50 microgram/mL injection, 5 x 1 mL ampoules

Alterations

Alteration – Note

11126N **BACLOFEN**, baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules (*Sintetica Baclofen Intrathecal*)
 5617P **BACLOFEN**, baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule (*Bacethecal, Lioresal Intrathecal*)
 11605T **INFLIXIMAB**, infliximab 100 mg injection, 1 vial (*Inflixtra, Renflexis*)
 11606W **INFLIXIMAB**, infliximab 100 mg injection, 1 vial (*Inflixtra, Remicade, Renflexis*)
 5758C **INFLIXIMAB**, infliximab 100 mg injection, 1 vial (*Inflixtra, Remicade, Renflexis*)
 13052Y **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)
 13068T **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)
 13083N **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)
 13084P **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)
 12606L **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)
 12614X **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)
 12943F **RISDIPLAM**, risdiplam 750 microgram/mL powder for oral liquid, 80 mL (*Evrysdi*)

Alteration – Restriction

5758C **INFLIXIMAB**, infliximab 100 mg injection, 1 vial (*Inflixtra, Remicade, Renflexis*)
 13068T **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)
 13084P **NUSINERSEN**, nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial (*Spinraza*)
 12942E **ONASEMNOGENE ABEPARVOVEC**, onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [4 x 8.3 mL vials], 1 pack (*Zolgensma*)
 12951P **ONASEMNOGENE ABEPARVOVEC**, onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 8.3 mL vials], 1 pack (*Zolgensma*)
 12956X **ONASEMNOGENE ABEPARVOVEC**, onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [7 x 8.3 mL vials], 1 pack (*Zolgensma*)
 12964H **ONASEMNOGENE ABEPARVOVEC**, onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [8 x 8.3 mL vials], 1 pack (*Zolgensma*)

12975X	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [6 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
12976Y	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
12977B	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [3 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
12940C	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [3 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
12941D	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [6 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
12950N	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [4 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
12952Q	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [7 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
12955W	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [8.3 mL vial], 1 pack (<i>Zolgensma</i>)
12957Y	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 2 x 8.3 mL vials (<i>Zolgensma</i>)
12958B	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
12963G	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5 x 8.3 mL vials], 1 pack (<i>Zolgensma</i>)
12961E	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 3 x 8.3 mL vials (<i>Zolgensma</i>)
12971Q	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 4 x 8.3 mL vials (<i>Zolgensma</i>)
12962F	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 5 x 8.3 mL vials (<i>Zolgensma</i>)
12973T	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 6 x 8.3 mL vials (<i>Zolgensma</i>)
12954T	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 7 x 8.3 mL vials (<i>Zolgensma</i>)
12974W	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 8 x 8.3 mL vials (<i>Zolgensma</i>)
12949M	ONASEMNOGENE ABEPARVOVEC , onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 9 x 8.3 mL vials (<i>Zolgensma</i>)
12606L	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)
12614X	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)
12943F	RISDIPLAM , risdiplam 750 microgram/mL powder for oral liquid, 80 mL (<i>Evrysdi</i>)

Alteration – Manufacturer Code

		<i>From</i>	<i>To</i>
11093W	<i>Apomine Solution for Infusion</i> – APOMORPHINE , apomorphine hydrochloride hemihydrate 100 mg/20 mL injection, 5 x 20 mL vials	PF	IT
11672H	<i>Apomine Intermittent</i> – APOMORPHINE , apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL cartridges	PF	IT

Advance Notices

1 November 2023

Deletion – Brand

9569P Valganciclovir Mylan, AF – **VALGANCICLOVIR**, valganciclovir 450 mg tablet, 60

1 December 2023

Deletion – Brand

5741E Neupogen, AN – **FILGRASTIM**, filgrastim 300 microgram/mL injection, 10 x 1 mL vials

5742F Neupogen, AN – **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes

5743G Neupogen, AN – **FILGRASTIM**, filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials

5744H Neupogen, AN – **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes

9510M Octreotide MaxRx, GQ – **OCTREOTIDE**, octreotide 500 microgram/mL injection, 5 x 1 mL ampoules

Highly Specialised Drugs Program (Community Access)

Additions

Addition – Brand

10279B Entecavir Viatris, AL – **ENTECAVIR**, entecavir 500 microgram tablet, 30

Advance Notices

1 November 2023

Deletion – Brand

10306K Valganciclovir Mylan, AF – **VALGANCICLOVIR**, valganciclovir 450 mg tablet, 60

Growth Hormone Program

Alterations

Alteration – Restriction

11493X **SOMATROPIN**, somatropin 5 mg injection [1 chamber] (&) inert substance diluent [1 mL chamber], 1 dual chamber pen device (*Genotropin GoQuick*)

11495B **SOMATROPIN**, somatropin 12 mg injection [1 chamber] (&) inert substance diluent [1 mL chamber], 1 dual chamber pen device (*Genotropin GoQuick*)

11895C **SOMATROPIN**, somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge (*Norditropin FlexPro*)

11650E **SOMATROPIN**, somatropin 10 mg/2 mL injection, 2 mL cartridge (*NutropinAq*)

Repatriation Pharmaceutical Benefits

Deletions

Deletion – Brand

2194L **ALENDRONATE PLUS D3 70mg/70ug APOTEX, GX – ALENDRONATE + COLECALCIFEROL**, alendronate 70 mg + colecalciferol 70 microgram (2800 units) tablet, 4

2224C **ALENDRONATE PLUS D3 70mg/140ug APOTEX, GX – ALENDRONATE + COLECALCIFEROL**, alendronate 70 mg + colecalciferol 140 microgram (5600 units) tablet, 4

Advance Notices

1 November 2023

Deletion – Brand

10598T **CHEMISTS' OWN OSTEO RELIEF PARACETAMOL, RF – PARACETAMOL**, paracetamol 665 mg modified release tablet, 96

General Pharmaceutical Benefits

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after

at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Pharmaceutical benefits that have the form adalimumab 40 mg/0.4 mL syringes and pharmaceutical benefits that have the form adalimumab 40 mg/0.8 mL syringes are equivalent for the purposes of substitution

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

11635

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required (STREAMLINED)

11606

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to their most recent course of treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL syringes

13217P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	676.79	30.00	^a Yuflyma [EW]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

12366W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	827.02	30.00	^a Amgevita [XT] ^a Hyrimoz [SZ]	^a Hadlima [RF] ^a Idacio [PK]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the

new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and

scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Pharmaceutical benefits that have the form adalimumab 40 mg/0.4 mL pen devices and pharmaceutical benefits that have the form adalimumab 40 mg/0.8 mL pen devices are equivalent for the purposes of substitution

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

11635

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required (STREAMLINED)

11606

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to their most recent course of treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL pen devices

13215M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	676.79	30.00	^a Yuflyma [EW]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

12403T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	827.02	30.00	^a Amgevita [XT] ^a Hyrimoz [SZ]	^a Hadlima [RF] ^a Idacio [PK]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion

of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Pharmaceutical benefits that have the form adalimumab 40 mg/0.4 mL pen devices and pharmaceutical benefits that have the form adalimumab 40 mg/0.8 mL pen devices are equivalent for the purposes of substitution

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheet and the face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR

- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheet and the face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL pen devices

12377K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	676.79	30.00	^a Humira [VE]	^a Yuflyma [EW]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9428F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	827.02	30.00	^a Amgevita [XT] ^a Hyrimoz [SZ]	^a Hadlima [RF] ^a Idacio [PK]

■ ADALIMUMAB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a

treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Pharmaceutical benefits that have the form adalimumab 40 mg/0.4 mL syringes and pharmaceutical benefits that have the form adalimumab 40 mg/0.8 mL syringes are equivalent for the purposes of substitution

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheet and the face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**

- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**

- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheet and the face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL syringes

12422T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	676.79	30.00	^a Humira [VE]	^a Yuflyma [EW]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9427E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	827.02	30.00	^a Amgevita [XT] ^a Hyrimoz [SZ]	^a Hadlima [RF] ^a Idacio [PK]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in

biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

An assessment of a patient's response to this initial course of treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks prior the completion of this course of treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

An assessment of a patient's response to this initial course of treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks prior the completion of this course of treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

adalimumab 80 mg/0.8 mL injection, 0.8 mL pen device

12447D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	676.79	30.00	Humira [VE]

adalimumab 80 mg/0.8 mL injection, 0.8 mL syringe

12394H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	676.79	30.00	Humira [VE]

■ ADALIMUMAB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Amgevita, Hadlima, Hyrimoz, Idacio or Yuflyma is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Pharmaceutical benefits that have the form adalimumab 40 mg/0.4 mL pen devices and pharmaceutical benefits that have the form adalimumab 40 mg/0.8 mL pen devices are equivalent for the purposes of substitution

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

An assessment of a patient's response to this initial course of treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks prior the completion of this course of treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

An assessment of a patient's response to this initial course of treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks prior the completion of this course of treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL pen devices

12342N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	4	..	676.79	30.00	^a Humira [VE]	^a Yuflyma [EW]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9426D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	4	..	827.02	30.00	^a Amgevita [XT] ^a Hyrimoz [SZ]	^a Hadlima [RF] ^a Idacio [PK]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in

biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Amgevita, Hadlima, Hyrimoz, Idacio or Yuflyma is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Pharmaceutical benefits that have the form adalimumab 40 mg/0.4 mL syringes and pharmaceutical benefits that have the form adalimumab 40 mg/0.8 mL syringes are equivalent for the purposes of substitution

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

An assessment of a patient's response to this initial course of treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks prior the completion of this course of treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).**Note** Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

An assessment of a patient's response to this initial course of treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks prior the completion of this course of treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:

(i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Prescribing the 80 mg presentation:

Two completed authority prescriptions may be submitted with every initial application for this biological medicine. One prescription for an induction dose of 80 mg and no repeats and the second prescription for 2 doses of 40 mg and 3 repeats.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.4 mL injection, 2 x 0.4 mL syringes

12421R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	4	..	676.79	30.00	^a Humira [VE]	^a Yuflyma [EW]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9425C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	4	..	827.02	30.00	^a Amgevita [XT] ^a Hyrimoz [SZ]	^a Hadlima [RF] ^a Idacio [PK]

■ AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES

Note Alfamino products that contain 2-fucosyllactose and lacto-N-neotetrose and Alfamino products that do not contain 2-fucosyllactose and lacto-N-neotetrose are equivalent for the purposes of substitution.

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have been assessed at least once or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have had failure to thrive prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist at intervals not greater than 12 months.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Cows' milk anaphylaxis

Treatment criteria:

- Must be treated by a specialist allergist or clinical immunologist, or in consultation with a specialist allergist or clinical immunologist.

Population criteria:

- Patient must be up to the age of 24 months.

Anaphylaxis is defined as a severe and/or potentially life threatening allergic reaction.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides) prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe intestinal malabsorption including short bowel syndrome

Clinical criteria:

- Patient must have failed to respond to protein hydrolysate formulae; OR
- Patient must have been receiving parenteral nutrition.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must have responded to an initial course of PBS-subsidised treatment.

Population criteria:

- Patient must be 18 years of age or less.

Response to initial treatment is demonstrated by oesophageal biopsy specimens obtained by endoscopy, where the most densely involved oesophageal biopsy had 5 or less eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies. The response criteria will not be deemed to have been met if oral steroids were commenced during initial treatment.

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides powder for oral liquid, 400 g

2900P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*316.37	30.00	^a Alfamino [NT]

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides with 2-fucosyllactose and lacto-N-neotetraose powder for oral liquid, 400 g

13615N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*316.37	30.00	^a Alfamino [NT]

■ AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES

Note Alfamino products that contain 2-fucosyllactose and lacto-N-neoteroose and Alfamino products that do not contain 2-fucosyllactose and lacto-N-neoteroose are equivalent for the purposes of substitution.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides).

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Initial treatment for up to 3 months

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must require an amino acid based formula as a component of a dietary elimination program.

Population criteria:

- Patient must be 18 years of age or less.

Treatment with oral steroids should not be commenced during the period of initial treatment.

Eosinophilic oesophagitis is demonstrated by the following criteria:

- (i) Chronic symptoms of reflux that persisted despite a 2-month trial of a proton pump inhibitor or chronic dysphagia; and
- (ii) A lack of demonstrable anatomic abnormality with the exception of stricture, which can be attributable to eosinophilic oesophagitis; and
- (iii) Eosinophilic infiltration of the oesophagus, demonstrated by oesophageal biopsy specimens obtained by endoscopy and where the most densely involved oesophageal biopsy had 20 or more eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies.

The date of birth of the patient must be included in the authority application.

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides powder for oral liquid, 400 g

2928D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*316.37	30.00	^a Alfamino [NT]

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides with 2-fucosyllactose and lacto-N-neotetraose powder for oral liquid, 400 g

13627F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*316.37	30.00	^a Alfamino [NT]

■ APREMILAST

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14417

Severe chronic plaque psoriasis

Clinical criteria:

- Patient must not have achieved adequate response after at least 6 weeks of treatment with methotrexate prior to initiating treatment with this drug; OR
- Patient must have a contraindication to methotrexate according to the Therapeutic Goods Administration (TGA) approved Product Information; OR
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate, **AND**
- The condition must have caused significant interference with quality of life, **AND**
- Patient must not be undergoing concurrent PBS-subsidised treatment for psoriasis with each of: (i) a biological medicine, (ii) ciclosporin, (iii) deucravacitinib.

Treatment criteria:

- Must be treated by a medical practitioner who is either: (i) a dermatologist, (ii) an accredited dermatology registrar in consultation with a dermatologist; OR
- Must be treated by a general practitioner who has been directed to continue treatment (not initiate treatment) by one of the above practitioner types.

Population criteria:

- Patient must be at least 18 years of age.

apremilast 30 mg tablet, 56

12223H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	653.71	30.00	Otezla [AN]

apremilast 10 mg tablet [4] (&) apremilast 20 mg tablet [4] (&) apremilast 30 mg tablet [19], 27

12218C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	273.20	30.00	Otezla [AN]

■ AZACITIDINE

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 6 months following cessation of therapy.

Oral azacitidine should not be used interchangeably with injectable azacitidine due to differences in the exposure, dose and schedule of treatment.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Dose escalation therapy - Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have, in order to extend the dose schedule as per the TGA-approved Product Information, between 5% to 15% blasts in either the: (i) bone marrow, (ii) peripheral blood, in conjunction with clinical assessment, **AND**
- Patient must not be receiving concomitant PBS-subsidised treatment with midostaurin.

Authority applications must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail:

If the application is submitted through HPOS form upload or mail, it must include:

- (a) a completed authority prescription form; and
- (b) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice)
- (c) details (date, unique identifying number/code or provider number) of the pathology report from an Approved Pathology Authority demonstrating the blast percentage.

All reports must be documented in the patient's medical records.

azacitidine 300 mg tablet, 7

13624C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	1	..	*20003.79	30.00	Onureg [CJ]

■ BIMEKIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**

- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the following:

(i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice) which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets, and the face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 24 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 24 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

bimekizumab 160 mg/mL injection, 2 x 1 mL pen devices

13644D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	4	..	3422.13	30.00	Bimzelx [UC]

▪ **BIMEKIZUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised

therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years). An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Grandfathered patient - Whole body (initial PBS-subsidised supply for continuing treatment in a patient commenced on non-PBS-subsidised therapy)

Clinical criteria:

- Patient must have a documented severe chronic plaque psoriasis where lesions have been present for at least 6 months prior to commencing non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have received non-PBS-subsidised treatment with this drug for this PBS indication prior to 1 October 2023, **AND**
- Patient must have a documented failure to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments prior to commencing non-PBS-subsidised treatment with this drug for this condition: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- Patient must have a documented Psoriasis Area and Severity Index (PASI) score of greater than 15 prior to commencing non-PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheets including the date of the assessment of the patient's condition at baseline (prior to initiation of therapy with this drug); and
- (c) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Note A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the continuing treatment criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Grandfathered patient - Face, hand, foot (initial PBS-subsidised supply for continuing treatment in a patient commenced on non-PBS-subsidised therapy)

Clinical criteria:

- Patient must have a documented severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where lesions have been present for at least 6 months prior to commencing non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have received non-PBS-subsidised treatment with this drug for this PBS indication prior to 1 October 2023, **AND**
- Patient must have a documented failure to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments prior to commencing non-PBS-subsidised treatment with this drug for this condition: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- Patient must have a documented Psoriasis Area and Severity Index (PASI) score of greater than 15 prior to commencing non-PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheets including the date of the assessment of the patient's condition at baseline (prior to initiation of therapy with this drug); and
- (c) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Note A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the continuing treatment criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Grandfathered patient - Face, hand, foot or Whole body - Balance of Supply

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must have received insufficient therapy with this drug for this condition under the Grandfathered patient - Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Grandfathered patient - Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

bimekizumab 160 mg/mL injection, 2 x 1 mL pen devices

13652M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3422.13	30.00	Bimzelx [UC]

■ CEFUROXIME**cefuroxime 125 mg/5 mL powder for oral liquid, 70 mL**

13643C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±2	1	..	*#115.51	30.00	Zinnat (UK) [RQ]

cefuroxime 125 mg/5 mL powder for oral liquid, 70 mL

13653N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±2	..	..	*#115.51	30.00	Zinnat (UK) [RQ]

DP

■ DEUCRAVACITINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**14384**

Severe chronic plaque psoriasis

Clinical criteria:

- Patient must not have achieved adequate response after at least 6 weeks of treatment with methotrexate prior to initiating treatment with this drug; OR
- Patient must have a contraindication to methotrexate according to the Therapeutic Goods Administration (TGA) approved Product Information; OR
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate, **AND**
- The condition must have caused significant interference with quality of life, **AND**
- Patient must not be undergoing concurrent PBS-subsidised treatment for psoriasis with each of: (i) a biological medicine, (ii) ciclosporin, (iii) apremilast.

Treatment criteria:

- Must be treated by a medical practitioner who is either: (i) a dermatologist, (ii) an accredited dermatology registrar in consultation with a dermatologist; OR
- Must be treated by a general practitioner who has been directed to continue treatment (not initiate treatment) by one of the above practitioner types.

Population criteria:

- Patient must be at least 18 years of age.

deucravacitinib 6 mg tablet, 28

13649J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1259.52	30.00	Sotyktu [BQ]

■ ETANERCEPT**Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological

medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that

they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Pharmaceutical benefits that have the form etanercept injection 50 mg/mL, 4 x 1 mL syringes and pharmaceutical benefits that have the form etanercept injection 50 mg/mL, 4 x 1 mL pen devices are equivalent for the purposes of substitution.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

8887

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required (STREAMLINED)

8955

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to their most recent course of treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11221N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	845.62	30.00	^a Brenzys [RF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11225T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	845.62	30.00	^a Brenzys [RF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

11223Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*845.63	30.00	Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence

such therapy (Initial 1 - New patient); or
(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**

- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

9429G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*845.63	30.00	Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was

approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and

scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Pharmaceutical benefits that have the form etanercept injection 50 mg/mL, 4 x 1 mL syringes and pharmaceutical benefits that have the form etanercept injection 50 mg/mL, 4 x 1 mL pen devices are equivalent for the purposes of substitution.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11222P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	845.62	30.00	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11224R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	845.62	30.00	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting

in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Pharmaceutical benefits that have the form etanercept injection 50 mg/mL, 4 x 1 mL syringes and pharmaceutical benefits that have the form etanercept injection 50 mg/mL, 4 x 1 mL pen devices are equivalent for the purposes of substitution.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9462B	Max. Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	845.62	30.00	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9431J	Max. Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	845.62	30.00	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT**Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

-
- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
- (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
- (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to Services Australia no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.
- The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to Services Australia no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and
(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack

9037P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*845.63	30.00	Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide

new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Pharmaceutical benefits that have the form etanercept injection 50 mg/mL, 4 x 1 mL syringes and pharmaceutical benefits that have the form etanercept injection 50 mg/mL, 4 x 1 mL pen devices are equivalent for the purposes of substitution.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to Services Australia no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to Services Australia no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available

on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online

Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9461Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	845.62	30.00	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9091L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	845.62	30.00	^a Brenzys [RF]	^a Enbrel [PF]

▪ **FOSNETUPITANT + PALONOSETRON**

Note This medicine is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Note Various sources of information outline the emetic risk associated with cancer treatment. Examples include the National Comprehensive Cancer Network guidelines (USA), eviQ guidelines and approved Product Information of individual drugs. These examples are not a comprehensive list of which anti-cancer drugs that have moderate to high emesis risk.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Nausea and vomiting

Clinical criteria:

- The treatment must be for prevention of nausea and vomiting associated with moderate to highly emetogenic anti-cancer therapy, **AND**
- The treatment must be in combination with dexamethasone, unless contraindicated, **AND**
- Patient must be unable to swallow; OR
- Patient must be contraindicated to oral anti-emetics.

fosnetupitant 235 mg + palonosetron 250 microgram injection, 1 vial

13640X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	111.12	30.00	Akynzeo IV [JZ]

▪ **GUSELKUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological

medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline

PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) **Recommencement of treatment after a 5-year break in PBS-subsidised therapy.**

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and
(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 20 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 20 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

guselkumab 100 mg/mL injection, 1 x 1 mL syringe

11614G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3796.75	30.00	Tremfya [JC]

▪ **INFLIXIMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4

weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Where there is already an approved authority prescription for the IV formulation, an authority application for this subcutaneously administered form can be made under one of:

(1) The Balance of Supply listing - apply under this listing if the treatment phase (Initial treatment, First Continuing treatment, Subsequent Continuing treatment) is yet to be completed; the following authority application is to be under a Continuing treatment listing.

(2) A Continuing treatment listing - apply under one of these types of listings where a treatment phase is concluding/has concluded; this typically occurs for the dose due at treatment week 16, 40, 64, 88 etc.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note The application should indicate which formulation eg. pre-filled pen or pre-filled syringe to ensure appropriate item is approved.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment (whole body, or, face/hand/foot) with subcutaneous form or switching from intravenous form to subcutaneous form

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- The treatment must have both: (i) provided the patient with an adequate response with the preceding supply, (ii) been assessed for response after at least 12 weeks of therapy, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Where the condition is affecting the whole body, an adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by at least 75%, or, is sustained at this level, when compared with the baseline value for this treatment cycle. State the qualifying PASI score in the authority application.

Where the condition is affecting the face/hand/foot, an adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) A reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or, sustained at this level, as compared to the baseline values. Indicate the rating (0=none, 1=slight) for each of these 3 observations in the authority application for each affected area; or

(ii) A reduction by at least 75% in the skin area affected, or, sustained at this level, as compared to the baseline value for this treatment cycle. State the qualifying numerical percentage figure in the authority application for each affected area.

All assessment findings must be no more than 1 month old at the time of application. Response assessments must be performed on the same affected area assessed at baseline.

infliximab 120 mg/mL injection, 1 mL pen device

13076F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*577.89	30.00	Remsima SC [EW]

infliximab 120 mg/mL injection, 1 mL syringe

13050W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*759.43	30.00	Remsima SC [EW]

▪ **INFLIXIMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Where there is already an approved authority prescription for the IV formulation, an authority application for this subcutaneously administered form can be made under one of:

- (1) The Balance of Supply listing - apply under this listing if the treatment phase (Initial treatment, First Continuing treatment, Subsequent Continuing treatment) is yet to be completed; the following authority application is to be under a Continuing treatment listing.
- (2) A Continuing treatment listing - apply under one of these types of listings where a treatment phase is concluding/has concluded; this typically occurs for the dose due at treatment week 16, 40, 64, 88 etc.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note The application should indicate which formulation eg. pre-filled pen or pre-filled syringe to ensure appropriate item is approved.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment with the subcutaneous form where a concurrent PBS authority application for the intravenously (IV) administered formulation is being made

Treatment criteria:

- Must be treated by a specialist prescriber who is the same prescriber completing the PBS authority application for the IV administered formulation of this drug/biological medicine, **AND**
- Patient must be undergoing treatment with this benefit where: (i) there is a concurrent PBS authority application for the IV administered formulation submitted for approval, (ii) the concurrent PBS authority application is approved/in the process of being approved.

Population criteria:

- Patient must be at least 18 years of age.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

The PBS administrator will confirm that:

- (i) there is a concurrent authority application for the intravenous (IV) formulation of this benefit for the patient;
- (ii) the concurrent authority application for the IV formulation is to be approved before approving this authority application.

infliximab 120 mg/mL injection, 1 mL pen device

13062L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*577.89	30.00	Remsima SC [EW]

infliximab 120 mg/mL injection, 1 mL syringe

13067R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*759.43	30.00	Remsima SC [EW]

■ INSULIN ASPART

insulin aspart 100 units/mL fast acting injection, 5 x 3 mL cartridges

13651L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	1	..	*167.07	30.00	Fiasp Penfill [NO]

NP

■ IXEKIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological

medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline

PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**

- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices

11033Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3259.13	30.00	Taltz [LY]

▪ **IXEKIZUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).
- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed

treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

- (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
- (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available

on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

- (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

- (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of restriction after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and

- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks

from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices

11032P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	3259.13	30.00	Taltz [LY]

■ LEVOTHYROXINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Eltroxin and Levothox are not interchangeable with Oroxine, Eutroxsig, Levoxine, Levothyroxine Lup, APO-Levothyroxine or Thyrox on a dose to dose basis, and dose titration may be required.

Note Pharmaceutical benefits that have the form levothyroxine sodium 75 microgram tablet, 200 (Eutroxsig, Oroxine, Levothyroxine Lup and APO-Levothyroxine brands) and pharmaceutical benefits that have the form levothyroxine sodium 75 microgram tablet, 2 x 100 (Levoxine and Thyrox brands) are equivalent for the purposes of substitution.

levothyroxine sodium 75 microgram tablet, 2 x 100

12960D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	23.26	24.66	^a LEVOXINE [RA]	^a Thyrox [IX]

■ LEVOTHYROXINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Eltroxin and Levothox are not interchangeable with Oroxine, Eutroxsig, Levoxine, Levothyroxine Lup, APO-Levothyroxine or Thyrox on a dose to dose basis, and dose titration may be required.

Note Pharmaceutical benefits that have the form levothyroxine sodium 100 microgram tablet, 200 (Eutroxsig, Oroxine, Levothyroxine Lup and APO-Levothyroxine brands) and pharmaceutical benefits that have the form levothyroxine sodium 100 microgram tablet, 2 x 100 (Levoxine and Thyrox brands) are equivalent for the purposes of substitution.

levothyroxine sodium 100 microgram tablet, 2 x 100

12968M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	23.45	24.85	^a LEVOXINE [RA]	^a Thyrox [IX]

■ LEVOTHYROXINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Eltroxin and Levothox are not interchangeable with Oroxine, Eutroxsig, Levoxine, Levothyroxine Lup, APO-Levothyroxine or Thyrox on a dose to dose basis, and dose titration may be required.

Note Pharmaceutical benefits that have the form levothyroxine sodium 50 microgram tablet, 200 (Eutroxsig, Oroxine, Levothyroxine Lup and APO-Levothyroxine brands) and pharmaceutical benefits that have the form levothyroxine sodium 50 microgram tablet, 2 x 100 (Levoxine and Thyrox brands) are equivalent for the purposes of substitution.

levothyroxine sodium 50 microgram tablet, 2 x 100

12969N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	23.09	24.49	^a LEVOXINE [RA]	^a Thyrox [IX]

■ LEVOTHYROXINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Eltroxin and Levothox are not interchangeable with Oroxine, Eutroxsig, Levoxine, Levothyroxine Lup, APO-Levothyroxine or Thyrox on a dose to dose basis, and dose titration may be required.

Note Pharmaceutical benefits that have the form levothyroxine sodium 200 microgram tablet, 200 (Eutroxsig, Oroxine, Levothyroxine Lup and APO-Levothyroxine brands) and pharmaceutical benefits that have the form levothyroxine sodium 200 microgram tablet, 2 x 100 (Levoxine and Thyrox brands) are equivalent for the purposes of substitution.

levothyroxine sodium 200 microgram tablet, 2 x 100

12970P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	25.26	26.66	^a LEVOXINE [RA]	^a Thyrox [IX]

■ LEVOTHYROXINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Eltroxin and Levothox are not interchangeable with Oroxine, Eutroxsig, Levoxine, Levothyroxine Lup, APO-Levothyroxine or Thyrox on a dose to dose basis, and dose titration may be required.

Note Pharmaceutical benefits that have the form levothyroxine sodium 200 microgram tablet, 200 (Eltroxin and Levothox brands) are equivalent for the purposes of substitution.

Note Pharmaceutical benefits that have the form levothyroxine sodium 200 microgram tablet, 200 (Eutroxsig, Oroxine, Levothyroxine Lup and APO-Levothyroxine brands) and pharmaceutical benefits that have the form levothyroxine sodium 200 microgram tablet, 2 x 100 (Levoxine and Thyrox brands) are equivalent for the purposes of substitution.

levothyroxine sodium 200 microgram tablet, 200

2173J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	25.26	26.66	^a APO-Levothyroxine [XT] ^a Levothyroxine Lup [GQ] ^b Eltroxin [LT]	^a Eutroxsig [LN] ^b Levothox [AF]
			^B 1.31	26.57	26.66	^a Oroxine [AS]	

■ LEVOTHYROXINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Eltroxin and Levothox are not interchangeable with Oroxine, Eutroxsig, Levoxine, Levothyroxine Lup, APO-Levothyroxine or Thyrox on a dose to dose basis, and dose titration may be required.

Note Pharmaceutical benefits that have the form levothyroxine sodium 50 microgram tablet, 200 (Eltroxin and Levothox brands) are equivalent for the purposes of substitution.

Note Pharmaceutical benefits that have the form levothyroxine sodium 50 microgram tablet, 200 (Eutroxsig, Oroxine, Levothyroxine Lup and APO-Levothyroxine brands) and pharmaceutical benefits that have the form levothyroxine sodium 50 microgram tablet, 2 x 100 (Levoxine and Thyrox brands) are equivalent for the purposes of substitution.

levothyroxine sodium 50 microgram tablet, 200

2174K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	23.09	24.49	^a APO-Levothyroxine [XT] ^a Levothyroxine Lup [GQ] ^b Eltroxin [LT]	^a Eutroxsig [LN] ^b Levothox [AF]
			^B 1.31	24.40	24.49	^a Oroxine [AS]	

■ LEVOTHYROXINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Eltroxin and Levothox are not interchangeable with Oroxine, Eutroxsig, Levoxine, Levothyroxine Lup, APO-Levothyroxine or Thyrox on a dose to dose basis, and dose titration may be required.

Note Pharmaceutical benefits that have the form levothyroxine sodium 100 microgram tablet, 200 (Eltroxin and Levothox brands) are equivalent for the purposes of substitution.

Note Pharmaceutical benefits that have the form levothyroxine sodium 100 microgram tablet, 200 (Eutroxsig, Oroxine, Levothyroxine Lup and APO-Levothyroxine brands) and pharmaceutical benefits that have the form levothyroxine sodium 100 microgram tablet, 2 x 100 (Levoxine and Thyrox brands) are equivalent for the purposes of substitution.

levothyroxine sodium 100 microgram tablet, 200

2175L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	23.45	24.85	^a APO-Levothyroxine [XT] ^a Levothyroxine Lup [GQ] ^b Eltroxin [LT]	^a Eutroxsig [LN] ^b Levothox [AF]
			^B 1.31	24.76	24.85	^a Oroxine [AS]	

■ LEVOTHYROXINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Eltroxin and Levothox are not interchangeable with Oroxine, Eutroxsig, Levoxine, Levothyroxine Lup, APO-Levothyroxine or Thyrox on a dose to dose basis, and dose titration may be required.

Note Pharmaceutical benefits that have the form levothyroxine sodium 75 microgram tablet, 200 (Eltroxin and Levothox brands) are equivalent for the purposes of substitution.

Note Pharmaceutical benefits that have the form levothyroxine sodium 75 microgram tablet, 200 (Eutroxsig, Oroxine, Levothyroxine Lup and APO-Levothyroxine brands) and pharmaceutical benefits that have the form levothyroxine sodium 75 microgram tablet, 2 x 100 (Levoxine and Thyrox brands) are equivalent for the purposes of substitution.

levothyroxine sodium 75 microgram tablet, 200

9287T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	23.26	24.66	^a APO-Levothyroxine [XT] ^a Levothyroxine Lup [GQ] ^b Eltroxin [LT]	^a Eutroxsig [LN] ^b Levothox [AF]
			^B 1.32	24.58	24.66	^a Oroxine [AS]	

■ NETUPITANT + PALONOSETRON

Note This medicine is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Note Various sources of information outline the emetic risk associated with cancer treatment. Examples include the National Comprehensive Cancer Network guidelines (USA), eviQ guidelines and approved Product Information of individual drugs. These examples are not a comprehensive list of which anti-cancer drugs that have moderate to high emesis risk.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

14443

Nausea and vomiting

Clinical criteria:

- The treatment must be in combination with dexamethasone, unless contraindicated, **AND**
- The treatment must be for prevention of nausea and vomiting associated with moderate to highly emetogenic anti-cancer therapy.

netupitant 300 mg + palonosetron 500 microgram capsule, 1

10731T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	100.79	30.00	Akynzeo [JZ]

■ RISANKIZUMAB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or commencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR

- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

risankizumab 75 mg/0.83 mL injection, 2 x 0.83 mL syringes

11858D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	5401.42	30.00	Skyrizi [VE]

▪ **RISANKIZUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment. A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or commencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 150 mg for weeks 0 and 4, then 150 mg every 12 weeks thereafter.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 150 mg for weeks 0 and 4, then 150 mg every 12 weeks thereafter.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 150 mg for weeks 0 and 4, then 150 mg every 12 weeks thereafter.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of prior biological treatment, including dosage, date and duration of treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 150 mg for weeks 0 and 4, then 150 mg every 12 weeks thereafter.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

risankizumab 75 mg/0.83 mL injection, 2 x 0.83 mL syringes

11827L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	5401.42	30.00	Skyrizi [VE]

▪ **SECUKINUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of

more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years). An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

secukinumab 150 mg/mL injection, 2 x 1 mL pen devices

10494H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1404.60	30.00	Cosentyx [NV]

▪ **SECUKINUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or commencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

secukinumab 150 mg/mL injection, 2 x 1 mL pen devices

10425Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1404.60	30.00	Cosentyx [NV]

■ SECUKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and

wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**

- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**

- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

secukinumab 150 mg/mL injection, 2 x 1 mL pen devices

10910F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*5428.41	30.00	Cosentyx [NV]

■ TILDRAKIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a

minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment. A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tildrakizumab 100 mg/mL injection, 1 mL syringe

11613F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3272.18	30.00	Ilumya [RA]

▪ **TILDRAKIZUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4

weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 100 mg for weeks 0 and 4, then 100 mg every 12 weeks thereafter.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.
- An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and

(ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 100 mg for weeks 0 and 4, then 100 mg every 12 weeks thereafter.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 100 mg for weeks 0 and 4, then 100 mg every 12 weeks thereafter.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 100 mg for weeks 0 and 4, then 100 mg every 12 weeks thereafter.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 28 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tildrakizumab 100 mg/mL injection, 1 mL syringe

11616J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3272.18	30.00	Ilumya [RA]

▪ **USTEKINUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised

therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years). An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 1 repeat will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 1 repeat will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial

9305R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3944.14	30.00	Stelara [JC]

■ USTEKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial

9304Q

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	2	..	3944.14	30.00	Stelara [JC]

Highly Specialised Drugs Program (Private Hospital)

■ BACLOFEN

Note Pharmaceutical benefits that have the form baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule and pharmaceutical benefits that have the form baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

9488

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity of cerebral origin.

Authority required (STREAMLINED)

9637

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to multiple sclerosis.

Authority required (STREAMLINED)

9489

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord injury.

Authority required (STREAMLINED)

9524

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord disease.

baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules

11128Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	801.58	^a Sintetica Baclofen Intrathecal [BZ]

baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule

6284R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	10	..	..	*801.57	^a Bacthecal [DZ]	^a Lioresal Intrathecal [NV]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where

they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or

have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11590B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	272.03	^a Inflectra [PF] ^a Renflexis [OQ]	^a Remicade [JC]

▪ **INFLIXIMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before

they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of

disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9602

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required (STREAMLINED)

9584

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11595G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1315.97	^a Inflectra [PF]	^a Renflexis [OQ]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion

of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

-
- (a) a completed authority prescription form(s); and
(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
(i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
(ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied by the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

9617E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	272.03	^a Inflectra [PF] ^a Renflexis [OQ]	^a Remicade [JC]

▪ **LENALIDOMIDE**

Caution This drug is a category X drug and must not be given to pregnant women. If lenalidomide is taken during pregnancy, a teratogenic effect of lenalidomide in humans cannot be ruled out.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note Patients receiving lenalidomide under the PBS listing must be registered in the risk management program relevant for the brand of lenalidomide being prescribed and dispensed: Revlimid - i-access program; Cipla Lenalidomide - Pregnancy Prevention Program; Lenalidomide Dr.Reddy's - Reddy-2-Assist Controlled Access Program; Lenalide - Juno Connected™; Lenalidomide Sandoz - MyCheckPoint Pregnancy Prevention Program; Lenalidomide Teva - Pregnancy Prevention Program; Lenalidomide Viatriis - Viatriis Care.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Triple combination therapy consisting of carfilzomib, lenalidomide and dexamethasone

Treatment criteria:

- Patient must be undergoing concurrent treatment with carfilzomib obtained through the PBS, **AND**
- Patient must not be undergoing simultaneous treatment with this drug obtained under another PBS listing.

lenalidomide 10 mg capsule, 21

13658W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	1569.67	Cipla Lenalidomide [LR] Lenalidomide Dr.Reddy's [RI] Lenalidomide-Teva [TB] Revlimid [CJ]	Lenalide [JU] Lenalidomide Sandoz [SZ] Lenalidomide Viatriis [AF]

lenalidomide 15 mg capsule, 21

13657T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	1880.62	Cipla Lenalidomide [LR] Lenalidomide Dr.Reddy's [RI] Lenalidomide-Teva [TB] Revlimid [CJ]	Lenalide [JU] Lenalidomide Sandoz [SZ] Lenalidomide Viatriis [AF]

lenalidomide 25 mg capsule, 21

13660Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	2445.58	Cipla Lenalidomide [LR] Lenalidomide Dr.Reddy's [RI] Lenalidomide-Teva [TB] Revlimid [CJ]	Lenalide [JU] Lenalidomide Sandoz [SZ] Lenalidomide Viatriis [AF]

lenalidomide 5 mg capsule, 21

13642B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	1209.63	Cipla Lenalidomide [LR] Lenalidomide Dr.Reddy's [RI] Lenalidomide-Teva [TB] Revlimid [CJ]	Lenalide [JU] Lenalidomide Sandoz [SZ] Lenalidomide Viatriis [AF]

■ NUSINERSEN

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Literature references for various instruments measuring motor function and quality of life in the context of spinal muscular atrophy are:

Revised Upper Limb Module

Mazzone et al. 2017. Revised upper limb module for spinal muscular atrophy: Development of a new module. **Muscle & Nerve** 55(6):869-874

Hammersmith Functional Motor Scale - Expanded

Ramsey et al. 2017. Revised Hammersmith Scale for spinal muscular atrophy: A SMA specific clinical outcome assessment tool. **PLoS ONE** 12(2): e0172346. doi:10.1371/journal.pone.0172346.

6-Minute Walk Test (6MWT)

American Thoracic Society. 2002. ATS statement: Guidelines for the six-minute walk test. **American Journal of Respiratory and Critical Care Medicine** 166(1), pp 111-117

The National Health Foundation of Australia has 6MWT test standardised instructions and recording forms located at:

<https://www.hearonline.org.au/resources/documents-and-links#exercise>

SMA Health Index

Zizzi et al. 2021. The Spinal Muscular Atrophy Health Index (SMA-HI): A Novel Outcome for Measuring How a Patient Feels and Functions. **Muscle & Nerve** 63(10), pp 837-844

SMA Functional Rating Scale

Elsheikh et al. 2018. Reliability of Spinal Muscular Atrophy Functional Rating Scale (SMAFRS) in Ambulatory Adults with Spinal Muscular Atrophy. **Neurology** April (15 Supplement) P4.452

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment in an adult where treatment was initiated in adulthood

Clinical criteria:

- The treatment must be each of: (i) occurring from week 104 onwards relative to the first administered dose, (ii) demonstrating a clinically meaningful response; OR
- The treatment must be occurring within the first 104 weeks from the first administered dose, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing continuation of existing PBS-subsidised treatment with this drug, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Where this authority application seeks to continue treatment beyond the first 104 weeks of treatment, comprehensive assessment must be undertaken periodically and documented, involving the patient and the treating physician to establish agreement that treatment is continuing to produce a clinically meaningful response.

A clinically meaningful response is present where an improvement, stabilisation or minimal decline in symptoms has occurred as a result of this drug treatment and where there is agreement between the treating physician and patient over what constitutes improvement, stabilisation, or minimal decline.

PBS subsidy must cease if there is no agreement on whether a clinically meaningful response is present.

Undertake re-assessments for a clinically meaningful response at least every six months. Document these re-assessments in the patient's medical records.

In undertaking comprehensive assessments, where practical, a clinically meaningful response assessment encompasses the patient's motor function as assessed using an instrument like the Revised Upper Limb Module (RULM), Hammersmith Functional Motor Scale - Expanded (HFMSE) or 6-minute walk test (6MWT), and the patient's quality of life including, but not limited to, level of independence. Quality of life may be informed by use of the SMA Health Index (SMA-HI) or SMA Functional Rating Scale (SMAFRS).

nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial

13045N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	110048.37	Spinraza [BD]

■ NUSINERSEN

Note Recognised hospitals in the management of SMA are Queensland Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment in a child or adult, but where treatment was initiated during childhood

Clinical criteria:

- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA, **AND**
- Patient must be undergoing continuation of existing PBS-subsidised treatment with this drug, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial

13091B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	110048.37	Spinraza [BD]

▪ **NUSINERSEN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Initial PBS-subsidised treatment in an adult who did not initiate PBS subsidy during childhood

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Population criteria:

- Patient must be at least 19 years of age at the time of this authority application, but never claimed PBS subsidy for a disease modifying treatment during childhood, **AND**
- Patient must have SMA where the onset of signs/symptoms (at least one) of SMA first occurred prior to their 19th birthday (SMA symptom onset after this age will be considered type IV SMA, which is not PBS-subsidised).

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing initial PBS-subsidised treatment for untreated disease - prescribe up to 3 repeat prescriptions to enable dosing occurring at days: 0 (original prescription), 14 (repeat 1), 28 (repeat 2), 63 (repeat 3) (i.e. the loading doses); OR
- Patient must be undergoing initial PBS-subsidised treatment, but the patient has initiated treatment via non-PBS supply (e.g. clinical trial, sponsor compassionate access) - prescribe zero repeat prescriptions where loading doses are complete, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Signs and symptoms of spinal muscular atrophy in the context of this PBS restriction are:

- (i) Failure to meet or regression in ability to perform age-appropriate motor milestones,
- (ii) Proximal weakness,
- (iii) Hypotonia,
- (iv) Absence of deep tendon reflexes,
- (v) Failure to gain weight appropriate for age,
- (vi) Any active denervation or chronic neurogenic changes found on electromyography,
- (vii) A compound muscle action potential below normative values for an age-matched child.

In this authority application, confirm:

- (1) the patient's medical history is consistent with a diagnosis of childhood onset spinal muscular atrophy,
- (2) which of the above (i to vii) (at least 1) were present during childhood,

(3) the age of the patient (rounded to the nearest year) when the first sign/symptom was observed.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Changing the prescribed therapy

Treatment criteria:

- Patient must be undergoing a change in prescribed SMA drug to this drug - the drug treatment being replaced was a PBS benefit initiated after the patient's 19th birthday, **AND**
- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- Patient must be untreated with gene therapy, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The prescriber has given consideration to whether a 'wash out' period is recommended or not prior to changing the prescribed therapy.

Note Subsequent changes in the prescribed drug where applicable are to occur under the 'Continuing treatment' phase listing of the drug that therapy is changing to.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial

13064N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	110048.37	Spinraza [BD]

▪ **NUSINERSEN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Initial PBS-subsidised treatment in a child

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Population criteria:

- Patient must be of an age that is prior to their 19th birthday at the time of this authority application, **AND**
- Patient must have SMA type III where the onset of signs/symptoms of SMA first occurred after their 3rd birthday, but before their 19th birthday (SMA type IIIB/IIIC).

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA, **AND**
- Patient must be undergoing initial PBS-subsidised treatment for untreated disease - prescribe up to 3 repeat prescriptions to enable dosing occurring at days: 0 (original prescription), 14 (repeat 1), 28 (repeat 2), 63 (repeat 3) (i.e. the loading doses); OR

- Patient must be undergoing initial PBS-subsidised treatment, but the patient has initiated treatment via non-PBS supply (e.g. clinical trial, sponsor compassionate access) - prescribe zero repeat prescriptions where loading doses are complete, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Signs and symptoms of spinal muscular atrophy in the context of this PBS restriction are:

- (i) Failure to meet or regression in ability to perform age-appropriate motor milestones,
- (ii) Proximal weakness,
- (iii) Hypotonia,
- (iv) Absence of deep tendon reflexes,
- (v) Any active denervation or chronic neurogenic changes found on electromyography,
- (vi) A compound muscle action potential below normative values for an age-matched child.

In this authority application, confirm:

- (1) the patient's medical history is consistent with a diagnosis of type IIIB/IIIC spinal muscular atrophy,
- (2) which of the above (i to vi) (at least 1) were present after their 3rd birthday, but before their 19th birthday,
- (3) the age of the patient (rounded to the nearest year) when the first sign/symptom was observed.

Note Recognised hospitals in the management of SMA are Queensland Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Changing the prescribed therapy

Treatment criteria:

- Patient must be undergoing a change in prescribed SMA drug to this drug - the drug treatment being replaced was a PBS benefit initiated prior to the patient's 19th birthday for SMA type IIIB/IIIC, **AND**
- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- Patient must be untreated with gene therapy, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The prescriber has given consideration to whether a 'wash out' period is recommended or not prior to changing the prescribed therapy.

Note Subsequent changes in the prescribed drug where applicable are to occur under the 'Continuing treatment' phase listing of the drug that therapy is changing to.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial

13111C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	110048.37	Spinraza [BD]

■ RISDIPLAM

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available

on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic Type I, II or IIIa spinal muscular atrophy (SMA)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must have experienced at least two of the defined signs and symptoms of SMA type I, II or IIIa prior to 3 years of age, **AND**
- The treatment must be given concomitantly with best supportive care for this condition, **AND**
- The treatment must not be in combination with PBS-subsidised treatment with nusinersen for this condition, **AND**
- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic.

Clinical criteria:

- Patient must be untreated with gene therapy.

Population criteria:

- Patient must be 18 years of age or under.

Defined signs and symptoms of type I SMA are:

- i) Onset before 6 months of age; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Hypotonia; or
- v) Absence of deep tendon reflexes; or
- vi) Failure to gain weight appropriate for age; or
- vii) Any active chronic neurogenic changes; or
- viii) A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type II SMA are:

- i) Onset between 6 and 18 months; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Weakness in trunk righting/derotation; or
- v) Hypotonia; or
- vi) Absence of deep tendon reflexes; or
- vii) Failure to gain weight appropriate for age; or
- viii) Any active chronic neurogenic changes; or
- ix) A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type IIIa SMA are:

- i) Onset between 18 months and 3 years of age; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Hypotonia; or
- v) Absence of deep tendon reflexes; or
- vi) Failure to gain weight appropriate for age; or
- vii) Any active chronic neurogenic changes; or
- viii) A compound muscle action potential below normative values for an age-matched child.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

Application for authorisation of initial treatment must be in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Spinal muscular atrophy PBS Authority Application Form which includes the following:
 - i) specification of SMA type (I, II or IIIa); and
 - ii) sign(s) and symptom(s) that the patient has experienced; and

- (iii) patient's age at the onset of sign(s) and symptom(s).
The approved Product Information recommended dosing is as follows:
- (i) 16 days to less than 2 months of age: 0.15 mg/kg
 - (ii) 2 months to less than 2 years of age: 0.20 mg/kg
 - (iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg
 - (iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

- 1 unit where (i) applies;
- 2 units where (ii) applies;
- 3 units where (iii) applies;
- 3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

12610Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	10890.26	Evrysdi [RO]

▪ **RISDIPLAM**

Note The maximum quantity of drug to be subsidised per dispensing, as well as the number of repeat prescriptions is to be as follows:

- Patient weight greater than 19 kg: up to 3 units per dispensing, with up to 5 repeat prescriptions
- Patient weight between 17 kg to 19 kg: up to 3 units per dispensing, with up to 4 repeat prescriptions
- Patient weight between 13 kg to 17 kg: up to 2 units per dispensing, with up to 5 repeat prescriptions
- Patient weight between 10 kg up to 13 kg: up to 2 units per dispensing, with up to 4 repeat prescriptions
- Patient weight less than 10 kg: up to 1 unit per dispensing, with up to 5 repeat prescriptions

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment with this drug of either symptomatic Type I, II or IIIa SMA, or, pre-symptomatic SMA

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition; OR
- Patient must be eligible for continuing PBS-subsidised treatment with nusinersen for this condition, **AND**
- The treatment must not be in combination with PBS-subsidised treatment with nusinersen for this condition, **AND**
- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug, **AND**
- The treatment must be given concomitantly with best supportive care for this condition.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, **AND**
- Patient must not be undergoing treatment through this 'Continuing treatment' listing where the most recent PBS authority approval for this PBS-indication has been for gene therapy.

Population criteria:

- Patient must have been 18 years of age or younger at the time of initial treatment with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

In a patient who wishes to switch from PBS-subsidised nusinersen to PBS-subsidised risdiplam for this condition a wash out period may be required.

The quantity of drug and number of repeat prescriptions prescribed is to be in accordance with the relevant 'Note' attached to this listing.

The approved Product Information recommended dosing is as follows:

- (i) 16 days to less than 2 months of age: 0.15 mg/kg
- (ii) 2 months to less than 2 years of age: 0.20 mg/kg
- (iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg
- (iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

- 1 unit where (i) applies;
- 2 units where (ii) applies;
- 3 units where (iii) applies;
- 3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

12609P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	5	..	10890.26	Evrysdi [RO]

■ RISDIPLAM

Note For the next authority application after this application, continue treatment through the 'Treatment phase: Continuing treatment' under the indication: 'Symptomatic Type I, II or IIIa spinal muscular atrophy (SMA)'.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Initial treatment occurring after onasemnogene abeparvovec therapy in a patient with Type 1 SMA

Clinical criteria:

- Patient must have experienced a regression in a developmental state listed below (see 'Definition') despite treatment with gene therapy - confirm that this: (i) not due to an acute concomitant illness; (ii) not due to non-compliance to best-supportive care, (iii) apparent for at least 3 months, (iv) verified by another clinician in the treatment team - state the full name of this clinician plus their profession (e.g. medical practitioner, nurse, physiotherapist; this is not an exhaustive list of examples), **AND**
- The treatment must not be a PBS-subsidised benefit where the condition has progressed to a point where invasive permanent assisted ventilation (i.e. ventilation via tracheostomy tube for at least 16 hours per day) is required in the absence of potentially reversible causes, **AND**
- The treatment must be given concomitantly with best supportive care for this condition, **AND**
- The treatment must not be in combination with PBS-subsidised treatment with nusinersen for this condition.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, **AND**
- Patient must be undergoing treatment under this Treatment phase listing once only - for continuing treatment beyond this authority application, refer to the drug's relevant 'Continuing treatment' listing for the patient's SMA type.

Population criteria:

- Patient must have a prior authority approval for any drug PBS-listed for symptomatic Type 1 SMA, with at least one approval having been for gene therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Do not resubmit previously submitted documentation concerning the diagnosis and type of SMA.

Confirm that a previous PBS authority application has been approved for symptomatic Type 1 SMA.

Definition:

Various childhood developmental states (1 to 9) are listed below, some followed by further observations (a up to d). Where at least one developmental state/observation is no longer present, that developmental state has regressed.

0. Absence of developmental states (1 to 9) listed below:

1. Rolls from side to side on back;
2. Child holds head erect for at least 3 seconds unsupported;
3. Sitting, but with assistance;
4. Sitting without assistance:
 - (a) Child sits up straight with the head erect for at least 10 seconds;
 - (b) Child does not use arms or hands to balance body or support position.
5. Hands and knees crawling:
 - (a) Child alternately moves forward or backwards on hands and knees;
 - (b) The stomach does not touch the supporting surface;
 - (c) There are continuous and consecutive movements at least 3 in a row.
6. Standing with assistance:
 - (a) Child stands in upright position on both feet, holding onto a stable object (e.g. furniture) with both hands and without leaning on object;
 - (b) The body does not touch the stable object, and the legs support most of the body weight;
 - (c) Child thus stands with assistance for at least 10 seconds.

7. Standing alone:

- (a) Child stands in upright position on both feet (not on the toes) with the back straight;
- (b) The leg supports 100% of the child's weight;
- (c) There is no contact with a person or object;

- (d) Child stands alone for at least 10 seconds.
8. Walking with assistance:
- (a) Child is in an upright position with the back straight;
 - (b) Child makes sideways or forced steps by holding onto a stable object (e.g. furniture) with 1 or both hands;
 - (c) One leg moves forward while the other supports part of the body weight;
 - (d) Child takes at least 5 steps in this manner.
9. Walking alone:
- (a) Child takes at least 5 steps independently in upright position with the back straight;
 - (b) One leg moves forward while the other supports most of the body weight;
 - (c) There is no contact with a person or object.

Confirm which developmental state has regressed by: (i) stating the overall developmental state (1 - 9) the patient was in at the time of gene therapy, or, the best developmental state achieved since gene therapy, and (ii) stating the patient's current overall developmental state (i.e. a number that is lower than stated in (i)).

Where the patient has neither regressed from a developmental state nor reached the next developmental state, PBS-subsidy of this benefit is not available.

The approved Product Information recommended dosing is as follows:

- (i) 16 days to less than 2 months of age: 0.15 mg/kg
- (ii) 2 months to less than 2 years of age: 0.20 mg/kg
- (iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg
- (iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

- 1 unit where (i) applies;
- 2 units where (ii) applies;
- 3 units where (iii) applies;
- 3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12959C	±1	..	10890.26	Evrysdi [RO]

■ RISDIPLAM

Note Recognised hospitals in the management of SMA are Queensland Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Pre-symptomatic spinal muscular atrophy (SMA)

Treatment Phase: Initial treatment with this drug of pre-symptomatic spinal muscular atrophy (SMA)

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA.

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- The condition must have genetic confirmation that there are 1 to 2 copies of the survival motor neuron 2 (SMN2) gene, **AND**
- The condition must be pre-symptomatic, **AND**
- The treatment must be given concomitantly with best supportive care for this condition, **AND**
- Patient must be untreated with gene therapy.

Population criteria:

- Patient must be aged under 36 months prior to commencing treatment. Application for authorisation of initial treatment must be in writing (lodged via postal service or electronic upload) and must include:

(a) a completed authority prescription form; and

(b) a completed Spinal muscular atrophy PBS Authority Application Form which includes the following:

(i) confirmation of genetic diagnosis of SMA; and

(ii) a copy of the results substantiating the number of SMN2 gene copies determined by quantitative polymerase chain reaction (qPCR) or multiple ligation dependent probe amplification (MLPA)

The quantity of drug and number of repeat prescriptions prescribed is to be in accordance with the relevant 'Note' attached to this listing.

The approved Product Information recommended dosing is as follows:

(i) 16 days to less than 2 months of age: 0.15 mg/kg

(ii) 2 months to less than 2 years of age: 0.20 mg/kg

(iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg

(iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

1 unit where (i) applies;

2 units where (ii) applies;

3 units where (iii) applies;

3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13647G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	10890.26	Evrysdi [RO]

■ RISDIPLAM

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Initial PBS-subsidised treatment with this drug in an adult who did not initiate PBS subsidy with this drug during childhood

Population criteria:

- Patient must be at least 19 years of age at the time of this authority application, but never claimed PBS subsidy for a disease modifying treatment during childhood, **AND**
- Patient must have SMA where the onset of signs/symptoms (at least one) of SMA first occurred prior to their 19th birthday (SMA symptom onset after this age will be considered type IV SMA, which is not PBS-subsidised).

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing initial PBS-subsidised treatment with this drug for untreated disease; OR
- Patient must be undergoing initial PBS-subsidised treatment, but the patient has initiated treatment via non-PBS supply (e.g. clinical trial, sponsor compassionate access), **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Signs and symptoms of spinal muscular atrophy in the context of this PBS restriction are:

- (i) Failure to meet or regression in ability to perform age-appropriate motor milestones,
- (ii) Proximal weakness,
- (iii) Hypotonia,
- (iv) Absence of deep tendon reflexes,
- (v) Failure to gain weight appropriate for age,
- (vi) Any active denervation or chronic neurogenic changes found on electromyography,
- (vii) A compound muscle action potential below normative values for an age-matched child.

In this authority application, confirm:

- (1) the patient's medical history is consistent with a diagnosis of childhood onset spinal muscular atrophy,
- (2) which of the above (i to vii) (at least 1) were present during childhood,
- (3) the age of the patient (rounded to the nearest year) when the first sign/symptom was observed.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13632L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
±3	7	..	*32574.03	Evrysdi [RO]	

■ RISDIPLAM

Note Literature references for various instruments measuring motor function and quality of life in the context of spinal muscular atrophy are:

Revised Upper Limb Module

Mazzone et al. 2017. Revised upper limb module for spinal muscular atrophy: Development of a new module. **Muscle & Nerve** 55(6):869-874

Hammersmith Functional Motor Scale - Expanded

Ramsey et al. 2017. Revised Hammersmith Scale for spinal muscular atrophy: A SMA specific clinical outcome assessment tool. **PLoS ONE** 12(2): e0172346. doi:10.1371/journal.pone.0172346.

6-Minute Walk Test (6MWT)

American Thoracic Society. 2002. ATS statement: Guidelines for the six-minute walk test. **American Journal of Respiratory and Critical Care Medicine** 166(1), pp 111-117

The National Heart Foundation of Australia has 6MWT test standardised instructions and recording forms located at:

<https://www.heartonline.org.au/resources/documents-and-links#exercise>

SMA Health Index

Zizzi et al. 2021. The Spinal Muscular Atrophy Health Index (SMA-HI): A Novel Outcome for Measuring How a Patient Feels and Functions. **Muscle & Nerve** 63(10), pp 837-844

SMA Functional Rating Scale

Elsheikh et al. 2018. Reliability of Spinal Muscular Atrophy Functional Rating Scale (SMAFRS) in Ambulatory Adults with Spinal Muscular Atrophy. **Neurology** April (15 Supplement) P4.452

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment in an adult where treatment was initiated in adulthood

Treatment criteria:

- Patient must be undergoing continuation of existing PBS-subsidised treatment with this drug; OR
- Patient must be undergoing a change in prescribed SMA drug to this drug - the drug treatment being replaced was a PBS benefit initiated after the patient's 19th birthday, **AND**
- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- The treatment must be each of: (i) occurring from week 104 onwards relative to the first administered dose, (ii) demonstrating a clinically meaningful response; OR
- The treatment must be occurring within the first 104 weeks from the first administered dose, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

Where this authority application seeks to continue treatment beyond the first 104 weeks of treatment, comprehensive assessment must be undertaken periodically and documented, involving the patient and the treating physician to establish agreement that treatment is continuing to produce a clinically meaningful response.

A clinically meaningful response is present where an improvement, stabilisation or minimal decline in symptoms has occurred as a result of this drug treatment and where there is agreement between the treating physician and patient over what constitutes improvement, stabilisation, or minimal decline.

PBS subsidy must cease if there is no agreement on whether a clinically meaningful response is present.

Undertake re-assessments for a clinically meaningful response at least every six months. Document these re-assessments in the patient's medical records.

In undertaking comprehensive assessments, where practical, a clinically meaningful response assessment encompasses the patient's motor function as assessed using an instrument like the Revised Upper Limb Module (RULM), Hammersmith Functional Motor Scale - Expanded (HFMSE) or 6-minute walk test (6MWT), and the patient's quality of life including, but not limited to, level of independence. Quality of life may be informed by use of the SMA Health Index (SMA-HI) or SMA Functional Rating Scale (SMAFRS).

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13646F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±3	5	..	*32574.03	Evrysdi [RO]

■ RISDIPLAM

Note The maximum quantity of drug to be subsidised per dispensing, as well as the number of repeat prescriptions is to be as follows:

Patient weight greater than 19 kg: up to 3 units per dispensing, with up to 5 repeat prescriptions

Patient weight between 17 kg to 19 kg: up to 3 units per dispensing, with up to 4 repeat prescriptions

Patient weight between 13 kg to 17 kg: up to 2 units per dispensing, with up to 5 repeat prescriptions

Patient weight between 10 kg up to 13 kg: up to 2 units per dispensing, with up to 4 repeat prescriptions

Patient weight less than 10 kg: up to 1 unit per dispensing, with up to 5 repeat prescriptions

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment in a child or adult, but where treatment was initiated during childhood

Treatment criteria:

- Patient must be undergoing continuation of existing PBS-subsidised treatment with this drug; OR
- Patient must be undergoing a change in prescribed SMA drug to this drug - the drug treatment being replaced was a PBS benefit initiated prior to the patient's 19th birthday for SMA type IIIB/IIIC, **AND**
- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The quantity of drug and number of repeat prescriptions prescribed is to be in accordance with the relevant 'Note' attached to this listing.

The approved Product Information recommended dosing is as follows:

(i) 16 days to less than 2 months of age: 0.15 mg/kg

(ii) 2 months to less than 2 years of age: 0.20 mg/kg

(iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg

(iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

1 unit where (i) applies;

2 units where (ii) applies;

3 units where (iii) applies;

3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13659X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±3	5	..	*32574.03	Evrysdi [RO]

▪ RISDIPLAM

Note The maximum quantity of drug to be subsidised per dispensing, as well as the number of repeat prescriptions is to be as follows:

Patient weight greater than 19 kg: up to 3 units per dispensing, with up to 5 repeat prescriptions

Patient weight between 17 kg to 19 kg: up to 3 units per dispensing, with up to 4 repeat prescriptions

Patient weight between 13 kg to 17 kg: up to 2 units per dispensing, with up to 5 repeat prescriptions

Patient weight between 10 kg up to 13 kg: up to 2 units per dispensing, with up to 4 repeat prescriptions

Patient weight less than 10 kg: up to 1 unit per dispensing, with up to 5 repeat prescriptions

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Initial PBS-subsidised treatment with this drug in a child

Population criteria:

- Patient must be of an age that is prior to their 19th birthday at the time of this authority application, **AND**
- Patient must have SMA type III where the onset of signs/symptoms of SMA first occurred after their 3rd birthday, but before their 19th birthday (SMA type IIIB/IIIC).

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing initial PBS-subsidised treatment with this drug for untreated disease; OR
- Patient must be undergoing initial PBS-subsidised treatment, but the patient has initiated treatment via non-PBS supply (e.g. clinical trial, sponsor compassionate access), **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Signs and symptoms of spinal muscular atrophy in the context of this PBS restriction are:

- (i) Failure to meet or regression in ability to perform age-appropriate motor milestones,
- (ii) Proximal weakness,
- (iii) Hypotonia,
- (iv) Absence of deep tendon reflexes,
- (v) Any active denervation or chronic neurogenic changes found on electromyography,
- (vi) A compound muscle action potential below normative values for an age-matched child.

In this authority application, confirm:

- (1) the patient's medical history is consistent with a diagnosis of type IIIB/IIIC spinal muscular atrophy,
- (2) which of the above (i to vi) (at least 1) were present after their 3rd birthday, but before their 19th birthday,
- (3) the age of the patient (rounded to the nearest year) when the first sign/symptom was observed.

The quantity of drug and number of repeat prescriptions prescribed is to be in accordance with the relevant 'Note' attached to this listing.

The approved Product Information recommended dosing is as follows:

- (i) 16 days to less than 2 months of age: 0.15 mg/kg

- (ii) 2 months to less than 2 years of age: 0.20 mg/kg
 (iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg
 (iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

- 1 unit where (i) applies;
 2 units where (ii) applies;
 3 units where (iii) applies;
 3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13639W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±3	5	..	*32574.03	Evrysdi [RO]

Highly Specialised Drugs Program (Public Hospital)

▪ BACLOFEN

Note Pharmaceutical benefits that have the form baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule and pharmaceutical benefits that have the form baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

6925

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity of cerebral origin.

Authority required (STREAMLINED)

6939

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to multiple sclerosis.

Authority required (STREAMLINED)

6940

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord injury.

Authority required (STREAMLINED)

6911

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord disease.

baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules

11126N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	762.70	^a Sintetica Baclofen Intrathecal [BZ]

baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule

5617P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	10	..	..	*762.70	^a Bacthecal [DZ]	^a Lioresal Intrathecal [NV]

▪ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where

they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or

have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11606W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	253.52	^a Inflectra [PF] ^a Renflexis [OQ]	^a Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before

they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of

disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum number of repeats may be authorised.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

8844

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required (STREAMLINED)

8940

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11605T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1267.60	^a Inflectra [PF]	^a Renflexis [OQ]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, bimekizumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab, and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab, and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab, 24 weeks of therapy for bimekizumab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of their course of initial treatment to ensure uninterrupted biological medicine supply.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment and no later than 4 weeks from the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab, adalimumab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion

of the most recent course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

A response to treatment is to be determined by comparison of current disease activity measurements relative to the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

-
- (a) a completed authority prescription form(s); and
(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
(i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
(ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 6 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks; (vi) deucravacitinib at a dose of 6 mg once daily for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, deucravacitinib, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast, deucravacitinib or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be at least 18 years of age.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Medicines webpage (www.health.gov.au/health-topics/medicines).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**

- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

5758C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	253.52	^a Inflectra [PF] ^a Renflexis [OQ]	^a Remicade [JC]

▪ **LENALIDOMIDE**

Caution This drug is a category X drug and must not be given to pregnant women. If lenalidomide is taken during pregnancy, a teratogenic effect of lenalidomide in humans cannot be ruled out.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note Patients receiving lenalidomide under the PBS listing must be registered in the risk management program relevant for the brand of lenalidomide being prescribed and dispensed: Revlimid - i-access program; Cipla Lenalidomide - Pregnancy Prevention Program; Lenalidomide Dr.Reddy's - Reddy-2-Assist Controlled Access Program; Lenalide - Juno Connected™; Lenalidomide Sandoz - MyCheckPoint Pregnancy Prevention Program; Lenalidomide Teva - Pregnancy Prevention Program; Lenalidomide Viatriis - Viatriis Care.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Triple combination therapy consisting of carfilzomib, lenalidomide and dexamethasone

Treatment criteria:

- Patient must be undergoing concurrent treatment with carfilzomib obtained through the PBS, **AND**
- Patient must not be undergoing simultaneous treatment with this drug obtained under another PBS listing.

lenalidomide 10 mg capsule, 21

13661B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	1521.30	Cipla Lenalidomide [LR] Lenalidomide Dr.Reddy's [RI] Lenalidomide-Teva [TB] Revlimid [CJ]	Lenalide [JU] Lenalidomide Sandoz [SZ] Lenalidomide Viatriis [AF]

lenalidomide 15 mg capsule, 21

13641Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	1832.25	Cipla Lenalidomide [LR] Lenalidomide Dr.Reddy's [RI] Lenalidomide-Teva [TB] Revlimid [CJ]	Lenalide [JU] Lenalidomide Sandoz [SZ] Lenalidomide Viatriis [AF]

lenalidomide 25 mg capsule, 21

13630J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	2397.21	Cipla Lenalidomide [LR] Lenalidomide Dr.Reddy's [RI] Lenalidomide-Teva [TB] Revlimid [CJ]	Lenalide [JU] Lenalidomide Sandoz [SZ] Lenalidomide Viatriis [AF]

lenalidomide 5 mg capsule, 21

13636Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	1161.26	Cipla Lenalidomide [LR] Lenalidomide Dr.Reddy's [RI] Lenalidomide-Teva [TB] Revlimid [CJ]	Lenalide [JU] Lenalidomide Sandoz [SZ] Lenalidomide Viatriis [AF]

■ NUSINERSEN

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Literature references for various instruments measuring motor function and quality of life in the context of spinal muscular atrophy are:

Revised Upper Limb Module

Mazzone et al. 2017. Revised upper limb module for spinal muscular atrophy: Development of a new module. **Muscle & Nerve** 55(6):869-874

Hammersmith Functional Motor Scale - Expanded

Ramsey et al. 2017. Revised Hammersmith Scale for spinal muscular atrophy: A SMA specific clinical outcome assessment tool. **PLoS ONE** 12(2): e0172346. doi:10.1371/journal.pone.0172346.

6-Minute Walk Test (6MWT)

American Thoracic Society. 2002. ATS statement: Guidelines for the six-minute walk test. **American Journal of Respiratory and Critical Care Medicine** 166(1), pp 111-117

The National Health Foundation of Australia has 6MWT test standardised instructions and recording forms located at:

<https://www.hearonline.org.au/resources/documents-and-links#exercise>

SMA Health Index

Zizzi et al. 2021. The Spinal Muscular Atrophy Health Index (SMA-HI): A Novel Outcome for Measuring How a Patient Feels and Functions. **Muscle & Nerve** 63(10), pp 837-844

SMA Functional Rating Scale

Elsheikh et al. 2018. Reliability of Spinal Muscular Atrophy Functional Rating Scale (SMAFRS) in Ambulatory Adults with Spinal Muscular Atrophy. **Neurology** April (15 Supplement) P4.452

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment in an adult where treatment was initiated in adulthood

Clinical criteria:

- The treatment must be each of: (i) occurring from week 104 onwards relative to the first administered dose, (ii) demonstrating a clinically meaningful response; OR
- The treatment must be occurring within the first 104 weeks from the first administered dose, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing continuation of existing PBS-subsidised treatment with this drug, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Where this authority application seeks to continue treatment beyond the first 104 weeks of treatment, comprehensive assessment must be undertaken periodically and documented, involving the patient and the treating physician to establish agreement that treatment is continuing to produce a clinically meaningful response.

A clinically meaningful response is present where an improvement, stabilisation or minimal decline in symptoms has occurred as a result of this drug treatment and where there is agreement between the treating physician and patient over what constitutes improvement, stabilisation, or minimal decline.

PBS subsidy must cease if there is no agreement on whether a clinically meaningful response is present.

Undertake re-assessments for a clinically meaningful response at least every six months. Document these re-assessments in the patient's medical records.

In undertaking comprehensive assessments, where practical, a clinically meaningful response assessment encompasses the patient's motor function as assessed using an instrument like the Revised Upper Limb Module (RULM), Hammersmith Functional Motor Scale - Expanded (HFMSE) or 6-minute walk test (6MWT), and the patient's quality of life including, but not limited to, level of independence. Quality of life may be informed by use of the SMA Health Index (SMA-HI) or SMA Functional Rating Scale (SMAFRS).

nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial

13068T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	110000.00	Spinraza [BD]

■ NUSINERSEN

Note Recognised hospitals in the management of SMA are Queensland Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment in a child or adult, but where treatment was initiated during childhood

Clinical criteria:

- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA, **AND**
- Patient must be undergoing continuation of existing PBS-subsidised treatment with this drug, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial

13084P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	110000.00	Spinraza [BD]

▪ **NUSINERSEN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Initial PBS-subsidised treatment in an adult who did not initiate PBS subsidy during childhood

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Population criteria:

- Patient must be at least 19 years of age at the time of this authority application, but never claimed PBS subsidy for a disease modifying treatment during childhood, **AND**
- Patient must have SMA where the onset of signs/symptoms (at least one) of SMA first occurred prior to their 19th birthday (SMA symptom onset after this age will be considered type IV SMA, which is not PBS-subsidised).

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing initial PBS-subsidised treatment for untreated disease - prescribe up to 3 repeat prescriptions to enable dosing occurring at days: 0 (original prescription), 14 (repeat 1), 28 (repeat 2), 63 (repeat 3) (i.e. the loading doses); OR
- Patient must be undergoing initial PBS-subsidised treatment, but the patient has initiated treatment via non-PBS supply (e.g. clinical trial, sponsor compassionate access) - prescribe zero repeat prescriptions where loading doses are complete, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Signs and symptoms of spinal muscular atrophy in the context of this PBS restriction are:

- (i) Failure to meet or regression in ability to perform age-appropriate motor milestones,
- (ii) Proximal weakness,
- (iii) Hypotonia,
- (iv) Absence of deep tendon reflexes,
- (v) Failure to gain weight appropriate for age,
- (vi) Any active denervation or chronic neurogenic changes found on electromyography,
- (vii) A compound muscle action potential below normative values for an age-matched child.

In this authority application, confirm:

- (1) the patient's medical history is consistent with a diagnosis of childhood onset spinal muscular atrophy,
- (2) which of the above (i to vii) (at least 1) were present during childhood,

(3) the age of the patient (rounded to the nearest year) when the first sign/symptom was observed.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Changing the prescribed therapy

Treatment criteria:

- Patient must be undergoing a change in prescribed SMA drug to this drug - the drug treatment being replaced was a PBS benefit initiated after the patient's 19th birthday, **AND**
- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- Patient must be untreated with gene therapy, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The prescriber has given consideration to whether a 'wash out' period is recommended or not prior to changing the prescribed therapy.

Note Subsequent changes in the prescribed drug where applicable are to occur under the 'Continuing treatment' phase listing of the drug that therapy is changing to.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial

13052Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	110000.00	Spinraza [BD]

▪ **NUSINERSEN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Initial PBS-subsidised treatment in a child

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Population criteria:

- Patient must be of an age that is prior to their 19th birthday at the time of this authority application, **AND**
- Patient must have SMA type III where the onset of signs/symptoms of SMA first occurred after their 3rd birthday, but before their 19th birthday (SMA type IIIB/IIIC).

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA, **AND**
- Patient must be undergoing initial PBS-subsidised treatment for untreated disease - prescribe up to 3 repeat prescriptions to enable dosing occurring at days: 0 (original prescription), 14 (repeat 1), 28 (repeat 2), 63 (repeat 3) (i.e. the loading doses); OR

- Patient must be undergoing initial PBS-subsidised treatment, but the patient has initiated treatment via non-PBS supply (e.g. clinical trial, sponsor compassionate access) - prescribe zero repeat prescriptions where loading doses are complete, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Signs and symptoms of spinal muscular atrophy in the context of this PBS restriction are:

- (i) Failure to meet or regression in ability to perform age-appropriate motor milestones,
- (ii) Proximal weakness,
- (iii) Hypotonia,
- (iv) Absence of deep tendon reflexes,
- (v) Any active denervation or chronic neurogenic changes found on electromyography,
- (vi) A compound muscle action potential below normative values for an age-matched child.

In this authority application, confirm:

- (1) the patient's medical history is consistent with a diagnosis of type IIIB/IIIC spinal muscular atrophy,
- (2) which of the above (i to vi) (at least 1) were present after their 3rd birthday, but before their 19th birthday,
- (3) the age of the patient (rounded to the nearest year) when the first sign/symptom was observed.

Note Recognised hospitals in the management of SMA are Queensland Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Changing the prescribed therapy

Treatment criteria:

- Patient must be undergoing a change in prescribed SMA drug to this drug - the drug treatment being replaced was a PBS benefit initiated prior to the patient's 19th birthday for SMA type IIIB/IIIC, **AND**
- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- Patient must be untreated with gene therapy, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The prescriber has given consideration to whether a 'wash out' period is recommended or not prior to changing the prescribed therapy.

Note Subsequent changes in the prescribed drug where applicable are to occur under the 'Continuing treatment' phase listing of the drug that therapy is changing to.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

nusinersen 12 mg/5 mL intrathecal injection, 5 mL vial

13083N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	110000.00	Spinraza [BD]

■ ONASEMNOGENE ABEPARVOVEC

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Other disease modifying therapies for this condition are: (i) nusinersen, (ii) risdiplam.

Note Recognised hospitals in the management of SMA are Queensland Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Note Accredited treatment centres and suppliers are those organisations accredited by the Gene Technology Regulator under section 92 of the **Gene Technology Act 2000**.

The following website provides a list of accredited organisations and may update without notice:
<https://www.ogtr.gov.au/what-weve-approved/accredited-organisations>

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Use in a patient untreated with disease modifying therapies for this condition

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- The condition must be pre-symptomatic SMA, with genetic confirmation that there are 3 copies of the **survival motor neuron 2 (SMN2)** gene, **AND**
- The treatment must not be a PBS-subsidised benefit where the condition has progressed to a point where invasive permanent assisted ventilation (i.e. ventilation via tracheostomy tube for at least 16 hours per day) is required in the absence of potentially reversible causes, **AND**
- The treatment must be given concomitantly with best supportive care for this condition.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA, **AND**
- Must be treated in a treatment centre that is each of: (i) recognised in the management of SMA, (ii) accredited in the use of this gene technology by the relevant authority, (iii) will(has) source(d) this product from an accredited supplier, as specified in the administrative notes to this listing, **AND**
- Patient must be undergoing treatment with this pharmaceutical benefit once only in a lifetime, **AND**
- Patient must not be undergoing treatment with this pharmaceutical benefit through this listing where prior treatment has occurred with any of: (i) nusinersen, (ii) risdiplam.

Population criteria:

- Patient must be no older than 9 months of age.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

State the weight of the patient in kilograms and request the appropriate product pack presentation with respect to the mix of 5.5 mL and 8.3 mL vials.

Confirm that genetic testing has been completed to demonstrate the following in support of an SMA diagnosis:

- (i) 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; or
- (ii) deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variance in the remaining single copy of the SMN1 gene.

Confirm that there is a genetic test finding that substantiates the number of SMN2 gene copies to be 3 and has been determined by quantitative polymerase chain reaction (qPCR) or multiple ligation dependent probe amplification (MLPA).

Quote the date, pathology provider name and any unique identifying serial number/code that links the genetic test result to the patient.

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 2 x 8.3 mL vials

13674Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13669K	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [3 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13682D	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [4 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13683E	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13681C	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [6 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13679Y	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [7 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13668J	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [8.3 mL vial], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13663D	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13671M	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 3 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13678X	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 4 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13665F	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [3 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13670L	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [4 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13672N	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 5 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13667H	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 6 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13664E	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 7 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13675R	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13680B	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [6 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13677W	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 8 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13673P	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 9 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13676T	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [7 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13662C	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [8 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
13666G	‡1	..	..	2527773.87	Zolgensma [NV]

■ ONASEMNOGENE ABEPARVOVEC

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Other disease modifying therapies for this condition are: (i) nusinersen, (ii) risdiplam.

Note Recognised hospitals in the management of SMA are Queensland Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Note Accredited treatment centres and suppliers are those organisations accredited by the Gene Technology Regulator under section 92 of the **Gene Technology Act 2000**.

The following website provides a list of accredited organisations and may update without notice:
<https://www.ogtr.gov.au/what-weve-approved/accredited-organisations>

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Use in a patient untreated with disease modifying therapies for this condition

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must have experienced at least two of the defined signs/symptoms of Type 1 SMA specified below; OR
- The condition must be pre-symptomatic SMA, with genetic confirmation that there are 1 to 2 copies of the survival motor neuron 2 (SMN2) gene, **AND**

- The treatment must not be a PBS-subsidised benefit where the condition has progressed to a point where invasive permanent assisted ventilation (i.e. ventilation via tracheostomy tube for at least 16 hours per day) is required in the absence of potentially reversible causes, **AND**
- The treatment must be given concomitantly with best supportive care for this condition.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA, **AND**
- Must be treated in a treatment centre that is each of: (i) recognised in the management of SMA, (ii) accredited in the use of this gene technology by the relevant authority, (iii) will(has) source(d) this product from an accredited supplier, as specified in the administrative notes to this listing, **AND**
- Patient must be undergoing treatment with this pharmaceutical benefit once only in a lifetime, **AND**
- Patient must not be undergoing treatment with this pharmaceutical benefit through this listing where prior treatment has occurred with any of: (i) nusinersen, (ii) risdiplam.

Population criteria:

- Patient must be no older than 9 months of age, **AND**
- Patient must have symptomatic Type 1 SMA; OR
- Patient must have pre-symptomatic SMA.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Prescribing Instructions:

In the relevant PBS Authority Application form, specify the following:

- the SMA type being treated: symptomatic Type 1 SMA, or, pre-symptomatic SMA;
- for Type 1 SMA, the signs/symptoms that the patient has experienced, together with the patient's age at the onset of these signs/symptoms.

State the weight of the patient in kilograms and request the appropriate product pack presentation with respect to the mix of 5.5 mL and 8.3 mL vials.

Confirm that genetic testing has been completed to demonstrate the following in support of an SMA diagnosis:

- 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; or
- deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variance in the remaining single copy of the SMN1 gene.

If the condition is pre-symptomatic SMA, confirm that there is genetic test finding that substantiates the number of SMN2 gene copies determined by quantitative polymerase chain reaction (qPCR) or multiple ligation dependent probe amplification (MLPA).

Quote the date, pathology provider name and any unique identifying serial number/code that links the genetic test result to the patient.

Defined signs and symptoms of type I SMA are:

- Onset before 6 months of age; and
- Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- Proximal weakness; or
- Hypotonia; or
- Absence of deep tendon reflexes; or
- Failure to gain weight appropriate for age; or
- Any active chronic neurogenic changes; or
- A compound muscle action potential below normative values for an age-matched child.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Use occurring after treatment with at least one disease modifying therapy for this condition (i.e. switching from nusinersen/risdiplam to onasemnogene abeparvovec)

Clinical criteria:

- The treatment must be given concomitantly with best supportive care for this condition, **AND**
- The treatment must not be a PBS-subsidised benefit where the condition has progressed to a point where invasive permanent assisted ventilation (i.e. ventilation via tracheostomy tube for at least 16 hours per day) is required in the absence of potentially reversible causes.

Treatment criteria:

- Patient must be undergoing treatment with this pharmaceutical benefit following prior PBS-subsidised treatment with at least one other disease modifying therapy for this condition, **AND**
- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA, **AND**
- Must be treated in a treatment centre that is each of: (i) recognised in the management of SMA, (ii) accredited in the use of this gene technology by the relevant authority, (iii) will(has) source(d) this product from an accredited supplier, as specified in the administrative notes to this listing, **AND**
- Patient must be undergoing treatment with this pharmaceutical benefit once only in a lifetime, **AND**

- Patient must be undergoing treatment with this pharmaceutical benefit with the intent that treatment with the replaced disease modifying agent is/has ceased.

Population criteria:

- Patient must be no older than 9 months of age, **AND**
- Patient must have symptomatic Type 1 SMA; OR
- Patient must have pre-symptomatic SMA.

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Do not resubmit previously submitted documentation concerning the diagnosis and type of SMA.

Confirm that a previous PBS authority application has been approved for one of the following:

- (i) Symptomatic Type 1 SMA; or
- (ii) Pre-symptomatic SMA.

State the weight of the patient in kilograms and request the appropriate product pack presentation with respect to the mix of 5.5 mL and 8.3 mL vials.

Adhere to any Product Information or local treatment guidelines with respect to treatment-free ('wash out') periods prior to administering this benefit.

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 2 x 8.3 mL vials

12957Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 8.3 mL vials], 1 pack

12958B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [3 x 8.3 mL vials], 1 pack

12940C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [4 x 8.3 mL vials], 1 pack

12950N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5 x 8.3 mL vials], 1 pack

12963G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [6 x 8.3 mL vials], 1 pack

12941D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [7 x 8.3 mL vials], 1 pack

12952Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 5.5 mL vials] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [8.3 mL vial], 1 pack

12955W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [2 x 8.3 mL vials], 1 pack

12951P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 3 x 8.3 mL vials

12961E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 4 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12971Q	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [3 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12977B	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [4 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12942E	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 5 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12962F	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 6 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12973T	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 7 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12954T	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12976Y	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [6 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12975X	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 8 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12974W	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection, 9 x 8.3 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12949M	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [7 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12956X	‡1	..	..	2527773.87	Zolgensma [NV]

onasemnogene abeparvovec 20 trillion vector genomes/mL injection [5.5 mL vial] (&) onasemnogene abeparvovec 20 trillion vector genomes/mL injection [8 x 8.3 mL vials], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
12964H	‡1	..	..	2527773.87	Zolgensma [NV]

■ RISDIPLAM

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic Type I, II or IIIa spinal muscular atrophy (SMA)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must have experienced at least two of the defined signs and symptoms of SMA type I, II or IIIa prior to 3 years of age, **AND**
- The treatment must be given concomitantly with best supportive care for this condition, **AND**
- The treatment must not be in combination with PBS-subsidised treatment with nusinersen for this condition, **AND**
- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic.

Clinical criteria:

- Patient must be untreated with gene therapy.

Population criteria:

- Patient must be 18 years of age or under.

Defined signs and symptoms of type I SMA are:

- i) Onset before 6 months of age; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Hypotonia; or
- v) Absence of deep tendon reflexes; or
- vi) Failure to gain weight appropriate for age; or
- vii) Any active chronic neurogenic changes; or
- viii) A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type II SMA are:

- i) Onset between 6 and 18 months; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Weakness in trunk righting/derotation; or
- v) Hypotonia; or
- vi) Absence of deep tendon reflexes; or
- vii) Failure to gain weight appropriate for age; or
- viii) Any active chronic neurogenic changes; or
- ix) A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type IIIa SMA are:

- i) Onset between 18 months and 3 years of age; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Hypotonia; or
- v) Absence of deep tendon reflexes; or
- vi) Failure to gain weight appropriate for age; or
- vii) Any active chronic neurogenic changes; or
- viii) A compound muscle action potential below normative values for an age-matched child.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

Application for authorisation of initial treatment must be in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Spinal muscular atrophy PBS Authority Application Form which includes the following:
 - i) specification of SMA type (I, II or IIIa); and
 - ii) sign(s) and symptom(s) that the patient has experienced; and
 - iii) patient's age at the onset of sign(s) and symptom(s).

The approved Product Information recommended dosing is as follows:

- (i) 16 days to less than 2 months of age: 0.15 mg/kg
- (ii) 2 months to less than 2 years of age: 0.20 mg/kg
- (iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg
- (iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

1 unit where (i) applies;

2 units where (ii) applies;

3 units where (iii) applies;
3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

12614X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	10841.89	Evrysdi [RO]

▪ **RISDIPLAM**

Note The maximum quantity of drug to be subsidised per dispensing, as well as the number of repeat prescriptions is to be as follows:

Patient weight greater than 19 kg: up to 3 units per dispensing, with up to 5 repeat prescriptions

Patient weight between 17 kg to 19 kg: up to 3 units per dispensing, with up to 4 repeat prescriptions

Patient weight between 13 kg to 17 kg: up to 2 units per dispensing, with up to 5 repeat prescriptions

Patient weight between 10 kg up to 13 kg: up to 2 units per dispensing, with up to 4 repeat prescriptions

Patient weight less than 10 kg: up to 1 unit per dispensing, with up to 5 repeat prescriptions

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment with this drug of either symptomatic Type I, II or IIIa SMA, or, pre-symptomatic SMA

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition; OR
- Patient must be eligible for continuing PBS-subsidised treatment with nusinersen for this condition, **AND**
- The treatment must not be in combination with PBS-subsidised treatment with nusinersen for this condition, **AND**
- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug, **AND**
- The treatment must be given concomitantly with best supportive care for this condition.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, **AND**
- Patient must not be undergoing treatment through this 'Continuing treatment' listing where the most recent PBS authority approval for this PBS-indication has been for gene therapy.

Population criteria:

- Patient must have been 18 years of age or younger at the time of initial treatment with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

In a patient who wishes to switch from PBS-subsidised nusinersen to PBS-subsidised risdiplam for this condition a wash out period may be required.

The quantity of drug and number of repeat prescriptions prescribed is to be in accordance with the relevant 'Note' attached to this listing.

The approved Product Information recommended dosing is as follows:

(i) 16 days to less than 2 months of age: 0.15 mg/kg

(ii) 2 months to less than 2 years of age: 0.20 mg/kg

(iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg

(iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

1 unit where (i) applies;

2 units where (ii) applies;

3 units where (iii) applies;

3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

12606L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	5	..	10841.89	Evrysdi [RO]

▪ **RISDIPLAM**

Note For the next authority application after this application, continue treatment through the 'Treatment phase: Continuing treatment' under the indication: 'Symptomatic Type I, II or IIIa spinal muscular atrophy (SMA)'.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Initial treatment occurring after onasemnogene abeparvovec therapy in a patient with Type 1 SMA

Clinical criteria:

- Patient must have experienced a regression in a developmental state listed below (see 'Definition') despite treatment with gene therapy - confirm that this: (i) not due to an acute concomitant illness; (ii) not due to non-compliance to best-supportive care, (iii) apparent for at least 3 months, (iv) verified by another clinician in the treatment team - state the full name of this clinician plus their profession (e.g. medical practitioner, nurse, physiotherapist; this is not an exhaustive list of examples), **AND**
- The treatment must not be a PBS-subsidised benefit where the condition has progressed to a point where invasive permanent assisted ventilation (i.e. ventilation via tracheostomy tube for at least 16 hours per day) is required in the absence of potentially reversible causes, **AND**
- The treatment must be given concomitantly with best supportive care for this condition, **AND**
- The treatment must not be in combination with PBS-subsidised treatment with nusinersen for this condition.

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic, **AND**
- Patient must be undergoing treatment under this Treatment phase listing once only - for continuing treatment beyond this authority application, refer to the drug's relevant 'Continuing treatment' listing for the patient's SMA type.

Population criteria:

- Patient must have a prior authority approval for any drug PBS-listed for symptomatic Type 1 SMA, with at least one approval having been for gene therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Do not resubmit previously submitted documentation concerning the diagnosis and type of SMA.

Confirm that a previous PBS authority application has been approved for symptomatic Type 1 SMA.

Definition:

Various childhood developmental states (1 to 9) are listed below, some followed by further observations (a up to d). Where at least one developmental state/observation is no longer present, that developmental state has regressed.

0. Absence of developmental states (1 to 9) listed below:

1. Rolls from side to side on back;
2. Child holds head erect for at least 3 seconds unsupported;
3. Sitting, but with assistance;
4. Sitting without assistance:
 - (a) Child sits up straight with the head erect for at least 10 seconds;
 - (b) Child does not use arms or hands to balance body or support position.
5. Hands and knees crawling:
 - (a) Child alternately moves forward or backwards on hands and knees;
 - (b) The stomach does not touch the supporting surface;
 - (c) There are continuous and consecutive movements at least 3 in a row.

6. Standing with assistance:

- (a) Child stands in upright position on both feet, holding onto a stable object (e.g. furniture) with both hands and without leaning on object;
- (b) The body does not touch the stable object, and the legs support most of the body weight;
- (c) Child thus stands with assistance for at least 10 seconds.

7. Standing alone:

- (a) Child stands in upright position on both feet (not on the toes) with the back straight;
- (b) The leg supports 100% of the child's weight;
- (c) There is no contact with a person or object;
- (d) Child stands alone for at least 10 seconds.

8. Walking with assistance:

- (a) Child is in an upright position with the back straight;
- (b) Child makes sideways or forced steps by holding onto a stable object (e.g. furniture) with 1 or both hands;
- (c) One leg moves forward while the other supports part of the body weight;
- (d) Child takes at least 5 steps in this manner.

9. Walking alone:

- (a) Child takes at least 5 steps independently in upright position with the back straight;

- (b) One leg moves forward while the other supports most of the body weight;
 (c) There is no contact with a person or object.

Confirm which developmental state has regressed by: (i) stating the overall developmental state (1 - 9) the patient was in at the time of gene therapy, or, the best developmental state achieved since gene therapy, and (ii) stating the patient's current overall developmental state (i.e. a number that is lower than stated in (i)).

Where the patient has neither regressed from a developmental state nor reached the next developmental state, PBS-subsidy of this benefit is not available.

The approved Product Information recommended dosing is as follows:

- (i) 16 days to less than 2 months of age: 0.15 mg/kg
- (ii) 2 months to less than 2 years of age: 0.20 mg/kg
- (iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg
- (iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

- 1 unit where (i) applies;
- 2 units where (ii) applies;
- 3 units where (iii) applies;
- 3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

12943F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	10841.89	Evrysdi [RO]

■ RISDIPLAM

Note Recognised hospitals in the management of SMA are Queensland Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
 Complex Drugs
 Reply Paid 9826
 HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Pre-symptomatic spinal muscular atrophy (SMA)

Treatment Phase: Initial treatment with this drug of pre-symptomatic spinal muscular atrophy (SMA)

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA.

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- The condition must have genetic confirmation that there are 1 to 2 copies of the survival motor neuron 2 (SMN2) gene, **AND**
- The condition must be pre-symptomatic, **AND**
- The treatment must be given concomitantly with best supportive care for this condition, **AND**
- Patient must be untreated with gene therapy.

Population criteria:

- Patient must be aged under 36 months prior to commencing treatment.

Application for authorisation of initial treatment must be in writing (lodged via postal service or electronic upload) and must include:

- (a) a completed authority prescription form; and
- (b) a completed Spinal muscular atrophy PBS Authority Application Form which includes the following:
 - (i) confirmation of genetic diagnosis of SMA; and
 - (ii) a copy of the results substantiating the number of SMN2 gene copies determined by quantitative polymerase chain reaction (qPCR) or multiple ligation dependent probe amplification (MLPA)

The quantity of drug and number of repeat prescriptions prescribed is to be in accordance with the relevant 'Note' attached to this listing.

The approved Product Information recommended dosing is as follows:

- (i) 16 days to less than 2 months of age: 0.15 mg/kg
- (ii) 2 months to less than 2 years of age: 0.20 mg/kg
- (iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg
- (iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

- 1 unit where (i) applies;
- 2 units where (ii) applies;
- 3 units where (iii) applies;
- 3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13655Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	10841.89	Evrysdi [RO]

▪ **RISDIPLAM**

Note The maximum quantity of drug to be subsidised per dispensing, as well as the number of repeat prescriptions is to be as follows:

- Patient weight greater than 19 kg: up to 3 units per dispensing, with up to 5 repeat prescriptions
- Patient weight between 17 kg to 19 kg: up to 3 units per dispensing, with up to 4 repeat prescriptions
- Patient weight between 13 kg to 17 kg: up to 2 units per dispensing, with up to 5 repeat prescriptions
- Patient weight between 10 kg up to 13 kg: up to 2 units per dispensing, with up to 4 repeat prescriptions
- Patient weight less than 10 kg: up to 1 unit per dispensing, with up to 5 repeat prescriptions

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment in a child or adult, but where treatment was initiated during childhood

Treatment criteria:

- Patient must be undergoing continuation of existing PBS-subsidised treatment with this drug; OR
- Patient must be undergoing a change in prescribed SMA drug to this drug - the drug treatment being replaced was a PBS benefit initiated prior to the patient's 19th birthday for SMA type IIIB/IIIC, **AND**
- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The quantity of drug and number of repeat prescriptions prescribed is to be in accordance with the relevant 'Note' attached to this listing.

The approved Product Information recommended dosing is as follows:

- (i) 16 days to less than 2 months of age: 0.15 mg/kg
- (ii) 2 months to less than 2 years of age: 0.20 mg/kg
- (iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg
- (iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

- 1 unit where (i) applies;
- 2 units where (ii) applies;
- 3 units where (iii) applies;
- 3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13631K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±3	5	..	*32525.67	Evrysdi [RO]

▪ **RISDIPLAM**

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Initial PBS-subsidised treatment with this drug in an adult who did not initiate PBS subsidy with this drug during childhood

Population criteria:

- Patient must be at least 19 years of age at the time of this authority application, but never claimed PBS subsidy for a disease modifying treatment during childhood, **AND**
- Patient must have SMA where the onset of signs/symptoms (at least one) of SMA first occurred prior to their 19th birthday (SMA symptom onset after this age will be considered type IV SMA, which is not PBS-subsidised).

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing initial PBS-subsidised treatment with this drug for untreated disease; OR
- Patient must be undergoing initial PBS-subsidised treatment, but the patient has initiated treatment via non-PBS supply (e.g. clinical trial, sponsor compassionate access), **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Signs and symptoms of spinal muscular atrophy in the context of this PBS restriction are:

- (i) Failure to meet or regression in ability to perform age-appropriate motor milestones,
- (ii) Proximal weakness,
- (iii) Hypotonia,
- (iv) Absence of deep tendon reflexes,
- (v) Failure to gain weight appropriate for age,
- (vi) Any active denervation or chronic neurogenic changes found on electromyography,
- (vii) A compound muscle action potential below normative values for an age-matched child.

In this authority application, confirm:

- (1) the patient's medical history is consistent with a diagnosis of childhood onset spinal muscular atrophy,
- (2) which of the above (i to vii) (at least 1) were present during childhood,
- (3) the age of the patient (rounded to the nearest year) when the first sign/symptom was observed.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13654P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±3	7	..	*32525.67	Evrysdi [RO]

■ RISDIPLAM

Note Literature references for various instruments measuring motor function and quality of life in the context of spinal muscular atrophy are:

Revised Upper Limb Module

Mazzone et al. 2017. Revised upper limb module for spinal muscular atrophy: Development of a new module. **Muscle & Nerve** 55(6):869-874

Hammersmith Functional Motor Scale - Expanded

Ramsey et al. 2017. Revised Hammersmith Scale for spinal muscular atrophy: A SMA specific clinical outcome assessment tool. **PLoS ONE** 12(2): e0172346. doi:10.1371/journal.pone.0172346.

6-Minute Walk Test (6MWT)

American Thoracic Society. 2002. ATS statement: Guidelines for the six-minute walk test. **American Journal of Respiratory and Critical Care Medicine** 166(1), pp 111-117

The National Heart Foundation of Australia has 6MWT test standardised instructions and recording forms located at: <https://www.heartonline.org.au/resources/documents-and-links#exercise>

SMA Health Index

Zizzi et al. 2021. The Spinal Muscular Atrophy Health Index (SMA-HI): A Novel Outcome for Measuring How a Patient Feels and Functions. **Muscle & Nerve** 63(10), pp 837-844

SMA Functional Rating Scale

Elsheikh et al. 2018. Reliability of Spinal Muscular Atrophy Functional Rating Scale (SMAFRS) in Ambulatory Adults with Spinal Muscular Atrophy. **Neurology** April (15 Supplement) P4.452

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Continuing/maintenance treatment in an adult where treatment was initiated in adulthood

Treatment criteria:

- Patient must be undergoing continuation of existing PBS-subsidised treatment with this drug; OR
- Patient must be undergoing a change in prescribed SMA drug to this drug - the drug treatment being replaced was a PBS benefit initiated after the patient's 19th birthday, **AND**
- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- The treatment must be each of: (i) occurring from week 104 onwards relative to the first administered dose, (ii) demonstrating a clinically meaningful response; OR
- The treatment must be occurring within the first 104 weeks from the first administered dose, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

Where this authority application seeks to continue treatment beyond the first 104 weeks of treatment, comprehensive assessment must be undertaken periodically and documented, involving the patient and the treating physician to establish agreement that treatment is continuing to produce a clinically meaningful response.

A clinically meaningful response is present where an improvement, stabilisation or minimal decline in symptoms has occurred as a result of this drug treatment and where there is agreement between the treating physician and patient over what constitutes improvement, stabilisation, or minimal decline.

PBS subsidy must cease if there is no agreement on whether a clinically meaningful response is present.

Undertake re-assessments for a clinically meaningful response at least every six months. Document these re-assessments in the patient's medical records.

In undertaking comprehensive assessments, where practical, a clinically meaningful response assessment encompasses the patient's motor function as assessed using an instrument like the Revised Upper Limb Module (RULM), Hammersmith Functional Motor Scale - Expanded (HFMSSE) or 6-minute walk test (6MWT), and the patient's quality of life including, but not limited to, level of independence. Quality of life may be informed by use of the SMA Health Index (SMA-HI) or SMA Functional Rating Scale (SMAFRS).

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13656R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±3	5	..	*32525.67	Evrysdi [RO]

■ RISDIPLAM

Note The maximum quantity of drug to be subsidised per dispensing, as well as the number of repeat prescriptions is to be as follows:

Patient weight greater than 19 kg: up to 3 units per dispensing, with up to 5 repeat prescriptions

Patient weight between 17 kg to 19 kg: up to 3 units per dispensing, with up to 4 repeat prescriptions

Patient weight between 13 kg to 17 kg: up to 2 units per dispensing, with up to 5 repeat prescriptions

Patient weight between 10 kg up to 13 kg: up to 2 units per dispensing, with up to 4 repeat prescriptions

Patient weight less than 10 kg: up to 1 unit per dispensing, with up to 5 repeat prescriptions

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note An outcome on the authority application is not immediate, but will follow in due course. Electronic upload is encouraged to reduce processing time.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Symptomatic type IIIB/IIIC spinal muscular atrophy (SMA)

Treatment Phase: Initial PBS-subsidised treatment with this drug in a child

Population criteria:

- Patient must be of an age that is prior to their 19th birthday at the time of this authority application, **AND**
- Patient must have SMA type III where the onset of signs/symptoms of SMA first occurred after their 3rd birthday, but before their 19th birthday (SMA type IIIB/IIIC).

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis/management of SMA; OR
- Must be treated by a medical practitioner who has been directed to prescribe this benefit by a specialist medical practitioner experienced in the diagnosis/management of SMA, **AND**
- Patient must be undergoing initial PBS-subsidised treatment with this drug for untreated disease; OR
- Patient must be undergoing initial PBS-subsidised treatment, but the patient has initiated treatment via non-PBS supply (e.g. clinical trial, sponsor compassionate access), **AND**
- Patient must be undergoing concomitant treatment with best supportive care, but this benefit is the sole PBS-subsidised disease modifying treatment.

Clinical criteria:

- The condition must have genetic confirmation of 5q homozygous deletion of the survival motor neuron 1 (SMN1) gene; OR
- The condition must have genetic confirmation of deletion of one copy of the SMN1 gene in addition to a pathogenic/likely pathogenic variant in the remaining single copy of the SMN1 gene, **AND**
- Patient must not be receiving invasive permanent assisted ventilation in the absence of a potentially reversible cause while being treated with this drug.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Signs and symptoms of spinal muscular atrophy in the context of this PBS restriction are:

- (i) Failure to meet or regression in ability to perform age-appropriate motor milestones,
- (ii) Proximal weakness,
- (iii) Hypotonia,
- (iv) Absence of deep tendon reflexes,
- (v) Any active denervation or chronic neurogenic changes found on electromyography,
- (vi) A compound muscle action potential below normative values for an age-matched child.

In this authority application, confirm:

- (1) the patient's medical history is consistent with a diagnosis of type IIIB/IIIC spinal muscular atrophy,
- (2) which of the above (i to vi) (at least 1) were present after their 3rd birthday, but before their 19th birthday,
- (3) the age of the patient (rounded to the nearest year) when the first sign/symptom was observed.

The quantity of drug and number of repeat prescriptions prescribed is to be in accordance with the relevant 'Note' attached to this listing.

The approved Product Information recommended dosing is as follows:

- (i) 16 days to less than 2 months of age: 0.15 mg/kg
- (ii) 2 months to less than 2 years of age: 0.20 mg/kg
- (iii) 2 years of age and older weighing less than 20 kg: 0.25 mg/kg
- (iv) 2 years of age and older weighing 20 kg or more: 5 mg

In this authority application, state which of (i) to (iv) above applies to the patient. Based on (i) to (iv), prescribe up to:

- 1 unit where (i) applies;
- 2 units where (ii) applies;
- 3 units where (iii) applies;
- 3 units where (iv) applies.

risdiplam 750 microgram/mL powder for oral liquid, 80 mL

13633M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	‡3	5	..	*32525.67	Evrysdi [RO]

Growth Hormone Program

▪ SOMATROPIN

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe growth hormone deficiency

Treatment Phase: Initial treatment of late onset growth hormone deficiency

Treatment criteria:

- Must be treated by an endocrinologist.

Clinical criteria:

- Patient must have onset of growth hormone deficiency secondary to organic hypothalamic or pituitary disease diagnosed at chronological age of 18 years or older; OR
- Patient must have onset of growth hormone deficiency diagnosed after skeletal maturity (bone age greater than or equal to 15.5 years in males or 13.5 years in females) and before chronological age of 18 years, **AND**
- Patient must have a diagnostic insulin tolerance test with maximum serum growth hormone (GH) less than 2.5 micrograms per litre; OR
- Patient must have a diagnostic arginine infusion test with maximum serum GH less than 0.4 micrograms per litre; OR
- Patient must have a diagnostic glucagon provocation test with maximum serum GH less than 3 micrograms per litre.

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Severe Growth Hormone Deficiency supporting information form; AND
3. Results of the growth hormone stimulation testing, including the date of testing, the type of test performed, the peak growth hormone concentration, and laboratory reference range for age/gender.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe growth hormone deficiency

Treatment Phase: Continuing treatment in a person with a mature skeleton or aged 18 years or older

Treatment criteria:

- Must be treated by an endocrinologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy with this drug for this condition under an initial treatment restriction applying to a documented childhood onset growth hormone deficiency due to a congenital, genetic or structural cause in a patient with a mature skeleton; OR
- Patient must have previously received PBS-subsidised therapy with this drug for this condition under an initial treatment restriction applying to late onset of growth hormone deficiency secondary to organic hypothalamic or pituitary disease in a patient with chronological age of 18 years or older; OR
- Patient must have previously received PBS-subsidised therapy with this drug for this condition under an initial treatment restriction applying to late onset of growth hormone deficiency diagnosed after skeletal maturity (bone age greater than or equal to 15.5 years in males or 13.5 years in females) and before chronological age of 18 years.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Severe growth hormone deficiency

Treatment Phase: Initial treatment of childhood onset growth hormone deficiency in a patient who has received PBS-subsidised treatment as a child

Treatment criteria:

- Must be treated by an endocrinologist.

Clinical criteria:

- Patient must have a documented childhood onset growth hormone deficiency due to a congenital, genetic or structural cause, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition as a child.

Population criteria:

- Patient must have a mature skeleton.

Somatropin is not PBS-subsidised for patients with Prader-Willi syndrome aged 18 years or older without a documented childhood onset Growth Hormone Deficiency.

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Severe Growth Hormone Deficiency supporting information form.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe growth hormone deficiency

Treatment Phase: Initial treatment of childhood onset growth hormone deficiency in a patient who has received non-PBS subsidised treatment as a child

Treatment criteria:

- Must be treated by an endocrinologist.

Clinical criteria:

- Patient must have a documented childhood onset growth hormone deficiency due to a congenital, genetic or structural cause, **AND**
- Patient must have previously received non-PBS subsidised treatment with this drug for this condition as a child, **AND**
- Patient must have current or historical evidence of an insulin tolerance test with maximum serum growth hormone (GH) less than 2.5 micrograms per litre; OR
- Patient must have current or historical evidence of an arginine infusion test with maximum serum GH less than 0.4 micrograms per litre; OR
- Patient must have current or historical evidence of a glucagon provocation test with maximum serum GH less than 3 micrograms per litre.

Population criteria:

- Patient must have a mature skeleton.

Somatropin is not PBS-subsidised for patients with Prader-Willi syndrome aged 18 years or older without a documented childhood onset Growth Hormone Deficiency.

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Severe Growth Hormone Deficiency supporting information form; AND
3. Results of the growth hormone stimulation testing, including the date of testing, the type of test performed, the peak growth hormone concentration, and laboratory reference range for age/gender.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

11895C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	165.94	30.00	Norditropin FlexPro [NO]

somatropin 12 mg injection [1 chamber] (&) inert substance diluent [1 mL chamber], 1 dual chamber pen device

11495B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	386.53	30.00	Genotropin GoQuick [PF]

somatropin 5 mg injection [1 chamber] (&) inert substance diluent [1 mL chamber], 1 dual chamber pen device

11493X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	165.94	30.00	Genotropin GoQuick [PF]

somatropin 10 mg/2 mL injection, 2 mL cartridge

11650E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	323.51	30.00	NutropinAq [IS]