



Australian Government

**Department of Health
and Aged Care**



Schedule of Pharmaceutical Benefits

Efficient Funding of Chemotherapy

Effective 1 June 2024



Contents

Summary of Changes	3
About the Supplement	5
Symbols used in the Efficient Funding of Chemotherapy supplement	5
Remuneration arrangements.....	5
Pharmaceutical Benefits Schedules	6
Chemotherapy items for Private Hospital use	7
Chemotherapy items for Public Hospital use.....	81
Related Pharmaceutical Benefits for Public Hospital use.....	155
Index of Manufacturers' Code	165
Generic/Proprietary Index	167

Summary of Changes

These changes to the Schedule of Pharmaceutical Benefits are effective from 1 June 2024. The Schedule is updated on the first day of each month and is available on the internet at www.pbs.gov.au.

Efficient Funding of Chemotherapy (Private Hospital)

Additions

Addition – Brand

12219D *BORTEZOMIB EUGIA, YG* – **BORTEZOMIB**, bortezomib 3.5 mg injection, 1 vial

Deletions

Deletion – Brand

12219D *Velcade, JC* – **BORTEZOMIB**, bortezomib 1 mg injection, 1 vial

12219D *Velcade, JC* – **BORTEZOMIB**, bortezomib 3 mg injection, 1 vial

12219D *Velcade, JC* – **BORTEZOMIB**, bortezomib 3.5 mg injection, 1 vial

Deletion – Note

10748Q **NIVOLUMAB**,
nivolumab 40 mg/4 mL injection, 4 mL vial (*Opdivo*);
nivolumab 100 mg/10 mL injection, 10 mL vial (*Opdivo*)

11532Y **NIVOLUMAB**,
nivolumab 40 mg/4 mL injection, 4 mL vial (*Opdivo*);
nivolumab 100 mg/10 mL injection, 10 mL vial (*Opdivo*)

13267G **PEMBROLIZUMAB**,
pembrolizumab 100 mg/4 mL injection, 4 mL vial (*Keytruda*)

Deletion – Restriction

10748Q **NIVOLUMAB**,
nivolumab 40 mg/4 mL injection, 4 mL vial (*Opdivo*);
nivolumab 100 mg/10 mL injection, 10 mL vial (*Opdivo*)

11532Y **NIVOLUMAB**,
nivolumab 40 mg/4 mL injection, 4 mL vial (*Opdivo*);
nivolumab 100 mg/10 mL injection, 10 mL vial (*Opdivo*)

13267G **PEMBROLIZUMAB**,
pembrolizumab 100 mg/4 mL injection, 4 mL vial (*Keytruda*)

Efficient Funding of Chemotherapy (Public Hospital)

Additions

Addition – Brand

12227M *BORTEZOMIB EUGIA, YG* – **BORTEZOMIB**, bortezomib 3.5 mg injection, 1 vial

Deletions

Deletion – Brand

12227M *Velcade, JC* – **BORTEZOMIB**, bortezomib 1 mg injection, 1 vial

12227M *Velcade, JC* – **BORTEZOMIB**, bortezomib 3 mg injection, 1 vial

12227M *Velcade, JC* – **BORTEZOMIB**, bortezomib 3.5 mg injection, 1 vial

Deletion – Note

10745M **NIVOLUMAB**,
nivolumab 40 mg/4 mL injection, 4 mL vial (*Opdivo*);
nivolumab 100 mg/10 mL injection, 10 mL vial (*Opdivo*)

11543M **NIVOLUMAB**,
nivolumab 40 mg/4 mL injection, 4 mL vial (*Opdivo*);
nivolumab 100 mg/10 mL injection, 10 mL vial (*Opdivo*)

13254N **PEMBROLIZUMAB**,
pembrolizumab 100 mg/4 mL injection, 4 mL vial (*Keytruda*)

Deletion – Restriction

10745M **NIVOLUMAB**,
nivolumab 40 mg/4 mL injection, 4 mL vial (*Opdivo*);
nivolumab 100 mg/10 mL injection, 10 mL vial (*Opdivo*)

11543M **NIVOLUMAB**,
nivolumab 40 mg/4 mL injection, 4 mL vial (*Opdivo*);
nivolumab 100 mg/10 mL injection, 10 mL vial (*Opdivo*)

13254N **PEMBROLIZUMAB**,
pembrolizumab 100 mg/4 mL injection, 4 mL vial (*Keytruda*)

Related Pharmaceutical Benefits for Public Hospital use

Additions

Addition – Brand

5967C *Ondansetron Tablets Viatris, AL* – **ONDANSETRON**, ondansetron 4 mg tablet, 4

5858H *Ondansetron ODT Viatris, AL* – **ONDANSETRON**, ondansetron 8 mg orally disintegrating tablet, 4

5968D *Ondansetron Tablets Viatris, AL* – **ONDANSETRON**, ondansetron 8 mg tablet, 4

About the Supplement

The Schedule of Pharmaceutical Benefits – Efficient Funding of Chemotherapy supplement lists items distributed under section 100 of the National Health Act 1953.

The Supplement is published and is effective on the first day of each month. For detailed information about the prescribing and supply of chemotherapy benefits go to www.pbs.gov.au

For information about the operational aspects of the Efficient Funding of Chemotherapy, such as, claiming, authority applications and stationery supplies contact Services Australia at www.servicesaustralia.gov.au

This supplement is split into three parts:

Chemotherapy items for private hospital use. This includes items subject to the revised arrangements, ie. chemotherapy drugs administered through infusion or injection

Chemotherapy items for public hospital use. This includes items subject to the revised arrangements, ie. chemotherapy drugs administered through infusion or injection

PBS products available for private and public hospital use may be dispensed in accordance with the relevant section 100 special arrangements through community pharmacy.

Related pharmaceutical benefits for public hospital use. This includes items such as antiemetics, antinauseants, immunostimulants and detoxifying agents for antineoplastic treatment

Symbols used in the Efficient Funding of Chemotherapy supplement

*	An asterisk in the dispensed price column indicates that the manufacturer's pack does not coincide with the maximum quantity
‡	A double dagger in the maximum quantity column indicates where the maximum quantity has been determined to match the manufacturer's pack. These packs cannot be broken and the maximum quantity should be supplied and claimed
^a or ^b	Located immediately before brand names of an item indicates that the brands are equivalent for the purposes of substitution. These brands may be interchanged without differences in clinical effect

Remuneration arrangements

Fees payable per item claimed:

Section 90 Community Pharmacy (incl. section 92 approved practitioners)

- Ready Prepared Dispensing Fee (\$8.37)
- Preparation fee (\$88.62)
- Distribution fee (\$29.15)
- Diluent fee (\$5.77)

Section 94 Approved Public Hospital Authority

- Preparation fee (\$88.62)

Section 94 Approved Private Hospital Authority

- Ready Prepared Dispensing Fee (\$8.37)
- Preparation fee (\$88.62)
- Distribution fee (\$29.15) (not payable where the drug is trastuzumab)
- Diluent fee (\$5.77)

Pharmaceutical Benefits Schedules

Chemotherapy items for Private Hospital use

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	8
ANTINEOPLASTIC AGENTS	8
ALKYLATING AGENTS	8
ANTIMETABOLITES	9
PLANT ALKALOIDS AND OTHER NATURAL PRODUCTS	11
CYTOTOXIC ANTIBIOTICS AND RELATED SUBSTANCES	14
MONOCLONAL ANTIBODIES AND ANTIBODY DRUG CONJUGATES	15
OTHER ANTINEOPLASTIC AGENTS	72

■ ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

■ ANTINEOPLASTIC AGENTS

ALKYLATING AGENTS

Nitrogen mustard analogues

■ BENDAMUSTINE

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7972

Previously untreated stage III or IV mantle cell lymphoma

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be CD20 positive, **AND**
- The treatment must be in combination with rituximab, **AND**
- The condition must be previously untreated, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for induction treatment purposes only, **AND**
- Patient must not receive more than 6 cycles (12 doses) of treatment under this restriction, **AND**
- Patient must not be eligible for stem cell transplantation.

Authority required (STREAMLINED)

7943

Previously untreated stage II bulky or stage III or IV indolent non-Hodgkin's lymphoma

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be CD20 positive, **AND**
- The condition must be previously untreated, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for induction treatment purposes only, **AND**
- The treatment must be in combination with rituximab or obinutuzumab, **AND**
- The treatment must not exceed 6 cycles (12 doses) with this drug under this restriction.

Authority required (STREAMLINED)

7944

Follicular lymphoma

Treatment Phase: Re-induction treatment

Clinical criteria:

- The condition must be CD20 positive, **AND**
- The condition must be refractory to treatment with rituximab for this condition, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for re-induction treatment purposes only, **AND**
- The treatment must be in combination with obinutuzumab, **AND**
- The treatment must not exceed 6 cycles (12 doses) with this drug under this restriction.

The condition is considered rituximab-refractory if the patient experiences less than a partial response or progression of disease within 6 months after completion of a prior rituximab-containing regimen.

Injection

10763L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	11	..	*1298.43	31.60	Bendamustine Juno [JU] (bendamustine hydrochloride 100 mg injection, 1 vial) Bendamustine Juno [JU] (bendamustine hydrochloride 25 mg injection, 1 vial) Bendamustine Sandoz [SZ] (bendamustine hydrochloride 100 mg injection, 1 vial) Bendamustine Sandoz [SZ] (bendamustine hydrochloride 25 mg injection, 1 vial) Bendamustine Viatrix [AF] (bendamustine hydrochloride 100 mg injection, 1 vial) Bendamustine Viatrix [AF] (bendamustine hydrochloride 25 mg injection, 1 vial) Ribomustin [JC] (bendamustine hydrochloride 100 mg injection, 1 vial) Ribomustin [JC] (bendamustine hydrochloride 25 mg injection, 1 vial)

■ **CYCLOPHOSPHAMIDE**

Injection

7226H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	2800 mg	17	..	*188.46	31.60	CYCLOPHOSPHAMIDE-REACH [RQ] (cyclophosphamide 1 g injection, 1 vial) CYCLOPHOSPHAMIDE-REACH [RQ] (cyclophosphamide 500 mg injection, 1 vial) Endoxan [BX] (cyclophosphamide 1 g injection, 1 vial) Endoxan [BX] (cyclophosphamide 2 g injection, 1 vial)

■ **IFOSFAMIDE**

Injection

7248L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	4000 mg	19	..	*287.03	31.60	Holoxan [BX] (ifosfamide 1 g injection, 1 vial) Holoxan [BX] (ifosfamide 2 g injection, 1 vial)

ANTIMETABOLITES

Folic acid analogues

■ **METHOTREXATE**

Injection

7250N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	250 mg	5	..	*157.76	31.60	DBL Methotrexate [PF] (methotrexate 1 g/10 mL injection, 10 mL vial) DBL Methotrexate [PF] (methotrexate 5 mg/2 mL injection, 5 x 2 mL vials) DBL Methotrexate [PF] (methotrexate 50 mg/2 mL injection, 5 x 2 mL vials) DBL Methotrexate [PF] (methotrexate 500 mg/20 mL injection, 20 mL vial) Methotrexate Accord [OD] (methotrexate 1 g/10 mL injection, 10 mL vial) Methotrexate Ebewe [SZ] (methotrexate 5 g/50 mL injection, 50 mL vial)

■ **METHOTREXATE**

Restricted benefit

Patients receiving treatment with a high dose regimen

Injection

7251P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	20000 mg	*895.87	31.60	DBL Methotrexate [PF] (methotrexate 1 g/10 mL injection, 10 mL vial) DBL Methotrexate [PF] (methotrexate 5 mg/2 mL injection, 5 x 2 mL vials) DBL Methotrexate [PF] (methotrexate 50 mg/2 mL injection, 5 x 2 mL vials) DBL Methotrexate [PF] (methotrexate 500 mg/20 mL injection, 20 mL vial) Methotrexate Accord [OD] (methotrexate 1 g/10 mL injection, 10 mL vial) Methotrexate Ebewe [SZ] (methotrexate 5 g/50 mL injection, 50 mL vial)

■ **PEMETREXED**

Injection

7255W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1100 mg	5	..	*191.09	31.60	Pemetrexed Accord [OD] (pemetrexed 1 g injection, 1 vial) Pemetrexed Accord [OD] (pemetrexed 100 mg injection, 1 vial) Pemetrexed Accord [OD] (pemetrexed 500 mg injection, 1 vial) Pemetrexed APOTEX [TX] (pemetrexed 500 mg injection, 1 vial) Pemetrexed Ever Pharma [IT] (pemetrexed 1 g/40 mL injection, 40 mL vial) Pemetrexed Ever Pharma [IT] (pemetrexed 100 mg/4 mL injection, 4 mL vial) Pemetrexed Ever Pharma [IT] (pemetrexed 500 mg/20 mL injection, 20 mL vial) Pemetrexed SUN [RA] (pemetrexed 1 g injection, 1 vial)

Pemetrexed SUN [RA] (pemetrexed 100 mg injection, 1 vial)
Pemetrexed SUN [RA] (pemetrexed 500 mg injection, 1 vial)

■ PRALATREXATE

Note No increase in the maximum number of repeats may be authorised.

Authority required

Relapsed or chemotherapy refractory Peripheral T-cell Lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be relapsed or chemotherapy refractory, **AND**
- Patient must have undergone appropriate prior front-line curative intent chemotherapy.

Injection

11271F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	80 mg	5	..	*4331.91	31.60	Folotyn [MF] (pralatrexate 20 mg/mL injection, 1 mL vial)

■ PRALATREXATE

Note No increase in the maximum number of repeats may be authorised.

Authority required

Relapsed or chemotherapy refractory Peripheral T-cell Lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be relapsed or chemotherapy refractory, **AND**
- Patient must not develop progressive disease whilst receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

Injection

11278N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	80 mg	11	..	*4331.91	31.60	Folotyn [MF] (pralatrexate 20 mg/mL injection, 1 mL vial)

■ RALTITREXED

Injection

7256X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	7 mg	8	..	*1189.35	31.60	Tomudex [PF] (raltitrexed 2 mg injection, 1 vial)

Purine analogues

■ CLADRIBINE

Authority required (STREAMLINED)

6265

Hairy cell leukaemia

Injection

7225G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	17 mg	6	..	*952.49	31.60	Leustatin [IX] (cladribine 10 mg/10 mL injection, 10 mL vial) Litak [AF] (cladribine 10 mg/5 mL injection, 5 mL vial)

■ FLUDARABINE

Note Pharmaceutical benefits that have the form fludarabine phosphate 50 mg injection and pharmaceutical benefits that have the form fludarabine phosphate 50 mg/2 mL injection are equivalent for the purposes of substitution.

Injection

7233Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	55 mg	29	..	*196.49	31.60	Fludarabine Ebewe [SZ] (fludarabine phosphate 50 mg/2 mL injection, 5 x 2 mL vials) Fludarabine Juno [JO] (fludarabine phosphate 50 mg injection, 1 vial)

Pyrimidine analogues

■ CYTARABINE

Injection

7227J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	7000 mg	15	..	*941.11	31.60	Pfizer Australia Pty Ltd [PF] (cytarabine 100 mg/5 mL injection, 5 x 5 mL vials)

■ FLUOROURACIL

Restricted benefit

Patients requiring administration of fluorouracil by intravenous infusion

Injection

7234R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	5500 mg	11	..	*155.07	31.60	Fluorouracil Accord [OC] (fluorouracil 1 g/20 mL injection, 20 mL vial) Fluorouracil Accord [OC] (fluorouracil 2.5 g/50 mL injection, 50 mL vial) Fluorouracil Accord [OC] (fluorouracil 5 g/100 mL injection, 100 mL vial) Fluorouracil Accord [OC] (fluorouracil 500 mg/10 mL injection, 10 mL vial) Fluorouracil Ebewe [SZ] (fluorouracil 1 g/20 mL injection, 20 mL vial) Fluorouracil Ebewe [SZ] (fluorouracil 5 g/100 mL injection, 100 mL vial)

▪ **FLUOROURACIL**

Restricted benefit

Patients requiring administration of fluorouracil by intravenous injection

Injection

7239B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	23	..	*137.83	31.60	Fluorouracil Accord [OC] (fluorouracil 1 g/20 mL injection, 20 mL vial) Fluorouracil Accord [OC] (fluorouracil 2.5 g/50 mL injection, 50 mL vial) Fluorouracil Accord [OC] (fluorouracil 5 g/100 mL injection, 100 mL vial) Fluorouracil Accord [OC] (fluorouracil 500 mg/10 mL injection, 10 mL vial) Fluorouracil Ebewe [SZ] (fluorouracil 1 g/20 mL injection, 20 mL vial) Fluorouracil Ebewe [SZ] (fluorouracil 5 g/100 mL injection, 100 mL vial)

▪ **GEMCITABINE**

Caution Pharmaceutical benefits containing gemcitabine may have different concentrations.

Injection

7246J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3000 mg	17	..	*195.84	31.60	DBL Gemcitabine Injection [PF] (gemcitabine 1 g/26.3 mL injection, 26.3 mL vial) DBL Gemcitabine Injection [PF] (gemcitabine 2 g/52.6 mL injection, 52.6 mL vial)

PLANT ALKALOIDS AND OTHER NATURAL PRODUCTS

Vinca alkaloids and analogues

▪ **VINBLASTINE**

Injection

7261E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	20 mg	17	..	*206.27	31.60	DBL Vinblastine [PF] (vinblastine sulfate 10 mg/10 mL injection, 5 x 10 mL vials)

▪ **VINCRIStINE**

Injection

7262F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	2 mg	7	..	*149.99	31.60	DBL Vincristine Sulfate [PF] (vincristine sulfate 1 mg/mL injection, 5 x 1 mL vials)

▪ **VINOReLBINE**

Injection

7263G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	70 mg	7	..	*194.67	31.60	Navelbine [FB] (vinorelbine 50 mg/5 mL injection, 5 mL vial) Vinorelbine Ebewe [SZ] (vinorelbine 10 mg/mL injection, 1 mL vial) Vinorelbine Ebewe [SZ] (vinorelbine 50 mg/5 mL injection, 5 mL vial)

Podophyllotoxin derivatives

■ ETOPOSIDE

Injection

7237X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	440 mg	14	..	*328.11	31.60	Etopophos [LM] (etoposide phosphate 1.136 g (etoposide 1 g) injection, 1 vial) Etoposide Ebewe [SZ] (etoposide 100 mg/5 mL injection, 5 x 5 mL vials)

Taxanes

■ CABAZITAXEL

Note Where the term 'novel hormonal drug' appears in this restriction, it refers to: (i) abiraterone, (ii) abiraterone and methylprednisolone, (iii) apalutamide, (iv) darolutamide, (v) enzalutamide.

Authority required (STREAMLINED)

13207

Castration resistant metastatic carcinoma of the prostate

Clinical criteria:

- The treatment must be in combination with prednisone or prednisolone, **AND**
- The condition must be resistant to treatment with docetaxel; OR
- Patient must have a documented intolerance necessitating permanent treatment withdrawal or a contraindication to docetaxel, **AND**
- The treatment must not be used in combination with a novel hormonal drug, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must not receive PBS-subsidised cabazitaxel if progressive disease develops while on cabazitaxel.

Injection

7236W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	55 mg	5	..	*221.82	31.60	Cabazitaxel Accord [OC] (cabazitaxel 60 mg/3 mL injection, 3 mL vial) Cabazitaxel Ever Pharma [IT] (cabazitaxel 60 mg/6 mL injection, 6 mL vial) Cabazitaxel Juno [JU] (cabazitaxel 60 mg/1.5 mL injection [1.5 mL vial] (&) inert substance diluent [4.5 mL vial], 1 pack) MSN Cabazitaxel [RQ] (cabazitaxel 60 mg/1.5 mL injection [1.5 mL vial] (&) inert substance diluent [4.5 mL vial], 1 pack)

■ DOCETAXEL

Note Pharmaceutical benefits that have the forms docetaxel solution concentrate for I.V. infusion 80 mg in 4 mL and docetaxel solution concentrate for I.V. infusion 80 mg in 8 mL are equivalent for the purposes of substitution.

Note Pharmaceutical benefits that have the forms docetaxel solution concentrate for I.V. infusion 160 mg in 8 mL and docetaxel solution concentrate for I.V. infusion 160 mg in 16 mL are equivalent for the purposes of substitution.

Injection

10158P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	250 mg	5	..	*199.55	31.60	DBL Docetaxel Concentrated Injection [PF] (docetaxel 160 mg/16 mL injection, 16 mL vial) DBL Docetaxel Concentrated Injection [PF] (docetaxel 80 mg/8 mL injection, 8 mL vial) Docetaxel Accord [OC] (docetaxel 160 mg/8 mL injection, 8 mL vial) Docetaxel Accord [OC] (docetaxel 80 mg/4 mL injection, 4 mL vial)

■ NANOPARTICLE ALBUMIN-BOUND PACLITAXEL

Authority required (STREAMLINED)

6106

Metastatic breast cancer

Authority required (STREAMLINED)

6119

HER2 positive breast cancer

Injection

7270P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	580 mg	5	..	*2220.39	31.60	Abraxane [TS] (paclitaxel (as nanoparticle albumin-bound) 100 mg injection, 1 vial)

■ NANOPARTICLE ALBUMIN-BOUND PACLITAXEL

Note Special Pricing Arrangements apply.

Note Not for use as neoadjuvant or adjuvant therapy.

Authority required (STREAMLINED)

4657

Stage IV (metastatic) adenocarcinoma of the pancreas

Clinical criteria:

- The treatment must be in combination with gemcitabine, **AND**
- The condition must not have been treated previously with PBS-subsidised therapy, **AND**
- Patient must have an Eastern Cooperative Oncology Group (ECOG) performance status score of 2 or less.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Injection

10150F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	275 mg	11	..	*1176.15	31.60	Abraxane [TS] (paclitaxel (as nanoparticle albumin-bound) 100 mg injection, 1 vial)

▪ **PACLITAXEL**

Injection

7254T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	450 mg	3	..	*206.49	31.60	Paclitaxel Accord [OC] (paclitaxel 300 mg/50 mL injection, 50 mL vial) Paclitaxel Ebewe [SZ] (paclitaxel 300 mg/50 mL injection, 50 mL vial)

Topoisomerase 1 (TOP1) inhibitors

▪ **IRINOTECAN**

Note In first-line usage, effectiveness and tolerance may be improved when irinotecan is combined with an infusional 5-fluorouracil regimen.

Injection

7249M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	800 mg	11	..	*202.91	31.60	Irinotecan Accord [OC] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial) Irinotecan Accord [OC] (irinotecan hydrochloride trihydrate 500 mg/25 mL injection, 25 mL vial) Irinotecan Alphapharm [AF] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial) Irinotecan Alphapharm [AF] (irinotecan hydrochloride trihydrate 500 mg/25 mL injection, 25 mL vial) IRINOTECAN BAXTER [BX] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial) Omegapharm Irinotecan [OE] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial) Omegapharm Irinotecan [OE] (irinotecan hydrochloride trihydrate 40 mg/2 mL injection, 2 mL vial)

▪ **TOPOTECAN**

Injection

7260D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3500 mcg	17	..	*164.44	31.60	Hycamtin [SZ] (topotecan 4 mg injection, 5 vials) Topotecan Accord [OC] (topotecan 4 mg/4 mL injection, 5 x 4 mL vials)

Other plant alkaloids and natural products

▪ **TRABECTEDIN**

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14196

Advanced (unresectable and/or metastatic) leiomyosarcoma or liposarcoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have an ECOG performance status of 2 or less, **AND**
- Patient must have received prior chemotherapy treatment including an anthracycline, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The condition must be one of the following subtypes for patients with liposarcoma: (i) dedifferentiated, (ii) myxoid, (iii) round-cell, (iv) pleomorphic.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Injection

13348M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3250 mcg	3	..	*3414.11	31.60	Yondelis [ZL] (trabectedin 1 mg injection, 1 vial)

▪ TRABECTEDIN

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14197

Advanced (unresectable and/or metastatic) leiomyosarcoma or liposarcoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Authority required (STREAMLINED)

14188

Advanced (unresectable and/or metastatic) leiomyosarcoma or liposarcoma

Treatment Phase: Transitioning from non-PBS to PBS-subsidised treatment - Grandfather arrangements

Clinical criteria:

- Patient must have been receiving treatment with this drug for this condition prior to 1 August 2023, **AND**
- Patient must have had a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 2 at the time non-PBS supply was initiated, **AND**
- Patient must have received chemotherapy treatment including an anthracycline, prior to initiating non-PBS-subsidised treatment, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The condition must be one of the following subtypes for patients with liposarcoma: (i) dedifferentiated, (ii) myxoid, (iii) round-cell, (iv) pleomorphic.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Injection

13344H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3250 mcg	7	..	*3414.11	31.60	Yondelis [ZL] (trabectedin 1 mg injection, 1 vial)

CYTOTOXIC ANTIBIOTICS AND RELATED SUBSTANCES

Anthracyclines and related substances

▪ DOXORUBICIN

Injection/intravesical

7229L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	135 mg	11	..	*169.14	31.60	Adriamycin [PF] (doxorubicin hydrochloride 200 mg/100 mL injection, 100 mL vial) Adriamycin [PF] (doxorubicin hydrochloride 50 mg/25 mL injection, 25 mL vial) Doxorubicin ACC [OC] (doxorubicin hydrochloride 200 mg/100 mL injection, 100 mL vial)

▪ DOXORUBICIN HYDROCHLORIDE (AS PEGYLATED LIPOSOMAL)

Injection

7230M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	100 mg	5	..	*800.27	31.60	Caelyx [BX] (doxorubicin hydrochloride (as pegylated liposomal) 20 mg/10 mL injection, 10 mL vial) Caelyx [BX] (doxorubicin hydrochloride (as pegylated liposomal) 50 mg/25 mL injection, 25 mL vial) Liposomal Doxorubicin SUN [RA] (doxorubicin hydrochloride (as pegylated liposomal) 20 mg/10 mL injection, 10 mL vial) Liposomal Doxorubicin SUN [RA] (doxorubicin hydrochloride (as pegylated liposomal) 50 mg/25 mL injection, 25 mL vial)

▪ EPIRUBICIN

Injection/intravesical

7231N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	220 mg	5	..	*312.31	31.60	Epirubicin Accord [OC] (epirubicin hydrochloride 200 mg/100 mL injection, 100 mL vial)

▪ IDARUBICIN

Restricted benefit

Acute myelogenous leukaemia (AML)

Injection

7247K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	30 mg	5	..	*313.89	31.60	Zavedos Solution [PF] (idarubicin hydrochloride 5 mg/5 mL injection, 5 mL vial)

▪ **MITOZANTRONE**

Injection

7252Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	30 mg	5	..	*225.81	31.60	Mitozantrone Ebewe [SZ] (mitozantrone 20 mg/10 mL injection, 10 mL vial) Onkotrone [BX] (mitozantrone 25 mg/12.5 mL injection, 12.5 mL vial)

Other cytotoxic antibiotics

▪ **BLEOMYCIN**

Restricted benefit

Germ cell neoplasms

Restricted benefit

Lymphoma

Injection

7244G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	30000 iu	11	..	*213.03	31.60	DBL Bleomycin Sulfate [PF] (bleomycin sulfate 15 000 international units injection, 1 vial)

MONOCLONAL ANTIBODIES AND ANTIBODY DRUG CONJUGATES

CD20 (Clusters of Differentiation 20) inhibitors

▪ **OBINUTUZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Stage II bulky or Stage III/IV follicular lymphoma

Treatment Phase: Maintenance therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug under the previously untreated initial restriction, **AND**
- The condition must be CD20 positive, **AND**
- Patient must have demonstrated a partial or complete response to PBS subsidised induction treatment with this drug for this condition, **AND**
- The treatment must be maintenance therapy, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed 12 doses or 2 years duration of treatment, whichever comes first, under this restriction, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Injection

11455X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	5	..	*4720.79	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ **OBINUTUZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Stage II bulky or Stage III/IV follicular lymphoma

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be CD20 positive, **AND**
- The condition must be previously untreated, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for induction treatment purposes only, **AND**
- The treatment must be in combination with chemotherapy, **AND**
- The treatment must not exceed 10 doses for induction treatment with this drug for this condition.

A patient may only qualify for PBS-subsidised initiation treatment once in a lifetime under:

- i) the previously untreated induction treatment restriction; or
- ii) the rituximab-refractory re-induction restriction.

Injection

11456Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	9	..	*4720.79	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

OBINUTUZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Follicular lymphoma

Treatment Phase: Re-induction treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised obinutuzumab, **AND**
- The condition must be CD20 positive, **AND**
- The condition must be refractory to treatment with rituximab for this condition, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for re-induction treatment purposes only, **AND**
- The treatment must be in combination with bendamustine, **AND**
- The treatment must not exceed 8 doses for re-induction treatment with this drug for this condition.

The condition is considered rituximab-refractory if the patient experiences less than a partial response or progression of disease within 6 months after completion of a prior rituximab-containing regimen.

A patient may only qualify for PBS-subsidised initiation treatment once in a lifetime under:

- i) the previously untreated induction treatment restriction; or
- ii) the rituximab-refractory re-induction restriction.

Injection

11460E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	7	..	*4720.79	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

OBINUTUZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Follicular lymphoma

Treatment Phase: Maintenance therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug under the rituximab refractory initial restriction, **AND**
- The condition must be CD20 positive, **AND**
- The condition must have been refractory to treatment with rituximab, **AND**
- Patient must have demonstrated a partial or complete response to PBS-subsidised re-induction treatment with this drug for this condition, **AND**
- The treatment must be maintenance therapy, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed 12 doses or 2 years duration of treatment, whichever comes first, under this restriction, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Injection

11473W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	5	..	*4720.79	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

OBINUTUZUMAB

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

11015

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: For combination use with venetoclax treatment cycles 1 to 6 inclusive in first-line therapy

Clinical criteria:

- The condition must be untreated, **AND**
- The treatment must be in combination with PBS-subsidised venetoclax.

Injection

12193R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	8	..	*4720.79	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ **OBINUTUZUMAB**

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

14764

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: For combination use with acalabrutinib from treatment cycles 2 to 7 inclusive in first-line therapy

Clinical criteria:

- The condition must be untreated, **AND**
- The treatment must be in combination with PBS-subsidised acalabrutinib (refer to Product Information for timing of obinutuzumab and acalabrutinib doses).

Injection

13793Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	7	..	*4720.79	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ **OBINUTUZUMAB**

Note The latest International Workshop on CLL (iwCLL) provides guidance on various aspects of management of CLL/SLL.

Notably, two of these are:

- (1) when to treat versus when to monitor the patient without therapy - see 'Indications for treatment' section; and
- (2) recognising progressive disease - see 'Definition of response, relapse, and refractory disease' section.

See the following literature reference for details:

Hallek, M et al. iwCLL guidelines for diagnosis, indications for treatment, response assessment, and supportive management of CLL. **Blood** vol. 131, 25 (2018): 2745-2760.

Note Obinutuzumab is not to be used as monotherapy or in combination with anti-cancer drugs other than chlorambucil under this restriction. For use with venetoclax, refer to the separate listing for this purpose.

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

14326

Chronic lymphocytic leukaemia (CLL)

Treatment Phase: Combination use with chlorambucil only

Clinical criteria:

- The condition must be CD20 positive, **AND**
 - The condition must be previously untreated, **AND**
 - The treatment must be in combination with chlorambucil, **AND**
 - The treatment must only be prescribed for a patient with active disease in accordance with the International Workshop on CLL (iwCLL) guidance (latest version) in relation to when to prescribe drug treatment for this condition.
- Treatment must be discontinued in patients who experience disease progression whilst on this treatment.

Injection

10418H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	7	..	*4720.79	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ **RITUXIMAB**

Note Prescribing/pharmacy claiming: prescribe/claim this benefit through the Section 100 Highly Specialised Drugs Program PBS item code(s) when administered for non-oncology indications. Prescribe/claim this benefit through the Efficient Funding of Chemotherapy PBS item code(s) when administered for oncology indications.

Injection

13090Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	800 mg	11	..	*482.67	31.60	Riximyo [SZ] (rituximab 100 mg/10 mL injection, 2 x 10 mL vials) Riximyo [SZ] (rituximab 500 mg/50 mL injection, 50 mL vial) Ruxience [PF] (rituximab 100 mg/10 mL injection, 10 mL vial) Ruxience [PF] (rituximab 500 mg/50 mL injection, 50 mL vial) Truxima [EW] (rituximab 100 mg/10 mL injection, 2 x 10 mL vials) Truxima [EW] (rituximab 500 mg/50 mL injection, 50 mL vial)

CD22 (Clusters of Differentiation 22) inhibitors
■ INOTUZUMAB OZOGAMICIN

Caution Careful monitoring of patients is required due to risk of developing hepatotoxicity, including life-threatening hepatic veno-occlusive disease, and the increased risk of post-haematopoietic stem cell transplant non-relapse mortality observed in patients treated with inotuzumab.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note A complete remission is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a full recovery of peripheral blood counts with platelets of greater than 100,000 per microliter, and absolute neutrophil count (ANC) of greater than 1,000 per microliter.

Note A complete remission with partial haematological recovery is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a partial recovery of peripheral blood counts with platelets of greater than 50,000 per microliter, and absolute neutrophil count (ANC) of greater than 500 per microliter.

Note Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time when an assessment is required must cease PBS-subsidised therapy with this agent.

Authority required

Acute Lymphoblastic leukaemia

Treatment Phase: Consolidation treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised induction treatment with this drug for this condition, **AND**
 - Patient must have achieved a complete remission; OR
 - Patient must have achieved a complete remission with partial haematological recovery, **AND**
 - The treatment must not be more than 5 treatment cycles under this restriction in a lifetime, **AND**
 - Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug.
- This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.
- The treatment must not exceed 0.5mg per m² for all doses within a treatment cycle
- Treatment with this drug for this condition must not exceed 6 treatment cycles in a lifetime.

Injection

11668D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	2820 mcg	4	..	*40010.22	31.60	Besponsa [PF] (inotuzumab ozogamicin 1 mg injection, 1 vial)

■ INOTUZUMAB OZOGAMICIN

Caution Careful monitoring of patients is required due to risk of developing hepatotoxicity, including life-threatening hepatic veno-occlusive disease, and the increased risk of post-haematopoietic stem cell transplant non-relapse mortality observed in patients treated with inotuzumab.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patients are eligible to receive a loading dose for the first dose of a treatment cycle while receiving induction treatment. Two prescriptions are required, the first prescription for the loading dose at a dose no higher than 0.8mg per m², and the second prescription for two doses at a dose no higher than 0.5mg per m². Both prescriptions must be submitted with the initial application.

Note Once a patient achieves complete remission or complete remission with partial haematological recovery, a new prescription must be written under the consolidation treatment phase.

Note A complete remission is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a full recovery of peripheral blood counts with platelets of greater than 100,000 per microliter, and absolute neutrophil count (ANC) of greater than 1,000 per microliter.

Note A complete remission with partial haematological recovery is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a partial recovery of peripheral blood counts with platelets of greater than 50,000 per microliter, and absolute neutrophil count (ANC) of greater than 500 per microliter.

Note Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time when an assessment is required must cease PBS-subsidised therapy with this agent.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acute lymphoblastic leukaemia
Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be relapsed or refractory B-precursor cell ALL, with an Eastern Cooperative Oncology Group (ECOG) performance status of 2 or less, **AND**
 - Patient must have received intensive combination chemotherapy for initial treatment of ALL or for subsequent salvage therapy, **AND**
 - Patient must not have received more than 1 line of salvage therapy, **AND**
 - Patient must have previously received a tyrosine kinase inhibitor (TKI) if the condition is Philadelphia chromosome positive, **AND**
 - The condition must be CD22-positive, **AND**
 - The condition must have more than 5% blasts in bone marrow, **AND**
 - The treatment must not be more than 3 treatment cycles under this restriction in a lifetime.
- This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

The authority application must be made in writing and must include:

- (1) two completed authority prescription forms;
- (2) a completed Acute Lymphoblastic Leukaemia PBS Authority Application - Supporting Information Form; and
- (3) evidence that the condition is CD22-positive; and
- (4) date of most recent chemotherapy, and if this was the initial chemotherapy regimen or salvage therapy, including what line of salvage; and
- (5) a copy of the most recent bone marrow biopsy report of no more than one month old at the time of application.

The treatment must not exceed 0.8mg per m² for the first dose of a treatment cycle (Day 1), and 0.5mg per m² for subsequent doses (Days 8 and 15) within a treatment cycle.

Treatment with this drug for this condition must not exceed 6 treatment cycles in a lifetime.

Injection

11673J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3384 mcg	2	..	*53302.99	31.60	Besponsa [PF] (inotuzumab ozogamicin 1 mg injection, 1 vial)

CD38 (Clusters of Differentiation 38) inhibitors

▪ **DARATUMUMAB**

Note This drug is not PBS-subsidised for use in patients with multiple myeloma who have received two or more prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent, or, who are refractory to both a PI and an immunomodulatory agent, as monotherapy.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment of second-line drug therapy for weeks 10 to 24 (administered every 3 weeks)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - The treatment must be in combination with bortezomib and dexamethasone, **AND**
 - Patient must not have developed disease progression while receiving treatment with this drug for this condition.
- Progressive disease is defined as at least 1 of the following:
- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
 - (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
 - (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
 - (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
 - (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
 - (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
 - (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

1225K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1920 mg	4	..	*11979.26	31.60	Darzalex [JC] (daratumumab 100 mg/5 mL injection, 5 mL vial) Darzalex [JC] (daratumumab 400 mg/20 mL injection, 20 mL vial)

▪ DARATUMUMAB

Note This drug is not PBS-subsidised for use in patients with multiple myeloma who have received two or more prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent, or, who are refractory to both a PI and an immunomodulatory agent, as monotherapy.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment of second-line drug therapy from week 25 until disease progression (administered every 4 weeks)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

12226L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1920 mg	5	..	*11979.26	31.60	Darzalex [JC] (daratumumab 100 mg/5 mL injection, 5 mL vial)
						Darzalex [JC] (daratumumab 400 mg/20 mL injection, 20 mL vial)

▪ DARATUMUMAB

Note This drug is not PBS-subsidised for use in patients with multiple myeloma who have received two or more prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent, or, who are refractory to both a PI and an immunomodulatory agent, as monotherapy.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Initial treatment as second-line drug therapy for weeks 1 to 9 (administered once weekly)

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with bortezomib and dexamethasone, **AND**
- Patient must have progressive disease after only one prior therapy (i.e. use must be as second-line drug therapy; use as third-line drug therapy or beyond is not PBS-subsidised).

Treatment criteria:

- Patient must be undergoing treatment with this drug in one of the following situations: (i) for the first time, irrespective of whether the diagnosis has been reclassified (i.e. the diagnosis has changed between multiple myeloma/amyloidosis), (ii) changing the drug's form (intravenous/subcutaneous) within the first 9 weeks of treatment for the same PBS indication.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or

(g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein. Details of: the histological diagnosis of multiple myeloma; prior treatments including name(s) of drug(s) and date of most recent treatment cycle; the basis of the diagnosis of progressive disease or failure to respond; and which disease activity parameters will be used to assess response, must be documented in the patient's medical records.

Confirmation of eligibility for treatment with current diagnostic reports of at least one of the following must be documented in the patient's medical records:

- (a) the level of serum monoclonal protein; or
- (b) Bence-Jones proteinuria - the results of 24-hour urinary light chain M protein excretion; or
- (c) the serum level of free kappa and lambda light chains; or
- (d) bone marrow aspirate or trephine; or
- (e) if present, the size and location of lytic bone lesions (not including compression fractures); or
- (f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
- (g) if present, the level of hypercalcaemia, corrected for albumin concentration.

As these parameters must be used to determine response, results for either (a) or (b) or (c) should be documented for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) must be documented in the patient's medical records. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (current serum M protein less than 10 g per L) must be documented in the patient's medical records.

A line of therapy is defined as 1 or more cycles of a planned treatment program. This may consist of 1 or more planned cycles of single-agent therapy or combination therapy, as well as a sequence of treatments administered in a planned manner.

A new line of therapy starts when a planned course of therapy is modified to include other treatment agents (alone or in combination) as a result of disease progression, relapse, or toxicity, with the exception to this being the need to attain a sufficient response for stem cell transplantation to proceed. A new line of therapy also starts when a planned period of observation off therapy is interrupted by a need for additional treatment for the disease.

Injection

	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
12230Q	1920 mg	8	..	*11979.26	31.60	Darzalex [JC] (daratumumab 100 mg/5 mL injection, 5 mL vial) Darzalex [JC] (daratumumab 400 mg/20 mL injection, 20 mL vial)

HER2 (Human Epidermal Growth Factor Receptor 2) inhibitors

▪ **PERTUZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) either in the primary tumour or a metastatic lesion, confirmed through a pathology report from an Approved Pathology Authority, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must not have received prior anti-HER2 therapy for this condition, **AND**
- Patient must not have received prior chemotherapy for this condition, **AND**
- The treatment must be in combination with trastuzumab and a taxane, **AND**
- The treatment must not be in combination with nab-paclitaxel, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Details (date, unique identifying number/code, or provider number) of the pathology report from an Approved Pathology Authority confirming evidence of HER2 gene amplification in the primary tumour or a metastatic lesion by in situ hybridisation (ISH) must be provided at the time of application.

The pathology report must be documented in the patient's medical records.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to seeking the initial authority approval.

Injection

	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
10334X	840 mg	*6051.13	31.60	Perjeta [RO] (pertuzumab 420 mg/14 mL injection, 14 mL vial)

■ PERTUZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note The criterion that limits breaks in treatment with pertuzumab under this restriction has been temporarily modified due to the current risk of COVID-19. This allows an extended break in therapy with PBS-subsidised pertuzumab in patients who are at risk of COVID-19.

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug, **AND**
- The treatment must be in combination with trastuzumab, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

The treatment must not exceed a lifetime total of one course. However, treatment breaks are permitted. A patient who has a treatment break in PBS-subsidised treatment with this drug for reasons other than disease progression is eligible to continue to receive PBS-subsidised treatment with this drug.

Where a patient has had a treatment break the length of the break is measured from the date the most recent treatment was stopped to the date of the application for further treatment.

Injection

10308M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	420 mg	3	..	*3091.52	31.60	Perjeta [RO] (pertuzumab 420 mg/14 mL injection, 14 mL vial)

■ TRASTUZUMAB

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

10296

Early HER2 positive breast cancer

Treatment Phase: Initial treatment (weekly regimen)

Clinical criteria:

- Patient must have undergone surgery (adjuvant) or be preparing for surgery (neoadjuvant), **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
- Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.

HER2 positivity must be demonstrated by in situ hybridisation (ISH).

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

Injection

7264H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	500 mg	*486.43	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

■ TRASTUZUMAB

Note Increased maximum quantity will be authorised where a patient requires a new loading dose due to a break in therapy of more than 1 week but less than 6 weeks from the last dose or a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

10213

Early HER2 positive breast cancer

Treatment Phase: Continuing treatment (weekly regimen)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
- Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.

Injection

7265J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	250 mg	9	..	*346.04	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB**

Note Increased maximum quantity will be authorised where a patient requires a new loading dose due to a break in therapy of more than 1 week but less than 6 weeks from the last dose or a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

10294

Early HER2 positive breast cancer

Treatment Phase: Continuing treatment (3 weekly regimen)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
- Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.

Injection

7267L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	750 mg	3	..	*729.71	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum quantity will be authorised where a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required (STREAMLINED)

9349

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Where a patient has a break in trastuzumab therapy of more than 1 week from when the last dose was due, a new loading dose may be required.

Injection

10383L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	750 mg	3	..	*729.71	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial)

Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial)
 Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial)
 Trazimera [PF] (trastuzumab 150 mg injection, 1 vial)
 Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

■ TRASTUZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum quantity will be authorised where a patient's weight is greater than 125 kg.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required (STREAMLINED)

9353

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) either in the primary tumour or a metastatic lesion, **AND**
- The treatment must not be in combination with nab-paclitaxel, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

Injection

10402L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	*870.10	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

■ TRASTUZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum quantity will be authorised where a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9573

Metastatic (Stage IV) HER2 positive adenocarcinoma of the stomach or gastro-oesophageal junction

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) positivity as demonstrated by immunohistochemistry 2+ or more in tumour material, **AND**
- Patient must have evidence of HER2 gene amplification as demonstrated by in situ hybridisation results based on more than 6 copies of HER2 in the same tumour tissue sample, **AND**
- Patient must have evidence of HER2 gene amplification as demonstrated by in situ hybridisation results based on the ratio of HER2 to chromosome 17 being more than 2 in the same tumour tissue sample, **AND**
- Patient must commence treatment in combination with platinum based chemotherapy and capecitabine; **OR**
- Patient must commence treatment in combination with platinum based chemotherapy and 5 fluorouracil, **AND**
- Patient must not have previously received this drug for this condition, **AND**
- Patient must not have received prior chemotherapy for this condition, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

Injection

10589H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	*870.10	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial)

Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum quantity will be authorised where a patient requires a new loading dose due to a break in therapy of more than 1 week but less than 6 weeks from the last dose or a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9571

Metastatic (Stage IV) HER2 positive adenocarcinoma of the stomach or gastro-oesophageal junction

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Injection

10597R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	750 mg	3	..	*729.71	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (&) inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

10293

Early HER2 positive breast cancer

Treatment Phase: Initial treatment (3 weekly regimen)

Clinical criteria:

- Patient must have undergone surgery (adjuvant) or be preparing for surgery (neoadjuvant), **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
- Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.

HER2 positivity must be demonstrated by in situ hybridisation (ISH).

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

Injection

7266K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	*870.10	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (&) inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB DERUXTECAN**

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Clinical criteria:

- Patient must have evidence of human epidermal growth factor (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) in either the primary tumour/a metastatic lesion - establish this finding once only with the first PBS prescription, **AND**
- The condition must have progressed following treatment with at least one prior HER2 directed regimen for metastatic breast cancer; OR
- The condition must have, at the time of treatment initiation with this drug, progressed during/within 6 months following adjuvant treatment with a HER2 directed therapy, **AND**
- Patient must have, at the time of initiating treatment with this drug, a WHO performance status no higher than 1, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this PBS indication, **AND**
- The treatment must not be prescribed where any of the following is present: (i) left ventricular ejection fraction of less than 50%, (ii) symptomatic heart failure; confirm cardiac function testing for the first PBS prescription only.

Treatment criteria:

- Patient must be undergoing initial treatment with this drug - the following are true: (i) this is the first prescription for this drug, (ii) this prescription seeks no more than 3 repeat prescriptions; OR
- Patient must be undergoing continuing treatment with drug - the following are true: (i) there has been an absence of further disease progression whilst on active treatment with this drug, (ii) this prescription does not seek to re-treat after disease progression, (iii) this prescription seeks no more than 8 repeat prescriptions.

Confirm that the following information is documented/retained in the patient's medical records once only with the first PBS prescription:

- 1) Evidence of HER2 gene amplification (evidence obtained in relation to past PBS treatment is acceptable).
- 2) Details of prior HER2 directed drug regimens prescribed for the patient.
- 3) Cardiac function test results (evidence obtained in relation to past PBS treatment is acceptable).

Injection

13713R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	675 mg	8	..	*17827.63	31.60	Enhertu [AP] (trastuzumab deruxtecan 100 mg injection, 1 vial)

▪ TRASTUZUMAB EMTANSINE

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) either in the primary tumour or a metastatic lesion, confirmed through a pathology report from an Approved Pathology Authority, **AND**
- The condition must have progressed following treatment with pertuzumab and trastuzumab in combination; OR
- The condition must have progressed during or within 6 months of completing adjuvant therapy with trastuzumab, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

The following information must be provided by the prescriber at the time of application:

(a) details (date, unique identifying number/code or provider number) of the pathology report from an Approved Pathology Authority confirming evidence of HER2 gene amplification in the primary tumour or a metastatic lesion by in situ hybridisation (ISH).

(b) dates of treatment with trastuzumab and pertuzumab;

(c) date of demonstration of progression following treatment with trastuzumab and pertuzumab; or

(d) date of demonstration of progression and date of completion of adjuvant trastuzumab treatment.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application.

All reports must be documented in the patient's medical records.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to seeking the initial authority approval.

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for metastatic (Stage IV) HER2 positive breast cancer, **AND**
- Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug, **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

The treatment must not exceed a lifetime total of one continuous course for this PBS indication.

Injection

10281D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	450 mg	8	..	*7412.41	31.60	Kadcyla [RO] (trastuzumab emtansine 100 mg injection, 1 vial) Kadcyla [RO] (trastuzumab emtansine 160 mg injection, 1 vial)

▪ **TRASTUZUMAB EMTANSINE**

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Authority required

Early HER2 positive breast cancer

Treatment Phase: Initial adjuvant treatment

Clinical criteria:

- The treatment must be prescribed within 12 weeks after surgery, **AND**
- Patient must have, prior to commencing treatment with this drug, evidence of residual invasive cancer in the breast and/or axillary lymph nodes following completion of surgery, as demonstrated by a pathology report, **AND**
- Patient must have completed systemic neoadjuvant therapy that included trastuzumab and taxane-based chemotherapy prior to surgery, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- The treatment must not extend beyond 42 weeks (14 cycles) duration under the initial and the continuing treatment restrictions combined.

Authority applications for initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail and must include:

(a) details (date, unique identifying number/code or provider number) of the pathology report from an Approved Pathology Authority demonstrating evidence of residual invasive carcinoma in the breast and/or axillary lymph nodes following completion of surgery.

The pathology report must be documented in the patient's medical records.

If the application is submitted through HPOS form upload or mail, it must include:

- (i) A completed authority prescription form; and
- (ii) A completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Early HER2 positive breast cancer

Treatment Phase: Continuing adjuvant treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- The treatment must not extend beyond 42 weeks (14 cycles) duration under the initial and the continuing treatment restrictions combined.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Injection

11956G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	450 mg	6	..	*7412.41	31.60	Kadcyla [RO] (trastuzumab emtansine 100 mg injection, 1 vial) Kadcyla [RO] (trastuzumab emtansine 160 mg injection, 1 vial)

EGFR (Epidermal Growth Factor Receptor) inhibitors**■ CETUXIMAB****Note** Special Pricing Arrangements apply.**Authority required (STREAMLINED)****12470**

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with PBS-subsidised encorafenib for this condition.

Injection

12817N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	550 mg	11	..	*1765.62	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

■ CETUXIMAB**Note** Special Pricing Arrangements apply.**Authority required (STREAMLINED)****12483**

Metastatic colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with PBS-subsidised encorafenib for this condition.

Injection

12821T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	880 mg	*2582.49	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

■ CETUXIMAB**Note** A maximum lifetime supply for this indication is limited to a maximum of 8 treatments per site and to 10 treatments per site for patients in whom radiotherapy is interrupted.**Authority required (STREAMLINED)****4788**

Stage III, IVa or IVb squamous cell cancer of the larynx, oropharynx or hypopharynx

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with radiotherapy, **AND**
- Patient must be unable to tolerate cisplatin; OR
- Patient must have a contraindication to cisplatin according to the TGA-approved Product Information.

Injection

7240C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	550 mg	5	..	*1765.62	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

■ CETUXIMAB**Note** No increase in the maximum number of repeats may be authorised.**Authority required (STREAMLINED)****4794**

Stage III, IVa or IVb squamous cell cancer of the larynx, oropharynx or hypopharynx

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be for the week prior to radiotherapy, **AND**
- Patient must have a contraindication to cisplatin according to the TGA-approved Product Information.

Authority required (STREAMLINED)**4785**

Stage III, IVa or IVb squamous cell cancer of the larynx, oropharynx or hypopharynx

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with radiotherapy, **AND**
- Patient must be unable to tolerate cisplatin.

Injection

7223E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	880 mg	*2582.49	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

▪ **CETUXIMAB**

Note Special Pricing Arrangements apply.

Note This drug is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Authority required (STREAMLINED)

12045

Metastatic colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have RAS wild-type metastatic colorectal cancer, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The condition must have failed to respond to first-line chemotherapy; OR
- The condition must have progressed following first-line treatment with pembrolizumab for dMMR mCRC, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on panitumumab are not eligible to receive PBS-subsidised cetuximab.

Patients who have developed intolerance to panitumumab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised cetuximab.

Authority required (STREAMLINED)

4908

Metastatic colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have RAS wild-type metastatic colorectal cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must be previously untreated, **AND**
- The treatment must be in combination with first-line chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Injection

7242E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	880 mg	*2582.49	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

▪ **CETUXIMAB**

Note Special Pricing Arrangements apply.

Note This drug is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Note This drug is not PBS-subsidised when chemotherapy partners are switched whilst maintaining an anti-EGFR antibody backbone in the face of progressive disease.

Note The treatment must not exceed a single course of therapy with this drug for metastatic colorectal cancer in a patient's lifetime.

Authority required (STREAMLINED)

4912

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for first-line treatment of RAS wild-type metastatic colorectal cancer, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must be in combination with first-line chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Injection

10265G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	550 mg	18	..	*1765.62	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial)

Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

■ CETUXIMAB

Note Special Pricing Arrangements apply.

Note This drug is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Note This drug is not PBS-subsidised when chemotherapy partners are switched whilst maintaining an anti-EGFR antibody backbone in the face of progressive disease.

Note The treatment must not exceed a single course of therapy with this drug for metastatic colorectal cancer in a patient's lifetime.

Authority required (STREAMLINED)

12016

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for treatment of RAS wild-type metastatic colorectal cancer after failure of first-line chemotherapy; OR
- Patient must have received an initial authority prescription for this drug for treatment of RAS wild-type metastatic colorectal cancer after failure of treatment with first-line pembrolizumab for dMMR mCRC, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on panitumumab are not eligible to receive PBS-subsidised cetuximab.

Patients who have developed intolerance to panitumumab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised cetuximab.

Injection

7273T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	550 mg	11	..	*1765.62	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

■ PANITUMUMAB

Note This drug is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Authority required (STREAMLINED)

12066

Metastatic colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have RAS wild-type metastatic colorectal cancer, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The condition must have failed to respond to first-line chemotherapy; OR
- The condition must have progressed following first-line treatment with pembrolizumab for dMMR mCRC, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on cetuximab are not eligible to receive PBS-subsidised panitumumab.

Patients who have developed intolerance to cetuximab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised panitumumab.

Authority required (STREAMLINED)

12035

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for treatment of RAS wild-type metastatic colorectal cancer after failure of first-line chemotherapy; OR
- Patient must have received an initial authority prescription for this drug for treatment of RAS wild-type metastatic colorectal cancer after failure of treatment with first-line pembrolizumab for dMMR mCRC, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on cetuximab are not eligible to receive PBS-subsidised panitumumab.

Patients who have developed intolerance to cetuximab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised panitumumab.

Note This drug is not PBS-subsidised when chemotherapy partners are switched whilst maintaining an anti-EGFR antibody backbone in the face of progressive disease.

Note The treatment must not exceed a single course of therapy with this drug for metastatic colorectal cancer in a patient's lifetime.

Injection

10069Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	720 mg	5	..	*3785.69	31.60	Vectibix [AN] (panitumumab 100 mg/5 mL injection, 5 mL vial) Vectibix [AN] (panitumumab 400 mg/20 mL injection, 20 mL vial)

▪ **PANITUMUMAB**

Note Special Pricing Arrangements apply.

Note Panitumumab is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Authority required (STREAMLINED)

5526

Metastatic colorectal cancer

Treatment Phase: Initial Treatment

Clinical criteria:

- Patient must have RAS wild-type metastatic colorectal cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must be previously untreated, **AND**
- The treatment must be in combination with first-line chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on cetuximab are not eligible to receive PBS-subsidised panitumumab.

Patients who have developed intolerance to cetuximab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised panitumumab.

Authority required (STREAMLINED)

5452

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for panitumumab for first-line treatment of RAS wild-type metastatic colorectal cancer, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must be in combination with first-line chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on cetuximab are not eligible to receive PBS-subsidised panitumumab.

Patients who have developed intolerance to cetuximab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised panitumumab.

Note This drug is not PBS-subsidised when chemotherapy partners are switched whilst maintaining an anti-EGFR antibody backbone in the face of progressive disease.

Note The treatment must not exceed a single course of therapy with this drug for metastatic colorectal cancer in a patient's lifetime.

Injection

10508C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	720 mg	9	..	*3785.69	31.60	Vectibix [AN] (panitumumab 100 mg/5 mL injection, 5 mL vial) Vectibix [AN] (panitumumab 400 mg/20 mL injection, 20 mL vial)

PD-1/PDL-1 (Programmed cell death protein 1/death ligand 1) inhibitors

▪ **ATEZOLIZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10297

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- Patient must have stable or responding disease.

Injection

11297N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	7	..	*6973.74	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

■ ATEZOLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10216**

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing first-line treatment of metastatic disease - 3 weekly treatment regimen

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab until disease progression, unless not tolerated.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug in this line of treatment, **AND**
- Patient must have stable or responding disease.

Injection

11801D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	7	..	*6973.74	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

■ ATEZOLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10215**

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Continuing treatment - 4 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have stable or responding disease.

Injection

11957H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	5	..	*9710.45	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

■ ATEZOLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10257**

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing first-line treatment of metastatic disease, as monotherapy, where concomitant bevacizumab has ceased due to intolerance - 4 weekly treatment regimen

Clinical criteria:

- Patient must have experienced intolerance to combination treatment with bevacizumab, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug in this line of treatment, **AND**
- Patient must have stable or responding disease, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

12098R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	5	..	*9710.45	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

■ ATEZOLIZUMAB

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**13443**

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The condition must have progressed on or after prior platinum based chemotherapy; OR
- The condition must have progressed after treatment with tepotinib.

Injection

11309F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	5	..	*6973.74	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ **ATEZOLIZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10206

Extensive-stage small cell lung cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be previously untreated, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be in combination with etoposide and a platinum-based antineoplastic drug.

Injection

11927R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	3	..	*6973.74	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ **ATEZOLIZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10521

Extensive-stage small cell lung cancer

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Injection

11928T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	4	..	*6973.74	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ **ATEZOLIZUMAB**

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13446

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Initial treatment - 4 weekly treatment regimen

Clinical criteria:

- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The condition must have progressed on or after prior platinum based chemotherapy; OR
- The condition must have progressed after treatment with tepotinib.

Injection

11940K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	3	..	*9710.45	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

ATEZOLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10509**

Extensive-stage small cell lung cancer

Treatment Phase: Continuing treatment - 4 weekly treatment regimen

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Injection

12076N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	3	..	*9710.45	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

ATEZOLIZUMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10917**

Advanced (unresectable) Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Continuing treatment of hepatocellular carcinoma - 3 weekly treatment regimen

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab until disease progression, unless not tolerated.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - Patient must not have developed disease progression while being treated with this drug for this condition.
- PBS supply of this drug must be through only one of the two continuing treatment regimens at any given time

Injection

12155R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	8	..	*6973.74	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

ATEZOLIZUMAB

Note No increase in the maximum amount or number of units may be authorised.

Note Increased repeats of up to 11 may be requested for doses of 840 mg administered every 2 weeks

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10972**

Advanced (unresectable) Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Continuing treatment where bevacizumab is discontinued - 4 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - Patient must not have developed disease progression while being treated with this drug for this condition.
- PBS supply of this drug must be through only one of the two continuing treatment regimens at any given time

Injection

12159Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	5	..	*9710.45	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

ATEZOLIZUMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**13451**

Resected early stage (Stage II to IIIA) non-small cell lung cancer (NSCLC)

Treatment Phase: 1,680 mg administered once every 4 weeks, or 840 mg every 2 weeks

Population criteria:

- Patient must be both: (i) initiating treatment, (ii) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy; OR
- Patient must be continuing existing PBS-subsidised treatment with this drug; OR
- Patient must be both: (i) transitioning from existing non-PBS to PBS subsidised supply of this drug, (ii) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy at the time this drug was initiated.

Clinical criteria:

- Patient must have/have had a WHO performance status score of no greater than 1 at treatment initiation with this drug. **AND**
- The treatment must be for the purpose of adjuvant therapy following all of: (i) surgical resection, (ii) platinum-based chemotherapy, **AND**
- The condition must have/have had, at treatment commencement, an absence of each of the following gene abnormalities confirmed via tumour material sampling: (i) an activating epidermal growth factor receptor (EGFR) gene mutation, (ii) an anaplastic lymphoma kinase (ALK) gene rearrangement, **AND**
- The condition must have/have had, at treatment commencement, confirmation of programmed cell death ligand 1 (PD-L1) expression on at least 50% of tumour cells, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment that does not occur beyond the following, whichever comes first: (i) the first instance of disease progression/recurrence, (ii) 12 months in total for this condition from the first administered dose; mark any remaining repeat prescriptions with the words 'cancelled' where (i)/(ii) has occurred.

Injection

13170E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	5	..	*9710.45	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

▪ **ATEZOLIZUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13442

Resected early stage (Stage II to IIIA) non-small cell lung cancer (NSCLC)

Treatment Phase: 1,200 mg administered once every 3 weeks

Population criteria:

- Patient must be both: (i) initiating treatment, (ii) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy; OR
- Patient must be continuing existing PBS-subsidised treatment with this drug; OR
- Patient must be both: (i) transitioning from existing non-PBS to PBS subsidised supply of this drug, (ii) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy at the time this drug was initiated.

Clinical criteria:

- Patient must have/have had a WHO performance status score of no greater than 1 at treatment initiation with this drug. **AND**
- The treatment must be for the purpose of adjuvant therapy following all of: (i) surgical resection, (ii) platinum-based chemotherapy, **AND**
- The condition must have/have had, at treatment commencement, an absence of each of the following gene abnormalities confirmed via tumour material sampling: (i) an activating epidermal growth factor receptor (EGFR) gene mutation, (ii) an anaplastic lymphoma kinase (ALK) gene rearrangement, **AND**
- The condition must have/have had, at treatment commencement, confirmation of programmed cell death ligand 1 (PD-L1) expression on at least 50% of tumour cells, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment that does not occur beyond the following, whichever comes first: (i) the first instance of disease progression/recurrence, (ii) 12 months in total for this condition from the first administered dose; mark any remaining repeat prescriptions with the words 'cancelled' where (i)/(ii) has occurred.

Injection

13172G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	7	..	*6973.74	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ **ATEZOLIZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13448

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment 1

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab and platinum-doublet chemotherapy.

Clinical criteria:

- The condition must be non-squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene mutation or an anaplastic lymphoma kinase (ALK) gene rearrangement in tumour material.

Authority required (STREAMLINED)**10125**

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment 2

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab and platinum-doublet chemotherapy.

Clinical criteria:

- The condition must be non-squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must have evidence of an activating epidermal growth factor receptor (EGFR) gene mutation or of an anaplastic lymphoma kinase (ALK) gene rearrangement in tumour material, **AND**
- Patient must have progressive disease following treatment with an epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI) OR an anaplastic lymphoma kinase (ALK) tyrosine kinase inhibitor (TKI), **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer.

Injection

11792P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	5	..	*6973.74	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

■ ATEZOLIZUMAB

Caution The safety of atezolizumab in combination with bevacizumab has not been established in patients who have incompletely treated varices, variceal bleeding within the previous 6 months or who are at high risk of bleeding. Patients should be assessed for risk of variceal bleeding prior to treatment with this combination.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10939**

Advanced (unresectable) Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Initial treatment

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab and atezolizumab until disease progression, unless not tolerated.

Clinical criteria:

- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must not be suitable for transarterial chemoembolisation, **AND**
- Patient must have Child Pugh class A, **AND**
- The condition must be untreated with systemic therapy; OR
- Patient must have developed intolerance to a vascular endothelial growth factor (VEGF) tyrosine kinase inhibitor (TKI) of a severity necessitating permanent treatment withdrawal.

Injection

12167J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	3	..	*6973.74	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

■ AVELUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

8947

Stage IV (metastatic) Merkel Cell Carcinoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed a total of 9 doses at a maximum dose of 10 mg per kg every 2 weeks under this restriction.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

11679Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	8	..	*8390.07	31.60	Bavencio [SG] (avelumab 200 mg/10 mL injection, 10 mL vial)

▪ **AVELUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10023

Stage IV (metastatic) Merkel Cell Carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 10 mg per kg every 2 weeks under this restriction.

Injection

11685B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	11	..	*8390.07	31.60	Bavencio [SG] (avelumab 200 mg/10 mL injection, 10 mL vial)

▪ **AVELUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13290

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Treatment Phase: Maintenance therapy - Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

13132E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	800 mg	11	..	*5637.35	31.60	Bavencio [SG] (avelumab 200 mg/10 mL injection, 10 mL vial)

▪ **AVELUMAB**

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13313

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Treatment Phase: Maintenance therapy - Initial treatment

Clinical criteria:

- Patient must have received first-line platinum-based chemotherapy, **AND**
- Patient must not have progressive disease following first-line platinum-based chemotherapy, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

13123Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	800 mg	7	..	*5637.35	31.60	Bavencio [SG] (avelumab 200 mg/10 mL injection, 10 mL vial)

■ CEMIPILIMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**15094**

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must not exceed a total of 7 doses under this restriction.

Injection

13160P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	350 mg	6	..	*7523.97	31.60	Libtayo [WM] (cemiplimab 350 mg/7 mL injection, 7 mL vial)

■ CEMIPILIMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**15063**

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a total of 35 cycles or up to 24 months of treatment under both initial and continuing treatment restrictions, whichever comes first.

Injection

13162R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	350 mg	6	..	*7523.97	31.60	Libtayo [WM] (cemiplimab 350 mg/7 mL injection, 7 mL vial)

■ CEMIPILIMAB

Caution In the first few months after starting immunotherapy, a transient tumour flare may occur that may be mistaken as disease progression despite an overall positive response to treatment.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Metastatic or locally advanced cutaneous squamous cell carcinoma (CSCC)

Treatment Phase: Initial treatment covering the first 3 treatment cycles

Clinical criteria:

- The condition must be unsuitable for each of: (i) curative surgical resection, (ii) curative radiotherapy, **AND**
- Patient must have had a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

13135H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	350 mg	2	..	*7523.97	31.60	Libtayo [WM] (cemiplimab 350 mg/7 mL injection, 7 mL vial)

▪ **CEMIPLIMAB**

Caution In the first few months after starting immunotherapy, a transient tumour flare may occur that may be mistaken as disease progression despite an overall positive response to treatment.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Metastatic or locally advanced cutaneous squamous cell carcinoma (CSCC)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Patient must not be undergoing treatment with this drug as a PBS benefit where the treatment duration extends beyond the following, whichever comes first: (i) disease progression despite treatment with this drug, (ii) 24 months from treatment initiation; annotate any remaining repeat prescriptions with the word 'cancelled' where this occurs.

Injection

13153G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	350 mg	7	..	*7523.97	31.60	Libtayo [WM] (cemiplimab 350 mg/7 mL injection, 7 mL vial)

▪ **DOSTARLIMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

15163

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Initial treatment covering the first 6 treatment cycles

Clinical criteria:

- Patient must have deficient mismatch repair (dMMR) endometrial cancer, as determined by immunohistochemistry test, **AND**
- The condition must be unsuitable for at least one of the following: (i) curative surgical resection, (ii) curative radiotherapy, **AND**
- The treatment must be initiated in combination with platinum-containing chemotherapy, **AND**
- The condition must be, at treatment initiation with this drug, either: (i) untreated with systemic therapy, (ii) treated with neoadjuvant/adjuvant systemic therapy, but the cancer has recurred or progressed after more than 6 months from the last dose of systemic therapy, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation.

Injection

14133W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	500 mg	5	..	*7886.48	31.60	Jemperli [GK] (dostarlimab 500 mg/10 mL injection, 10 mL vial)

▪ **DOSTARLIMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

15205

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 36 cumulative months from the first administered dose, once in a lifetime.

Authority required (STREAMLINED)

15196

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Transitioning from non-PBS to PBS-subsidised treatment - Grandfather treatment

Clinical criteria:

- Patient must have deficient mismatch repair (dMMR) endometrial cancer, as determined by immunohistochemistry test, **AND**
- Patient must have received non-PBS-subsidised treatment with this drug for this condition prior to 1 May 2024, **AND**
- The condition must be, prior to initiation of non-PBS-subsidised treatment with this drug, unsuitable for at least one of the following: (i) curative surgical resection, (ii) curative radiotherapy, **AND**
- The condition must be, prior to initiation of non-PBS-subsidised treatment with this drug, either: (i) untreated with systemic therapy, (ii) treated with neoadjuvant/adjuvant systemic therapy, but the cancer has recurred or progressed after more than 6 months from the last dose of systemic therapy, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation, **AND**
- The treatment must be, at initiation of non-PBS-subsidised treatment with this drug, used in combination with platinum-containing chemotherapy, **AND**
- Patient must not have developed disease progression while receiving non-PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 36 cumulative months from the first administered dose, once in a lifetime.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Injection

14109N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	3	..	*15641.05	31.60	Jemperli [GK] (dostarlimab 500 mg/10 mL injection, 10 mL vial)

■ DURVALUMAB

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10126**

Unresectable Stage III non-small cell lung cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have received platinum based chemoradiation therapy, **AND**
- The condition must not have progressed following platinum based chemoradiation therapy, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

Authority required (STREAMLINED)**12271**

Unresectable Stage III non-small cell lung cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The treatment must not exceed 12 months in total for this condition under the initial and continuing restriction combined, **AND**
- The treatment must be once in a lifetime with this drug for this condition.

Injection

11911X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1500 mg	4	..	*11619.27	31.60	Imfinzi [AP] (durvalumab 120 mg/2.4 mL injection, 2.4 mL vial) Imfinzi [AP] (durvalumab 500 mg/10 mL injection, 10 mL vial)

■ DURVALUMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**14708**

Locally advanced, metastatic or recurrent biliary tract cancer (intrahepatic cholangiocarcinoma, extrahepatic cholangiocarcinoma, and gallbladder cancer)

Population criteria:

- Patient must have either of the following at treatment initiation: (i) locally advanced biliary tract cancer that is untreated with systemic anti-cancer therapy in the unresectable setting, (ii) metastatic biliary tract cancer that is untreated with systemic anti-cancer therapy in the metastatic setting.

Clinical criteria:

- Patient must have/have had a WHO performance status score of no greater than 1 at treatment initiation with this drug. **AND**
- The treatment must be/have been initiated with both: (i) gemcitabine, (ii) cisplatin (refer to Product Information of gemcitabine and cisplatin for dosing information), **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Injection

13745K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1500 mg	5	..	*11619.27	31.60	Imfinzi [AP] (durvalumab 120 mg/2.4 mL injection, 2.4 mL vial)
						Imfinzi [AP] (durvalumab 500 mg/10 mL injection, 10 mL vial)

▪ **DURVALUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10509

Extensive-stage small cell lung cancer

Treatment Phase: Continuing treatment - 4 weekly treatment regimen

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Injection

13766M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1500 mg	5	..	*11619.27	31.60	Imfinzi [AP] (durvalumab 120 mg/2.4 mL injection, 2.4 mL vial)
						Imfinzi [AP] (durvalumab 500 mg/10 mL injection, 10 mL vial)

▪ **DURVALUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10206

Extensive-stage small cell lung cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be previously untreated, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be in combination with etoposide and a platinum-based antineoplastic drug.

Injection

13779F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1500 mg	3	..	*11619.27	31.60	Imfinzi [AP] (durvalumab 120 mg/2.4 mL injection, 2.4 mL vial)
						Imfinzi [AP] (durvalumab 500 mg/10 mL injection, 10 mL vial)

▪ **NIVOLUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

11477

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Continuing treatment as second-line drug therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- Patient must have stable or responding disease. Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11152Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**9299**

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11157F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**9252**

Recurrent or metastatic squamous cell carcinoma of the oral cavity, pharynx or larynx

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have stable or responding disease, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11425H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**9321**

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Maintenance treatment

Clinical criteria:

- Patient must have previously received of up to maximum 4 doses of PBS-subsidised combined therapy with nivolumab and ipilimumab as induction for this condition, **AND**
- The treatment must be as monotherapy for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

11626X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

9298

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Authority required (STREAMLINED)

13839

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Maintenance treatment

Clinical criteria:

- Patient must have previously received of up to maximum 4 doses of PBS-subsidised combined therapy with nivolumab and ipilimumab as induction for this condition, **AND**
- The treatment must be as monotherapy for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this PBS indication.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

10748Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

14816

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
- Patient must not have experienced disease progression whilst on adjuvant PD-1 inhibitor treatment or disease recurrence within 6 months of completion of adjuvant PD-1 inhibitor treatment if treated for resected Stage IIIB, IIIC, IIID or IV melanoma, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

10775D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	8	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13445

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Initial treatment as second-line drug therapy

Clinical criteria:

- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**

- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The condition must have progressed on or after prior platinum based chemotherapy; OR
- The condition must have progressed after treatment with tepotinib.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11143L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	8	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

9216

Recurrent or metastatic squamous cell carcinoma of the oral cavity, pharynx or larynx

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The condition must have progressed within 6 months of the last dose of prior platinum based chemotherapy, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor for this condition.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated. Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11434T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	8	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14830

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Induction treatment

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
- Patient must have an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1, **AND**
- The condition must not be ocular or uveal melanoma, **AND**
- The treatment must be in combination with PBS-subsidised treatment with ipilimumab as induction for this condition. Induction treatment with nivolumab must not exceed a total of 4 doses at a maximum dose of 1 mg per kg every 3 weeks. Induction treatment with ipilimumab must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks.

Injection

11532Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	3	..	*2532.57	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.
Stable disease (SD) is small changes that do not meet above criteria.

Authority required (STREAMLINED)

9312

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial Treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must have progressive disease according to the Response Evaluation Criteria in Solid Tumours (RECIST) following prior treatment with a tyrosine kinase inhibitor; OR
- Patient must have developed intolerance to a tyrosine kinase inhibitor of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11159H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	8	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13433

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial combination treatment (with ipilimumab) as first-line drug therapy

Clinical criteria:

- The condition must be squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must be in combination with platinum-based chemotherapy for the first two cycles, **AND**
- The treatment must be in combination with ipilimumab.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

11468

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing combination treatment (with ipilimumab) of first-line drug therapy

Clinical criteria:

- The condition must be squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed 24 months in total, measured from the initial dose, or, must not extend beyond disease progression, whichever comes first, **AND**
- The treatment must be in combination with ipilimumab.

Injection

12315E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	360 mg	13	..	*7333.85	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note An increase of number of repeats may be authorised up to 11 if the patient is receiving a weight based dosing of 3mg/kg every 2 weeks.

Authority required (STREAMLINED)

11985

Unresectable malignant mesothelioma

Clinical criteria:

- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be in combination with PBS-subsidised ipilimumab, unless an intolerance to ipilimumab of a severity necessitating permanent treatment withdrawal of ipilimumab, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a maximum total of 24 months in a lifetime for this condition.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

12574T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	360 mg	8	..	*7333.85	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Caution In the first few months after starting immunotherapy, a transient tumour flare may occur that may be mistaken as disease progression despite an overall positive response to treatment.

Note The stated maximum amount in this listing is based on this drug's approved Product Information recommended dosing in specific cancer types - the drug may be prescribed in a quantity up to this amount, but need not be this amount for every cancer type. Refer to this drug's approved Product Information (Dose and Method of Administration or Clinical Trials sections) for the recommended dosing in the specific cancer type.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14676

Advanced or metastatic gastro-oesophageal cancers

Clinical criteria:

- Patient must have/have had, at the time of initiating treatment with this drug, a WHO performance status no higher than 1, **AND**
- Patient must be untreated (up until initiating this drug) with programmed cell death-1/ligand-1 (PD-1/PD-L1) inhibitor therapy for gastro-oesophageal cancer.

Treatment criteria:

- Patient must not be undergoing treatment with this drug as a PBS benefit where the treatment duration extends beyond the following, whichever comes first: (i) disease progression despite treatment with this drug, (ii) 24 months from treatment initiation; annotate any remaining repeat prescriptions with the word 'cancelled' where this occurs.

Population criteria:

- Patient must be in one of the three population subsets described below.

Population 1

Conditions: gastric cancer, gastro-oesophageal junction cancer, oesophageal adenocarcinoma

Concomitant therapies: chemotherapy containing at least a fluoropyrimidine drug plus a platinum drug

Line of treatment: first-line drug treatment

Additional clinical finding: HER2 negative

Population 2

Condition: oesophageal squamous cell carcinoma (can be recurrent)

Concomitant therapies: chemotherapy containing at least a fluoropyrimidine drug plus a platinum drug

Line of treatment: first-line drug treatment

Additional clinical finding: unresectable

Population 3

Condition: oesophageal squamous cell carcinoma (can be recurrent)

Line of treatment: second-line drug treatment after chemotherapy containing at least a fluoropyrimidine drug plus a platinum drug

Additional clinical finding: unresectable

Injection

13117J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	13	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Up to an additional 2 repeat prescriptions (7 in total) may be sought only where dosing is on a 2-weekly schedule in the first 16 weeks of treatment. This listing's stated number of repeat prescriptions is based on 4-weekly dosing.

Note No increase in the maximum amount or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Adjuvant treatment of stage II or III oesophageal cancer or gastro-oesophageal junction cancer

Clinical criteria:

- The condition must have histological evidence confirming a diagnosis of a least one of: (i) adenocarcinoma, (ii) squamous cell cancer; document this evidence in the patient's medical records, **AND**
- The condition must have been treated with neoadjuvant platinum-based chemoradiotherapy, **AND**
- The treatment must be for the purposes of adjuvant use following complete surgical resection that occurred within 16 weeks prior to initiating this drug, **AND**
- The condition must have evidence, through resected specimen, that residual disease meets the Tumour Nodes Metastases (TNM) staging system (as published by the Union for International Cancer Control) of either: (i) at least ypT1, (ii) at least ypN1; document this evidence in the patient's medical records, **AND**
- Patient must have/have had, at the time of initiating treatment with this drug, a WHO performance status no higher than 1, **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment with a dosing regimen as set out in the drug's approved Australian Product Information, **AND**
- Patient must not be undergoing PBS-subsidised treatment with this drug where this prescription extends treatment beyond whichever comes first: (i) 12 months from treatment initiation, irrespective of whether initial treatment was PBS-subsidised/non-PBS-subsidised, (ii) disease recurrence despite treatment with this drug; annotate any remaining repeat prescriptions with the word 'cancelled' where this occurs.

Injection

13246E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	5	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note A prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk score can be calculated here: <https://www.mdcalc.com/imdc-international-metastatic-ccc-database-consortium-risk-model-metastatic-renal-cell-carcinoma>.

One point is assigned for each of:

- (i) a time of diagnosis to systemic therapy of less than 1 year
- (ii) a Karnofsky Performance Status of less than 80%
- (iii) a haemoglobin less than the lower limit of normal
- (iv) a corrected calcium level greater than the upper limit of normal
- (v) a neutrophil count greater than the upper limit of normal
- (vi) a platelet count greater than the upper limit of normal

Stated normal reference ranges may vary depending on the laboratory providing the measurement. 'Normal' here refers to the individual laboratory's stated normal reference range.

Favourable IMDC risk is a score of 0.

Intermediate IMDC risk is a score of 1 to 2.

Poor IMDC risk is a score of 3 to 6.

Document any IMDC risk score assessment in the patient's medical records.

Authority required (STREAMLINED)

14001

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must not have previously been treated, **AND**
- Patient must have a prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk classification score at treatment initiation with this drug of either: (i) 1 to 2 (intermediate risk), (ii) 3 to 6 (poor risk); document the IMDC risk classification score in the patient's medical records, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must be in combination with PBS-subsidised treatment with ipilimumab as induction for this condition. Induction treatment with nivolumab must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks. The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

11627Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	360 mg	3	..	*7333.85	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Resected Stage IIIB, IIIC, IIID or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be adjuvant to complete surgical resection, **AND**
- Patient must have a WHO performance status of 1 or less, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- The treatment must commence within 12 weeks of complete resection, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy. Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required

Resected Stage IIIB, IIIC, IIID or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for adjuvant treatment following complete surgical resection, **AND**
- Patient must not have experienced disease recurrence, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy. Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11906P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	5	..	*9734.47	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

10705

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease.

Injection

10424P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	7	..	*7886.47	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

10701

Unresectable Stage III or Stage IV malignant melanoma
Treatment Phase: Continuing treatment - 6 weekly treatment regimen

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease.

Injection

12123C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	3	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14727

Stage II or Stage III triple negative breast cancer

Clinical criteria:

- The treatment must be initiated in combination with neoadjuvant chemotherapy, **AND**
- The condition must not have progressed/recurred whilst on treatment with this drug.

Treatment criteria:

- Patient must not be undergoing treatment with this drug beyond 52 cumulative weeks under this restriction, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 7 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 4 repeat prescriptions.

Injection

13739D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	7	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

14818

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
- Patient must not have experienced disease progression whilst on adjuvant PD-1 inhibitor treatment or disease recurrence within 6 months of completion of adjuvant PD-1 inhibitor treatment if treated for resected Stage IIIB, IIIC, IIID or IV melanoma, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed a total of 6 doses under this restriction.

Injection

10475H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	5	..	*7886.47	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

14817

Unresectable Stage III or Stage IV malignant melanoma
Treatment Phase: Initial treatment - 6 weekly treatment regimen

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
- Patient must not have experienced disease progression whilst on adjuvant PD-1 inhibitor treatment or disease recurrence within 6 months of completion of adjuvant PD-1 inhibitor treatment if treated for resected Stage IIIB, IIIC, IIID or IV melanoma, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed a total of 3 doses under this restriction.

Injection

12122B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	2	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note A Combined Positive Score (CPS) is determined by:

The number of PD-L1-stained cells (tumour cells, lymphocytes, macrophages) divided by the number of all viable tumour cells (i.e. the total number of: PD-L1-positive tumour cells plus PD-L1-negative tumour cells).

Although the result of the CPS calculation can exceed 100, the maximum score is defined as CPS 100.

A minimum of 100 viable tumour cells in the PD-L1-stained slide is required for the specimen to be considered adequate for PD-L1 evaluation.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**14324**

Recurrent, unresectable or metastatic triple negative breast cancer

Clinical criteria:

- The condition must have been (up until this drug therapy) untreated in the unresectable/metastatic disease stage, **AND**
- The condition must have been (up until this drug therapy) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy in breast cancer, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation, **AND**
- The treatment must be in combination with chemotherapy, **AND**
- The condition must have both: (i) programmed cell death ligand 1 (PD-L1) expression confirmed by a validated test, (ii) a Combined Positive Score (CPS) of at least 10 at treatment initiation.

Treatment criteria:

- Patient must be undergoing initial treatment with this drug - this is the first prescription for this drug; OR
- Patient must be undergoing continuing treatment with this drug - both the following are true: (i) the condition has not progressed on active treatment with this drug, (ii) this prescription does not extend PBS subsidy beyond 24 cumulative months from the first administered dose, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Injection

13626E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)**13726**

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have undergone an autologous stem cell transplant (ASCT) for this condition and have experienced relapsed or refractory disease post ASCT; OR
- Patient must not be suitable for ASCT for this condition and have experienced relapsed or refractory disease following at least 2 prior treatments for this condition, **AND**

- Patient must not have received prior treatment with a PD-1 (programmed cell death-1) inhibitor for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required (STREAMLINED)

13741

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

11352L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13727

Relapsed or refractory primary mediastinal B-cell lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as primary mediastinal B-cell lymphoma through histological investigation combined with at least one of: (i) positron emission tomography - computed tomography (PET-CT) scan, (ii) PET scan, (iii) CT scan, **AND**

- Patient must have been treated with rituximab-based chemotherapy for this condition, **AND**
- Patient must be experiencing relapsed/refractory disease, **AND**
- Patient must be autologous stem cell transplant (ASCT) ineligible following a single line of treatment; OR
- Patient must have undergone an autologous stem cell transplant (ASCT); OR
- Patient must have been treated with at least 2 chemotherapy treatment lines for this condition, one of which must include rituximab-based chemotherapy, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required (STREAMLINED)

13732

Relapsed or refractory primary mediastinal B-cell lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR

- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

12126F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13431

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must not exceed a total of 7 doses under this restriction.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13432

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a total of 35 cycles or up to 24 months of treatment under both initial and continuing treatment restrictions, whichever comes first.

Injection

11492W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	6	..	*7886.47	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13739

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The condition must have progressed on or after prior platinum based chemotherapy; OR
- The condition must have progressed on or within 12 months of completion of adjuvant platinum-containing chemotherapy following cystectomy for localised muscle-invasive urothelial cancer; OR
- The condition must have progressed on or within 12 months of completion of neoadjuvant platinum-containing chemotherapy prior to cystectomy for localised muscle-invasive urothelial cancer, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13736

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

11632F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13436

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment - 6 weekly treatment regimen

Clinical criteria:

- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must not exceed a total of 4 doses under this restriction.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13437

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment - 6 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a total of 18 cycles or up to 24 months of treatment under both initial and continuing treatment restrictions, whichever comes first.

Injection

12121Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	3	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Initial treatment - 6 weekly treatment regimen

Clinical criteria:

- The treatment must be adjuvant to complete surgical resection, **AND**
- Patient must have a WHO performance status of 1 or less, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- The treatment must commence within 12 weeks of complete resection, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Continuing treatment - 6 weekly treatment regimen

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for adjuvant treatment following complete surgical resection, **AND**
- Patient must not have experienced disease recurrence, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

Injection

12125E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	3	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Unresectable or metastatic deficient mismatch repair (dMMR) colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be untreated for this PBS indication (i.e untreated for each of: (i) unresectable disease, (ii) metastatic disease), **AND**
- Patient must not have received prior treatment for colorectal cancer with each of: (i) a programmed cell death-1 (PD-1) inhibitor, (ii) a programmed cell death ligand-1 (PD-L1) inhibitor, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must have deficient mismatch repair (dMMR) colorectal cancer, as determined by immunohistochemistry test.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required

Unresectable or metastatic deficient mismatch repair (dMMR) colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR

- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

12605K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13735

Recurrent or metastatic squamous cell carcinoma of the oral cavity, pharynx or larynx

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be incurable by local therapies in the locally advanced setting, **AND**
- Patient must not have had systemic therapy for this condition in the recurrent or metastatic setting prior to initiating PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have experienced disease recurrence within 6 months of completion of systemic therapy if previously treated in the locally advanced setting, **AND**
- Patient must have had a WHO performance status of 0 or 1, **AND**
- The treatment must be either: (i) the sole PBS-subsidised therapy where the condition expresses programmed cell death ligand 1 (PD-L1) with a combined positive score (CPS) greater than or equal to 20 in the tumour sample, (ii) in combination with platinum-based chemotherapy, unless contraindicated or not tolerated.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13731

Recurrent or metastatic squamous cell carcinoma of the oral cavity, pharynx or larynx

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

13114F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13948

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk classification score at treatment initiation with this drug of either: (i) 1 to 2 (intermediate risk), (ii) 3 to 6 (poor risk); document the IMDC risk classification score in the patient's medical records, **AND**
- The condition must be untreated, **AND**
- Patient must have a WHO performance status of 2 or less.

Treatment criteria:

- Patient must be undergoing combination therapy consisting of: (i) pembrolizumab, (ii) lenvatinib; OR
- Patient must be undergoing monotherapy with this drug due to a contraindication/intolerance to the other drug in the combination mentioned above, requiring temporary/permanent discontinuation; document the details in the patient's medical records, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note A prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk score can be calculated here: <https://www.mdcalc.com/imdc-international-metastatic-rcc-database-consortium-risk-model-metastatic-renal-cell-carcinoma>.

One point is assigned for each of:

- a time of diagnosis to systemic therapy of less than 1 year
- a Karnofsky Performance Status of less than 80%
- a haemoglobin less than the lower limit of normal
- a corrected calcium level greater than the upper limit of normal
- a neutrophil count greater than the upper limit of normal
- a platelet count greater than the upper limit of normal

Stated normal reference ranges may vary depending on the laboratory providing the measurement. 'Normal' here refers to the individual laboratory's stated normal reference range.

Favourable IMDC risk is a score of 0.

Intermediate IMDC risk is a score of 1 to 2.

Poor IMDC risk is a score of 3 to 6.

Document any IMDC risk score assessment in the patient's medical records.

Authority required (STREAMLINED)

13949

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing combination therapy consisting of: (i) pembrolizumab, (ii) lenvatinib; OR
- Patient must be undergoing monotherapy with this drug due to a contraindication/intolerance to the other drug in the combination mentioned above, requiring temporary/permanent discontinuation; document the details in the patient's medical records, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

13267G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- The treatment must be in addition to complete surgical resection, **AND**
- Patient must have a WHO performance status of 1 or less, **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- The treatment must commence within 12 weeks of complete resection, **AND**
- Patient must not have received more than 12 months of therapy (irrespective of whether therapy has been partly PBS-subsidised/non-PBS-subsidised).

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note Commencing treatment within 12 weeks of complete resection means either 12 weeks after resection or 12 weeks prior to resection.

Where non-PBS-subsidised supply has occurred, the total amount of PBS-subsidised supply is intended to be the balance of 18 doses less the number of non-PBS-subsidised doses.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Treatment criteria:

- Patient must be undergoing continuing PBS-subsidised treatment commenced through an 'Initial treatment' listing.

Clinical criteria:

- Patient must not have experienced disease recurrence, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received more than 12 months of therapy (irrespective of whether therapy has been partly PBS-subsidised/non-PBS-subsidised).

Injection

12120X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	7	..	*7886.47	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14027

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have received prior treatment with platinum-based chemotherapy, **AND**
- The condition must be untreated with each of: (i) programmed cell death-1/ligand-1 (PD-1/PDL-1) inhibitor therapy, (ii) tyrosine kinase inhibitor therapy, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation.

Treatment criteria:

- Patient must be undergoing combination therapy consisting of: (i) pembrolizumab, (ii) lenvatinib; OR
- Patient must be undergoing monotherapy with this drug due to a contraindication/intolerance to the other drug in the combination mentioned above, requiring temporary/permanent discontinuation; document the details in the patient's medical records, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

14044

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing combination therapy consisting of: (i) pembrolizumab, (ii) lenvatinib; OR
- Patient must be undergoing monotherapy with this drug due to a contraindication/intolerance to the other drug in the combination mentioned above, requiring temporary/permanent discontinuation; document the details in the patient's medical records, **AND**

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Authority required (STREAMLINED)

14028

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply - Grandfather arrangements

Clinical criteria:

- Patient must have received non-PBS-subsidised treatment with this drug for this condition prior to 1 June 2023, **AND**
- The treatment must be occurring in a patient where each of the following is true: (i) the patient had received prior treatment with platinum-based chemotherapy, (ii) the patient was untreated at treatment initiation with each of: (a) programmed cell death-1/ligand-1 (PD-1/PDL-1) inhibitor therapy, (b) tyrosine kinase inhibitor therapy, (iii) the patient's WHO performance status was no higher than 1 at treatment initiation, (iv) this drug is being prescribed in either: (a) a combination of pembrolizumab plus lenvatinib only, (b) as monotherapy where there was a contraindication/intolerance to the other drug in the combination - document the details in the patient's medical records, (v) disease progression has not occurred whilst on treatment, (vi) this prescription does not extend treatment beyond 24 months from the first administered dose.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Injection

13287H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14403

Advanced carcinoma of the cervix

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be at least one of (i) persistent carcinoma, (ii) recurrent carcinoma, (iii) metastatic carcinoma of the cervix, **AND**
- The condition must be unsuitable for curative treatment with either of (i) surgical resection, (ii) radiation, **AND**
- Patient must have WHO performance status no higher than 1, **AND**
- Patient must not have received prior treatment for this PBS indication.

Treatment criteria:

- Patient must be undergoing concomitant treatment with chemotherapy, containing a minimum of: (i) a platinum-based chemotherapy agent, plus (ii) paclitaxel, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required (STREAMLINED)

14404

Advanced carcinoma of the cervix

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a total of (i) 24 months, (ii) 35 doses (based on a 3-weekly dose regimen), (iii) 17 doses (based on a 6-weekly dose regimen) whichever comes first from the first dose of this drug regardless if it was PBS/non-PBS subsidised.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required (STREAMLINED)

14405

Advanced carcinoma of the cervix

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply - Grandfather arrangements

Clinical criteria:

- Patient must be currently receiving non-PBS-subsidised treatment with this drug for this condition, with treatment having commenced prior to 1 October 2023, **AND**
- Patient must have met all other PBS eligibility criteria that a non-Grandfather patient would ordinarily be required to meet, meaning that at the time non-PBS supply was commenced, the patient: (i) had either one of (1) persistent carcinoma, (2) recurrent carcinoma, (3) metastatic carcinoma of the cervix; (ii) had a WHO performance status no higher than 1; (iii) was unsuitable for curative treatment with either of (1) surgical resection, (2) radiation; (iv) had not received prior treatment for this PBS indication; (v) was treated concomitantly with platinum-based chemotherapy agent, plus paclitaxel, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a total of (i) 24 months, (ii) 35 doses (based on a 3-weekly dose regimen), (iii) 17 doses (based on a 6-weekly dose regimen) whichever comes first from the first dose of this drug regardless if it was PBS/non-PBS subsidised.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Injection

13645E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15641.03	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

VEGF/VEGFR (Vascular Endothelial Growth Factor) inhibitors

▪ **BEVACIZUMAB**

Injection

12508H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1800 mg	7	..	*1780.35	31.60	Abevmy [SZ] (bevacizumab 100 mg/4 mL injection, 4 mL vial) Abevmy [SZ] (bevacizumab 400 mg/16 mL injection, 16 mL vial) Bevaciptin [LR] (bevacizumab 100 mg/4 mL injection, 4 mL vial) Bevaciptin [LR] (bevacizumab 400 mg/16 mL injection, 16 mL vial) Mvasi [AN] (bevacizumab 100 mg/4 mL injection, 4 mL vial) Mvasi [AN] (bevacizumab 400 mg/16 mL injection, 16 mL vial)

Other monoclonal antibodies and antibody drug conjugates

▪ **BLINATUMOMAB**

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome, neurological toxicities and reactivation of John Cunningham virus (JC) viral infection.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be relapsed or refractory B-precursor cell ALL, with an Eastern Cooperative Oncology Group (ECOG) performance status of 2 or less, **AND**
- The condition must not be present in the central nervous system or testis, **AND**
- Patient must have previously received a tyrosine kinase inhibitor (TKI) if the condition is Philadelphia chromosome positive, **AND**
- Patient must have received intensive combination chemotherapy for initial treatment of ALL or for subsequent salvage therapy, **AND**
- Patient must not have received more than 1 line of salvage therapy, **AND**
- The condition must be one of the following: (i) untreated with this drug for measurable residual disease, (ii) treated with this drug for measurable residual disease, but the condition has not relapsed within 6 months of completing that course of treatment, **AND**
- The condition must have more than 5% blasts in bone marrow, **AND**
- The treatment must not be more than 2 treatment cycles under this restriction in a lifetime.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 9 days of the first cycle and the first 2 days of the second cycle. For all subsequent cycle starts and re-initiation (e.g. if treatment is interrupted for 4 or more hours), supervision by a health care professional or hospitalisation is recommended.

An amount of 651 microgram will be sufficient for a continuous infusion of blinatumomab over 28 days in cycle 1. An amount of 784 microgram, which may be obtained under Induction treatment - balance of supply restriction, will be sufficient for a continuous infusion of blinatumomab over 28 days in cycle 2.

Blinatumomab is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Acute Lymphoblastic Leukaemia PBS Authority Application - Supporting Information Form; and
- (3) date of most recent chemotherapy, and if this was the initial chemotherapy regimen or salvage therapy, including what line of salvage; and
- (4) if applicable, the date of completion of blinatumomab treatment for measurable residual disease and the date of the patient's subsequent relapse; and
- (5) the percentage blasts in bone marrow count that is no more than 4 weeks old at the time of application.

Injection

11116C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	651 mcg	*67287.75	31.60	Blinicyto [AN] (blinatumomab 38.5 microgram injection [1 vial] (&) inert substance solution [10 mL vial], 1 pack)

■ BLINATUMOMAB

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome, neurological toxicities and reactivation of John Cunningham virus (JC) viral infection.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Induction treatment - balance of supply

Clinical criteria:

- The condition must be relapsed or refractory B-precursor cell ALL, with an Eastern Cooperative Oncology Group (ECOG) performance status of 2 or less, **AND**
- The condition must not be present in the central nervous system or testis, **AND**
- Patient must have previously received a tyrosine kinase inhibitor (TKI) if the condition is Philadelphia chromosome positive, **AND**
- Patient must have received insufficient therapy with this agent for this condition under the Induction treatment restriction to complete a maximum of 2 treatment cycles in a lifetime.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 9 days of the first cycle and the first 2 days of the second cycle. For all subsequent cycle starts and re-initiation (e.g. if treatment is interrupted for 4 or more hours), supervision by a health care professional or hospitalisation is recommended.

An amount of 784 mcg will be sufficient for a continuous infusion of blinatumomab over 28 days in cycle 2.

Blinatumomab is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Injection

11119F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	784 mcg	*78480.39	31.60	Blinicyto [AN] (blinatumomab 38.5 microgram injection [1 vial] (&) inert substance solution [10 mL vial], 1 pack)

■ BLINATUMOMAB

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome, neurological toxicities and reactivation of John Cunningham virus (JC) viral infection.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note A complete remission is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a full recovery of peripheral blood counts with platelets of greater than 100,000 per microliter, and absolute neutrophil count (ANC) of greater than 1,000 per microliter.

Note A complete remission with partial haematological recovery is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a partial recovery of peripheral blood counts with platelets of greater than 50,000 per microliter, and absolute neutrophil count (ANC) of greater than 500 per microliter.

Note Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time when an assessment is required must cease PBS-subsidised therapy with this agent.

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Consolidation treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised induction treatment with this drug for this condition, **AND**
- Patient must have achieved a complete remission; OR
- Patient must have achieved a complete remission with partial haematological recovery, **AND**
- The treatment must not be more than 3 treatment cycles under this restriction in a lifetime, **AND**
- Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug.

Injection

11115B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	784 mcg	2	..	*78480.39	31.60	Blinicyto [AN] (blinatumomab 38.5 microgram injection [1 vial] (&) inert substance solution [10 mL vial], 1 pack)

▪ **BLINATUMOMAB**

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome, neurological toxicities and reactivation of John Cunningham virus (JC) viral infection.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note A complete remission is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a full recovery of peripheral blood counts with platelets of greater than 100,000 per microliter, and absolute neutrophil count (ANC) of greater than 1,000 per microliter.

Authority required

Measurable residual disease of precursor B-cell acute lymphoblastic leukaemia (Pre-B-cell ALL)

Treatment Phase: Initial treatment of measurable residual disease of Pre-B-cell ALL

Treatment criteria:

- Must be treated by a physician experienced in the treatment of haematological malignancies.

Clinical criteria:

- Patient must have an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1, **AND**
- The condition must not be present in the central nervous system or testis, **AND**
- Patient must have achieved complete remission following intensive combination chemotherapy for initial treatment of acute lymphoblastic leukaemia (ALL) or for subsequent salvage therapy, **AND**
- Patient must have measurable residual disease based on measurement in bone marrow, documented after an interval of at least 2 weeks from the last course of systemic chemotherapy given as intensive combination chemotherapy treatment of ALL/as subsequent salvage therapy, whichever was the later, measured using flow cytometry/molecular methods,

AND

- The treatment must not be more than 2 treatment cycles under this restriction in a lifetime.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 days of the first cycle and the first 2 days of the second cycle.

For all subsequent cycle starts and re-initiation (e.g. if treatment is interrupted for four or more hours), supervision by a health care professional or hospitalisation is recommended.

An amount of 784 mcg will be sufficient for a continuous infusion of blinatumomab over 28 days in each cycle.

Blinatumomab is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Measurable residual disease positive Acute Lymphoblastic Leukaemia PBS Authority Application - Supporting Information Form; and
- (3) date of most recent chemotherapy, and if this was the initial chemotherapy regimen or salvage therapy; and
- (4) the percentage blasts in bone marrow count that is no more than 4 weeks old at the time of application.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Measurable residual disease of precursor B-cell acute lymphoblastic leukaemia (Pre-B-cell ALL)
Treatment Phase: Continuing treatment of previously measurable residual disease of Pre-B-cell ALL

Treatment criteria:

- Must be treated by a physician experienced in the treatment of haematological malignancies.

Clinical criteria:

- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition, **AND**
- Patient must have achieved a complete remission, **AND**
- The condition must be negative for measurable residual disease using the same method used to determine initial PBS eligibility, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition, **AND**
- The treatment must not be more than 2 treatment cycles under this restriction in a lifetime.

For all subsequent cycle starts and re-initiation (e.g. if treatment is interrupted for four or more hours), supervision by a health care professional or hospitalisation is recommended.

An amount of 784 microgram will be sufficient for a continuous infusion of blinatumomab over 28 days in each cycle.

Blinatumomab is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Injection

11867N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	784 mcg	1	..	*78480.39	31.60	Blinicyto [AN] (blinatumomab 38.5 microgram injection [1 vial] (&) inert substance solution [10 mL vial], 1 pack)

■ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

CD30 positive systemic anaplastic large cell lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be for curative intent, **AND**
- Patient must have undergone appropriate prior front-line curative intent chemotherapy, **AND**
- Patient must demonstrate relapsed or chemotherapy-refractory disease, **AND**
- Patient must have responded to PBS-subsidised treatment with this drug if previously used for initial treatment of CD30 positive peripheral T-cell lymphoma, non-cutaneous type, **AND**
- The treatment must not exceed 4 cycles under this restriction.

Applications for authorisation of initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail and must include:

- (a) details (date, unique identifying number or provider number) of a histology report showing evidence of the tumour's CD30 positivity; and
- (b) The date of initial diagnosis of systemic anaplastic large cell lymphoma; and
- (c) Dates of commencement and completion of front-line curative intent chemotherapy; and

(d) a declaration of whether the patient's disease is relapsed or refractory, and the date and means by which the patient's disease was assessed as being relapsed or refractory.

All reports must be documented in the patient's medical records.

If the application is submitted through HPOS form upload or mail, it must include:

(i) A completed authority prescription form; and

(ii) A completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

10172J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	3	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

CD30 positive systemic anaplastic large cell lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed 12 cycles under this restriction in a lifetime.

Injection

10180T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	11	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have undergone a primary autologous stem cell transplant (ASCT) for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 12 cycles of treatment under this restriction.
The treatment must not exceed a total of 16 cycles of combined initial and continuing treatment in a lifetime.

Injection

11067L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	11	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Relapsed or Refractory Hodgkin lymphoma
Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have undergone an autologous stem cell transplant (ASCT) for this condition, **AND**
- Patient must not be suitable for ASCT for this condition; OR
- Patient must not be suitable for treatment with multi-agent chemotherapy for this condition, **AND**
- Patient must have experienced a relapsed CD30+ Hodgkin lymphoma following at least two prior treatments for this condition; OR
- Patient must have experienced a refractory CD30+ Hodgkin lymphoma following at least two prior treatments for this condition, **AND**
- Patient must not receive more than 4 cycles of treatment under this restriction.

Applications for authorisation of initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail.

If the application is submitted through HPOS upload or mail, it must include:

- (a) a completed authority prescription form; and
- (b) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
11080E	200 mg	3	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed or Refractory Hodgkin lymphoma
Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must not have undergone an autologous stem cell transplant (ASCT) for this condition, **AND**
- Patient must not be suitable for ASCT for this condition; OR
- Patient must not be suitable for treatment with multi-agent chemotherapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 12 cycles of treatment under this restriction.

The treatment must not exceed a total of 16 cycles of combined initial and continuing treatment in a lifetime.

Injection

	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
11086L	200 mg	11	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)
Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have undergone a primary autologous stem cell transplant (ASCT), **AND**
- Patient must have experienced a relapsed CD30+ Hodgkin lymphoma post ASCT; OR
- Patient must have experienced a refractory CD30+ Hodgkin lymphoma post ASCT, **AND**
- Patient must not receive more than 4 cycles of treatment under this restriction.

Applications for authorisation of initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail.

If the application is submitted through HPOS upload or mail, it must include:

- (a) a completed authority prescription form; and
- (b) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

11089P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	3	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

CD30 positive cutaneous T-cell lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have pathologically confirmed CD30 positive cutaneous T-cell lymphoma, **AND**
- Patient must have CD30 positivity of at least 3% of malignant cells, **AND**
- Patient must have a diagnosis of mycosis fungoides; OR
- Patient must have a diagnosis of Sezary syndrome; OR
- Patient must have a diagnosis of primary cutaneous anaplastic large cell lymphoma, **AND**
- Patient must have received prior systemic treatment for this condition, **AND**
- The condition must be relapsed or refractory, **AND**
- The treatment must not exceed 4 cycles under this restriction in a lifetime, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

The authority application must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail and must include:

- (a) details (date, unique identifying number/code or provider number) of the histopathology report from an Approved Pathology Authority demonstrating the patient has a diagnosis of either mycosis fungoides, Sezary syndrome or primary cutaneous anaplastic large cell lymphoma; and
- (b) details (date, unique identifying number/code or provider number) of a histology report on the tumour sample or of a flow cytometric analysis of lymphoma cells of the blood showing CD30 positivity of at least 3% of malignant cells; and
- (c) Date of commencement and completion of the most recent prior systemic treatment.

All reports must be documented in the patient's medical records.

If the application is submitted through HPOS form upload or mail, it must include:

- (i) A completed authority prescription form; and
- (ii) A completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

11651F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	180 mg	3	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

CD30 positive cutaneous T-cell lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have achieved an objective response with this drug, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The treatment must not exceed 12 cycles under this restriction in a lifetime.

An objective response is defined as the demonstration of response by clinical observation of skin lesions, or response by positron-emission tomography (PET) and/or computed tomography (CT) standard criteria.

Injection

11661R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	180 mg	11	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note This product is not PBS-subsidised for the treatment of previously untreated CD30 positive cutaneous T-cell lymphoma.

Authority required

CD30 positive peripheral T-cell lymphoma, non-cutaneous type

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with cyclophosphamide, doxorubicin and prednisone, **AND**
- Patient must have completed 6 initial cycles of PBS-subsidised treatment with this drug for this indication, **AND**
- Patient must have achieved at least a partial response to the 6 initial cycles of treatment with a combination of this drug and cyclophosphamide, doxorubicin and prednisone for this indication, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be more than 2 treatment cycles under this restriction in a lifetime.

Partial response is defined using Lugano Response Criteria for Non-Hodgkin Lymphoma as:

- (a) Positron emission tomography-based response: lymph nodes and extralymphatic sites - a score of 4 (uptake moderately > liver), or 5 (uptake markedly higher than liver and/or new lesions), with reduced uptake compared with baseline and residual mass(es) of any size; non-measured lesions - not applicable; organ enlargement - not applicable; new lesions - none; bone marrow - residual uptake higher than uptake in normal marrow but reduced compared with baseline (diffuse uptake compatible with reactive changes from chemotherapy allowed). If there are persistent focal changes in the marrow in the context of a nodal response, consideration should be given to further evaluation with MRI or biopsy or an interval scan; OR
- (b) Computed tomography-based response: lymph nodes and extralymphatic sites - greater than or equal to 50% decrease in the sum of the product of the perpendicular diameters for multiple lesions, of up to six (6) target measurable nodes and extranodal sites; non-measured lesions - absent/normal, regressed but no increase; new lesions - none; bone marrow - not applicable.

Injection

12632W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	1	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note This product is not PBS-subsidised for the treatment of previously untreated CD30 positive cutaneous T-cell lymphoma.

Authority required

CD30 positive peripheral T-cell lymphoma, non-cutaneous type

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have histological confirmation of CD30 expression in at least 3% of malignant cells, **AND**
- The treatment must be for first line therapy for this condition, **AND**
- The treatment must be for curative intent, **AND**
- The treatment must be in combination with cyclophosphamide, doxorubicin and prednisone, **AND**
- The treatment must not be more than 6 treatment cycles under this restriction in a lifetime.

Applications for authorisation of initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail and must include:

- (a) details (date, unique identifying number/code or provider number) of a histology report on the tumour sample from an Approved Pathology Authority showing CD30 positivity of at least 3% malignant cells; and
- (b) The date of initial diagnosis of Peripheral T-cell lymphoma.

All reports must be documented in the patient's medical records.

If the application is submitted through HPOS form upload or mail, it must include:

- (i) A completed authority prescription form; and
- (ii) A completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

12656D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	5	..	*19012.59	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **ELOTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with lenalidomide and dexamethasone, **AND**
- Patient must have progressive disease after at least one prior therapy, **AND**
- Patient must have undergone or be ineligible for a stem cell transplant, **AND**
- Patient must not have previously received this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

12990Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	9	..	*5151.24	31.60	Empliciti [BQ] (elotuzumab 300 mg injection, 1 vial) Empliciti [BQ] (elotuzumab 400 mg injection, 1 vial)

▪ **ELOTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Relapsed and/or refractory multiple myeloma
Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - The treatment must be in combination with lenalidomide and dexamethasone, **AND**
 - Patient must not have developed disease progression while receiving treatment with this drug for this condition. Progressive disease is defined as at least 1 of the following:
 - (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
 - (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
 - (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
 - (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
 - (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
 - (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
 - (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).
- Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

12995Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	5	..	*5151.24	31.60	Empliciti [BQ] (elotuzumab 300 mg injection, 1 vial) Empliciti [BQ] (elotuzumab 400 mg injection, 1 vial)

▪ **ENFORTUMAB VEDOTIN**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14416

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Clinical criteria:

- The condition must have progressed on/following both: (i) platinum-based chemotherapy, (ii) programmed cell death 1/ligand 1 (PD-1/PD-L1) inhibitor therapy; OR
- The condition must have progressed on/following platinum-based chemotherapy, whilst PD-1/PD-L1 inhibitor therapy resulted in an intolerance that required treatment cessation, **AND**
- Patient must have/have had a WHO performance status score of no greater than 1 at treatment initiation with this drug. **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this PBS indication.

Treatment criteria:

- Patient must be undergoing treatment with this drug for the first time; OR
- Patient must be undergoing continuing treatment with this drug, with each of the following being true: (i) all other PBS eligibility criteria in this restriction are met, (ii) disease progression is absent.

Injection

13634N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	125 mg	8	..	*6341.02	31.60	Padcev [LL] (enfortumab vedotin 20 mg injection, 1 vial) Padcev [LL] (enfortumab vedotin 30 mg injection, 1 vial)

▪ **GEMTUZUMAB OZOGAMICIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Acute Myeloid Leukaemia
Treatment Phase: Induction treatment

Clinical criteria:

- Patient must have confirmed CD33-positive AML prior to initiation of treatment, **AND**
- The condition must be de novo, **AND**
- The condition must be previously untreated at the time of initiation (except for prior essential treatment with hydroxyurea or leukapheresis for patients with hyperleukocytic AML), **AND**

- Patient must have confirmed intermediate/favourable cytogenetic risk; OR
- Patient must have unknown cytogenetic risk due to inconclusive test results, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score of 2 or less, **AND**
- The condition must not be acute promyelocytic leukaemia, **AND**
- The treatment must be in combination with standard intensive remission induction chemotherapy for this condition, which must include cytarabine and an anthracycline, **AND**
- The treatment must not be used in combination with a tyrosine kinase inhibitor, **AND**
- The condition must not be internal tandem duplication (ITD) or tyrosine kinase domain (TKD) FMS tyrosine kinase 3 (FLT3) mutation positive, **AND**
- Patient must not receive more than 1 induction cycle under this restriction in a lifetime.
This drug is not PBS-subsidised if it is prescribed to an in-patient in a public hospital setting.

Injection

12878T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	5 mg	2	..	*9427.25	31.60	Mylotarg [PF] (gemtuzumab ozogamicin 5 mg injection, 1 vial)

▪ **GEMTUZUMAB OZOGAMICIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Consolidation treatment

Clinical criteria:

- Patient must have achieved a complete remission following induction treatment with this drug for this condition, **AND**
- The treatment must be in combination with standard intensive remission consolidation chemotherapy for this condition, which must include cytarabine and an anthracycline, **AND**
- Patient must not receive more than 2 consolidation cycles under this restriction in a lifetime.

This drug is not PBS-subsidised if it is prescribed to an in-patient in a public hospital setting.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Complete remission following induction is defined as fewer than 5% blasts in a normocellular marrow and an absolute neutrophil count of more than 1.0×10^9 cells/L with a platelet count of 100×10^9 /L or more in the peripheral blood in the absence of transfusion.

Progressive disease is defined as the presence of any of the following:

- a) Leukaemic cells in the CSF;
- b) Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- c) Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- d) Extramedullary leukaemia.

Injection

12904E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	5 mg	1	..	*9427.25	31.60	Mylotarg [PF] (gemtuzumab ozogamicin 5 mg injection, 1 vial)

▪ **IPILIMUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

11478

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial combination treatment (with nivolumab) as first-line drug therapy

Clinical criteria:

- The condition must be squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must not have previously been treated for this condition in the metastatic setting, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must be in combination with platinum-based chemotherapy for the first two cycles, **AND**
- The treatment must be in combination with nivolumab.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)**11391**

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing combination treatment (with nivolumab) of first-line drug therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed 24 months in total, measured from the initial dose, or, must not extend beyond disease progression, whichever comes first, **AND**
- The treatment must be in combination with nivolumab.

Injection

12304N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	4	..	*16390.23	31.60	Yervoy [BQ] (ipilimumab 50 mg/10 mL injection, 10 mL vial)

■ IPILIMUMAB

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)**11930**

Unresectable malignant mesothelioma

Clinical criteria:

- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be in combination with PBS-subsidised nivolumab for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a maximum total of 24 months in a lifetime for this condition.

Injection

12601F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	3	..	*16390.23	31.60	Yervoy [BQ] (ipilimumab 50 mg/10 mL injection, 10 mL vial)

■ IPILIMUMAB

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note A prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk score can be calculated here: <https://www.mdcalc.com/imdc-international-metastatic-renal-cell-carcinoma>.

One point is assigned for each of:

- a time of diagnosis to systemic therapy of less than 1 year
- a Karnofsky Performance Status of less than 80%
- a haemoglobin less than the lower limit of normal
- a corrected calcium level greater than the upper limit of normal
- a neutrophil count greater than the upper limit of normal
- a platelet count greater than the upper limit of normal

Stated normal reference ranges may vary depending on the laboratory providing the measurement. 'Normal' here refers to the individual laboratory's stated normal reference range.

Favourable IMDC risk is a score of 0.

Intermediate IMDC risk is a score of 1 to 2.

Poor IMDC risk is a score of 3 to 6.

Document any IMDC risk score assessment in the patient's medical records.

Authority required (STREAMLINED)**8555**

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must not have previously been treated, **AND**

- The condition must be classified as intermediate to poor risk according to the International Metastatic Renal Cell Carcinoma Database Consortium (IMDC), **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must be in combination with PBS-subsidised treatment with nivolumab as induction therapy for this condition.

Induction treatment with ipilimumab must not exceed a total of 4 doses at a maximum dose of 1 mg per kg every 3 weeks. The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

11644W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	3	..	*16390.23	31.60	Yervoy [BQ] (ipilimumab 50 mg/10 mL injection, 10 mL vial)

▪ **IPIILIMUMAB**

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

6562

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Induction treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
 - Patient must not have received prior treatment with ipilimumab, **AND**
 - The treatment must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks.
- The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Note For patients who commence therapy with ipilimumab:

- Decisions concerning efficacy should await completion of the entire induction regimen (four doses) and should be made in conjunction with established criteria for immunological responses. However induction may be ceased or delayed if symptomatic progressive disease or intolerable adverse events occur and if, in the opinion of the clinician, continuation of treatment poses a risk to the patient;
- Tumour responses may occur beyond the initial 12 week induction phase and evaluation for potential later responses should be undertaken regularly for the first year.

Authority required (STREAMLINED)

6585

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Re-induction treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have progressive disease after achieving an initial objective response to the most recent course of ipilimumab treatment (induction or re-induction), **AND**
- The treatment must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks.

An initial objective response to treatment is defined as either:

- sustained stable disease of greater than or equal to 3 months duration measured from at least 2 weeks after the date of completion of the most recent course of ipilimumab; or
- a partial or complete response.

The patient's body weight must be documented in the patient's medical records at the time treatment with ipilimumab is initiated.

Authority required (STREAMLINED)

14808

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Induction treatment

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
- Patient must have an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1, **AND**
- The condition must not be ocular or uveal melanoma, **AND**
- The treatment must be in combination with PBS-subsidised treatment with nivolumab as induction therapy for this condition.

Induction treatment with nivolumab must not exceed a total of 4 doses at a maximum dose of 1 mg per kg every 3 weeks.

Induction treatment with ipilimumab must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

2638W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	360 mg	3	..	*43487.43	31.60	Yervoy [BQ] (ipilimumab 200 mg/40 mL injection, 40 mL vial) Yervoy [BQ] (ipilimumab 50 mg/10 mL injection, 10 mL vial)

▪ SACITUZUMAB GOVITECAN

Caution This medicine contains a cytotoxic component and causes chemotherapy-like toxicity, in particular, it can cause severe or life-threatening neutropenia and severe diarrhoea. For further information, refer to the Product Information.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12656

Unresectable locally advanced or metastatic triple-negative breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have progressive disease following two or more prior systemic therapies, at least one of them in the locally advanced or metastatic setting, **AND**
- The condition must be inoperable, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation, **AND**
- The treatment must be the sole PBS-subsidised therapy for this PBS indication.

Injection

12944G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	7	..	*10669.50	31.60	Trodely [GI] (sacituzumab govitecan 180 mg injection, 1 vial)

▪ SACITUZUMAB GOVITECAN

Caution This medicine contains a cytotoxic component and causes chemotherapy-like toxicity, in particular, it can cause severe or life-threatening neutropenia and severe diarrhoea. For further information, refer to the Product Information.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12669

Unresectable locally advanced or metastatic triple-negative breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this PBS indication.

Injection

12965J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	13	..	*10669.50	31.60	Trodely [GI] (sacituzumab govitecan 180 mg injection, 1 vial)

OTHER ANTINEOPLASTIC AGENTS

Platinum compounds

▪ CARBOPLATIN

Injection

7222D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	900 mg	5	..	*185.81	31.60	Carboplatin Accord [OC] (carboplatin 450 mg/45 mL injection, 45 mL vial)

▪ CISPLATIN

Injection

7224F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	220 mg	14	..	*181.57	31.60	Cisplatin Accord [OC] (cisplatin 100 mg/100 mL injection, 100 mL vial) Cisplatin Accord [OC] (cisplatin 50 mg/50 mL injection, 50 mL vial)

▪ OXALIPLATIN

Injection

7253R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	300 mg	11	..	*177.44	31.60	Oxaliplatin Accord [OC] (oxaliplatin 100 mg/20 mL injection, 20 mL vial) Oxaliplatin SUN [RA] (oxaliplatin 100 mg/20 mL injection, 20 mL vial) Oxaliplatin SUN [RA] (oxaliplatin 200 mg/40 mL injection, 40 mL vial)

Proteasome inhibitors

▪ **BORTEZOMIB**

Restricted benefit

Multiple myeloma

Restricted benefit

Newly diagnosed systemic light chain amyloidosis

Treatment Phase: Administration on Days 1, 8, 15 and 22 of six treatment cycles (28 days per cycle) in total

Treatment criteria:

- Patient must be undergoing concurrent treatment with PBS-subsidised daratumumab for this PBS indication.

Injection

12219D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3000 mcg	15	..	*210.46	31.60	Bortezom [CR] (bortezomib 3.5 mg injection, 1 vial) Bortezomib Accord [OC] (bortezomib 1 mg injection, 1 vial) Bortezomib Accord [OC] (bortezomib 2.5 mg/mL injection, 1 mL vial) Bortezomib Accord [OC] (bortezomib 3.5 mg injection, 1 vial) Bortezomib Accord [OC] (bortezomib 3.5 mg/1.4 mL injection, 1.4 mL vial) Bortezomib Baxter [BX] (bortezomib 3.5 mg injection, 1 vial) BORTEZOMIB EUGIA [YG] (bortezomib 3.5 mg injection, 1 vial) Bortezomib Ever Pharma [IT] (bortezomib 2.5 mg/mL injection, 1 mL vial) Bortezomib Ever Pharma [IT] (bortezomib 3.5 mg/1.4 mL injection, 1.4 mL vial) Bortezomib Juno [JU] (bortezomib 2.5 mg injection, 1 vial) Bortezomib Juno [JU] (bortezomib 3.5 mg injection, 1 vial) Bortezomib Sandoz [SZ] (bortezomib 3.5 mg injection, 1 vial) BORTEZOMIB-TEVA [TB] (bortezomib 3.5 mg injection, 1 vial) DBL Bortezomib [PF] (bortezomib 1 mg injection, 1 vial) DBL Bortezomib [PF] (bortezomib 2.5 mg injection, 1 vial) DBL Bortezomib [PF] (bortezomib 3 mg injection, 1 vial) DBL Bortezomib [PF] (bortezomib 3.5 mg injection, 1 vial)

▪ **CARFILZOMIB**

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum amount or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12934

Multiple myeloma

Treatment Phase: Initial treatment - twice weekly treatment regimen

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must have progressive disease after at least one prior therapy, **AND**
- Patient must have undergone or be ineligible for a stem cell transplant, **AND**
- Patient must not have previously received this drug for this condition, **AND**
- Patient must not receive more than three cycles of treatment under this restriction.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Authority required (STREAMLINED)

12930

Multiple myeloma

Treatment Phase: Continuing treatment - twice weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must not develop disease progression while receiving treatment with this drug for this condition, **AND**
- Patient must not receive more than 3 cycles of treatment per continuing treatment course authorised under this restriction.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

11230C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	17	..	*2576.65	31.60	Kyprolis [AN] (carfilzomib 10 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 30 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 60 mg injection, 1 vial)

▪ **CARFILZOMIB**

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum amount or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12694

Multiple myeloma

Treatment Phase: Initial treatment - once weekly treatment regimen

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must have progressive disease after at least one prior therapy, **AND**
- Patient must have undergone or be ineligible for a stem cell transplant, **AND**
- Patient must not have previously received this drug for this condition, **AND**
- Patient must not receive more than three cycles of treatment under this restriction.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Authority required (STREAMLINED)

12849

Multiple myeloma

Treatment Phase: Continuing treatment - once weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must not develop disease progression while receiving treatment with this drug for this condition, **AND**
- Patient must not receive more than 3 cycles of treatment per continuing treatment course authorised under this restriction.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or

- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
 - (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
 - (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
 - (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
 - (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
 - (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).
- Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

12243J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	160 mg	8	..	*3391.57	31.60	Kyprolis [AN] (carfilzomib 10 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 30 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 60 mg injection, 1 vial)

▪ **CARFILZOMIB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14389

Relapsed and/or refractory multiple myeloma

Treatment Phase: Initial treatment for Cycles 1 to 3

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with lenalidomide and dexamethasone, **AND**
- Patient must have progressive disease after at least one prior therapy, **AND**
- Patient must not have previously received this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Provide details of the histological diagnosis of multiple myeloma, prior treatments including name(s) of drug(s) and date of the most recent treatment cycle; the basis of the diagnosis of progressive disease or failure to respond; and which disease activity parameters will be used to assess response once only through the Authority application for lenalidomide.

Authority required (STREAMLINED)

14363

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment for Cycles 3 to 12

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with lenalidomide and dexamethasone, **AND**
- Patient must not have progressive disease while receiving treatment with this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or

(g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Authority required (STREAMLINED)

14364

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment for Cycles 13 onwards

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with lenalidomide and dexamethasone, **AND**
- Patient must not have progressive disease while receiving treatment with this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

13637R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	60 mg	17	..	*1354.28	31.60	Kyprolis [AN] (carfilzomib 10 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 30 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 60 mg injection, 1 vial)

Other antineoplastic agents

■ **ARSENIC**

Authority required (STREAMLINED)

6018

Acute promyelocytic leukaemia

Treatment Phase: Induction and consolidation treatment

Clinical criteria:

- The condition must be characterised by the presence of the t(15:17) translocation or PML/RAR-alpha fusion gene transcript.

Injection

10699D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	18 mg	140	..	*250.47	31.60	Arsenic Trioxide Accord [OC] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials) Arsenic Trioxide-AFT [AE] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL ampoules) Arsenic Trioxide Juno [JU] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials) Phenasen [FF] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials)

■ **ARSENIC**

Authority required (STREAMLINED)

4793

Acute promyelocytic leukaemia

Treatment Phase: Induction and consolidation treatment

Clinical criteria:

- The condition must be characterised by the presence of the t(15:17) translocation or PML/RAR-alpha fusion gene transcript, **AND**
- The condition must be relapsed, **AND**
- Patient must be arsenic naive at induction.

Authority required (STREAMLINED)

5997

Acute promyelocytic leukaemia

Clinical criteria:

- The condition must be characterised by the presence of the t(15:17) translocation or PML/RAR-alpha fusion gene transcript.

Injection

7241D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	18 mg	89	..	*250.47	31.60	Arsenic Trioxide Accord [OC] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials) Arsenic Trioxide-AFT [AE] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL ampoules) Arsenic Trioxide Juno [JU] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials) Phenasen [FF] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials)

▪ **ERIBULIN**

Note A patient who has progressive disease with eribulin is no longer eligible for PBS-subsidised eribulin.

Authority required (STREAMLINED)

4649

Locally advanced or metastatic breast cancer

Clinical criteria:

- Patient must have progressive disease, **AND**
- Patient must have failed at least two prior chemotherapeutic regimens for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

10140Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3 mg	13	..	*825.48	31.60	Halaven [EI] (eribulin mesilate 1 mg/2 mL injection, 2 mL vial)

▪ **ERIBULIN**

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7258

Advanced (unresectable and/or metastatic) liposarcoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have an ECOG performance status of 2 or less, **AND**
- The condition must be dedifferentiated, myxoid, round-cell or pleomorphic subtype, **AND**
- Patient must have received prior chemotherapy treatment including an anthracycline and ifosfamide (unless contraindicated) for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Population criteria:

- Patient must be aged 18 years or older.

Authority required (STREAMLINED)

7280

Advanced (unresectable and/or metastatic) liposarcoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop progressive disease while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Population criteria:

- Patient must be aged 18 years or older.

Injection

11199K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3 mg	7	..	*825.48	31.60	Halaven [EI] (eribulin mesilate 1 mg/2 mL injection, 2 mL vial)

▪ **TEBENTAFUSP**

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome (CRS).

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

15085

Advanced (unresectable or metastatic) uveal melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition; OR

- Patient must have previously received inpatient treatment with this drug for this condition in the public hospital setting, **AND**
- Patient must not receive PBS-subsidised treatment with this drug for this condition if it is no longer determined to be clinically beneficial by the treating clinician.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 doses (on Days 1, 8 and 15) and for at least 16 hours after each infusion is completed. If the patient does not experience hypotension that is Grade 2 or worse (requiring medical intervention) with the third dose, subsequent doses can be administered in an appropriate outpatient/ambulatory care setting. Supervision by a health care professional is recommended for a minimum of 30 minutes following each infusion.

Injection

13824N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	136 mcg	7	..	*35791.37	31.60	Kimmtrak [WM] (tebentafusp 100 microgram/0.5 mL injection, 0.5 mL vial)

■ TEBENTAFUSP

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome (CRS).

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Advanced (unresectable or metastatic) uveal melanoma

Treatment Phase: Initial treatment - day 1

Clinical criteria:

- Patient must have HLA-A*02:01-positive disease, **AND**
- Patient must have uveal melanoma that has been confirmed either (i) histologically, (ii) cytologically, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received prior systemic therapy for metastatic disease.

Population criteria:

- Patient must be at least 18 years of age.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 doses (on Days 1, 8 and 15) and for at least 16 hours after each infusion is completed. If the patient does not experience hypotension that is Grade 2 or worse (requiring medical intervention) with the third dose, subsequent doses can be administered in an appropriate outpatient/ambulatory care setting. Supervision by a health care professional is recommended for a minimum of 30 minutes following each infusion.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Positive HLA-A*02:01 assessment must be documented in the patient's medical records.

Injection

13818G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	20 mcg	*17961.64	31.60	Kimmtrak [WM] (tebentafusp 100 microgram/0.5 mL injection, 0.5 mL vial)

■ TEBENTAFUSP

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome (CRS).

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14825

Advanced (unresectable or metastatic) uveal melanoma

Treatment Phase: Initial treatment - day 15

Clinical criteria:

- Patient must have HLA-A*02:01-positive disease, **AND**
- Patient must have previously received PBS-subsidised initial day 8 treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 doses (on Days 1, 8 and 15) and for at least 16 hours after each infusion is completed. If the patient does not experience hypotension that is Grade 2 or worse (requiring medical intervention) with the third dose, subsequent doses can be administered in an appropriate outpatient/ambulatory care setting. Supervision by a health care professional is recommended for a minimum of 30 minutes following each infusion.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Positive HLA-A*02:01 assessment must be documented in the patient's medical records.

Injection

13822L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	68 mcg	*17961.64	31.60	Kimmtrak [WM] (tebentafusp 100 microgram/0.5 mL injection, 0.5 mL vial)

▪ **TEBENTAFUSP**

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome (CRS).

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14821

Advanced (unresectable or metastatic) uveal melanoma

Treatment Phase: Initial treatment - day 8

Clinical criteria:

- Patient must have HLA-A*02:01-positive disease, **AND**
- Patient must have previously received PBS-subsidised initial day 1 treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 doses (on Days 1, 8 and 15) and for at least 16 hours after each infusion is completed. If the patient does not experience hypotension that is Grade 2 or worse (requiring medical intervention) with the third dose, subsequent doses can be administered in an appropriate outpatient/ambulatory care setting. Supervision by a health care professional is recommended for a minimum of 30 minutes following each infusion.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Positive HLA-A*02:01 assessment must be documented in the patient's medical records.

Injection

13833C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	30 mcg	*17961.64	31.60	Kimmtrak [WM] (tebentafusp 100 microgram/0.5 mL injection, 0.5 mL vial)

Chemotherapy items for Public Hospital use

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	82
ANTINEOPLASTIC AGENTS	82
ALKYLATING AGENTS	82
ANTIMETABOLITES	83
PLANT ALKALOIDS AND OTHER NATURAL PRODUCTS	85
CYTOTOXIC ANTIBIOTICS AND RELATED SUBSTANCES	88
MONOCLONAL ANTIBODIES AND ANTIBODY DRUG CONJUGATES	89
OTHER ANTINEOPLASTIC AGENTS	146

■ **ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS**

■ **ANTINEOPLASTIC AGENTS**

ALKYLATING AGENTS

Nitrogen mustard analogues

■ **BENDAMUSTINE**

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7972

Previously untreated stage III or IV mantle cell lymphoma

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be CD20 positive, **AND**
- The treatment must be in combination with rituximab, **AND**
- The condition must be previously untreated, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for induction treatment purposes only, **AND**
- Patient must not receive more than 6 cycles (12 doses) of treatment under this restriction, **AND**
- Patient must not be eligible for stem cell transplantation.

Authority required (STREAMLINED)

7943

Previously untreated stage II bulky or stage III or IV indolent non-Hodgkin's lymphoma

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be CD20 positive, **AND**
- The condition must be previously untreated, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for induction treatment purposes only, **AND**
- The treatment must be in combination with rituximab or obinutuzumab, **AND**
- The treatment must not exceed 6 cycles (12 doses) with this drug under this restriction.

Authority required (STREAMLINED)

7944

Follicular lymphoma

Treatment Phase: Re-induction treatment

Clinical criteria:

- The condition must be CD20 positive, **AND**
- The condition must be refractory to treatment with rituximab for this condition, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for re-induction treatment purposes only, **AND**
- The treatment must be in combination with obinutuzumab, **AND**
- The treatment must not exceed 6 cycles (12 doses) with this drug under this restriction.

The condition is considered rituximab-refractory if the patient experiences less than a partial response or progression of disease within 6 months after completion of a prior rituximab-containing regimen.

Injection

10760H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	11	..	*1239.04	31.60	Bendamustine Juno [JU] (bendamustine hydrochloride 100 mg injection, 1 vial) Bendamustine Juno [JU] (bendamustine hydrochloride 25 mg injection, 1 vial) Bendamustine Sandoz [SZ] (bendamustine hydrochloride 100 mg injection, 1 vial) Bendamustine Sandoz [SZ] (bendamustine hydrochloride 25 mg injection, 1 vial) Bendamustine Viatrix [AF] (bendamustine hydrochloride 100 mg injection, 1 vial) Bendamustine Viatrix [AF] (bendamustine hydrochloride 25 mg injection, 1 vial) Ribomustin [JC] (bendamustine hydrochloride 100 mg injection, 1 vial) Ribomustin [JC] (bendamustine hydrochloride 25 mg injection, 1 vial)

■ **CYCLOPHOSPHAMIDE**

Injection

4327R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	2800 mg	17	..	*144.39	31.60	CYCLOPHOSPHAMIDE-REACH [RQ] (cyclophosphamide 1 g injection, 1 vial) CYCLOPHOSPHAMIDE-REACH [RQ] (cyclophosphamide 500 mg injection, 1 vial) Endoxan [BX] (cyclophosphamide 1 g injection, 1 vial) Endoxan [BX] (cyclophosphamide 2 g injection, 1 vial)

■ **IFOSFAMIDE**

Injection

4448D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	4000 mg	19	..	*241.58	31.60	Holoxan [BX] (ifosfamide 1 g injection, 1 vial) Holoxan [BX] (ifosfamide 2 g injection, 1 vial)

ANTIMETABOLITES

Folic acid analogues

■ **METHOTREXATE**

Injection

4502Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	250 mg	5	..	*114.12	31.60	DBL Methotrexate [PF] (methotrexate 1 g/10 mL injection, 10 mL vial) DBL Methotrexate [PF] (methotrexate 5 mg/2 mL injection, 5 x 2 mL vials) DBL Methotrexate [PF] (methotrexate 50 mg/2 mL injection, 5 x 2 mL vials) DBL Methotrexate [PF] (methotrexate 500 mg/20 mL injection, 20 mL vial) Methotrexate Accord [OD] (methotrexate 1 g/10 mL injection, 10 mL vial) Methotrexate Ebewe [SZ] (methotrexate 5 g/50 mL injection, 50 mL vial)

■ **METHOTREXATE**

Restricted benefit

Patients receiving treatment with a high dose regimen

Injection

4512L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	20000 mg	*842.02	31.60	DBL Methotrexate [PF] (methotrexate 1 g/10 mL injection, 10 mL vial) DBL Methotrexate [PF] (methotrexate 5 mg/2 mL injection, 5 x 2 mL vials) DBL Methotrexate [PF] (methotrexate 50 mg/2 mL injection, 5 x 2 mL vials) DBL Methotrexate [PF] (methotrexate 500 mg/20 mL injection, 20 mL vial) Methotrexate Accord [OD] (methotrexate 1 g/10 mL injection, 10 mL vial) Methotrexate Ebewe [SZ] (methotrexate 5 g/50 mL injection, 50 mL vial)

■ **PEMETREXED**

Injection

4600D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1100 mg	5	..	*147.01	31.60	Pemetrexed Accord [OD] (pemetrexed 1 g injection, 1 vial) Pemetrexed Accord [OD] (pemetrexed 100 mg injection, 1 vial) Pemetrexed Accord [OD] (pemetrexed 500 mg injection, 1 vial) Pemetrexed APOTEX [TX] (pemetrexed 500 mg injection, 1 vial) Pemetrexed Ever Pharma [IT] (pemetrexed 1 g/40 mL injection, 40 mL vial) Pemetrexed Ever Pharma [IT] (pemetrexed 100 mg/4 mL injection, 4 mL vial) Pemetrexed Ever Pharma [IT] (pemetrexed 500 mg/20 mL injection, 20 mL vial) Pemetrexed SUN [RA] (pemetrexed 1 g injection, 1 vial)

Pemetrexed SUN [RA] (pemetrexed 100 mg injection, 1 vial)
Pemetrexed SUN [RA] (pemetrexed 500 mg injection, 1 vial)

■ PRALATREXATE

Note No increase in the maximum number of repeats may be authorised.

Authority required

Relapsed or chemotherapy refractory Peripheral T-cell Lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be relapsed or chemotherapy refractory, **AND**
- Patient must not develop progressive disease whilst receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

Injection

11272G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	80 mg	11	..	*4230.62	31.60	Folotyn [MF] (pralatrexate 20 mg/mL injection, 1 mL vial)

■ PRALATREXATE

Note No increase in the maximum number of repeats may be authorised.

Authority required

Relapsed or chemotherapy refractory Peripheral T-cell Lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be relapsed or chemotherapy refractory, **AND**
- Patient must have undergone appropriate prior front-line curative intent chemotherapy.

Injection

11293J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	80 mg	5	..	*4230.62	31.60	Folotyn [MF] (pralatrexate 20 mg/mL injection, 1 mL vial)

■ RALTITREXED

Injection

4610P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	7 mg	8	..	*1131.46	31.60	Tomudex [PF] (raltitrexed 2 mg injection, 1 vial)

Purine analogues

■ CLADRIBINE

Authority required (STREAMLINED)

6265

Hairy cell leukaemia

Injection

4326Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	17 mg	6	..	*897.88	31.60	Leustatin [IX] (cladribine 10 mg/10 mL injection, 10 mL vial) Litak [AF] (cladribine 10 mg/5 mL injection, 5 mL vial)

■ FLUDARABINE

Note Pharmaceutical benefits that have the form fludarabine phosphate 50 mg injection and pharmaceutical benefits that have the form fludarabine phosphate 50 mg/2 mL injection are equivalent for the purposes of substitution.

Injection

4393F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	55 mg	29	..	*152.30	31.60	Fludarabine Ebewe [SZ] (fludarabine phosphate 50 mg/2 mL injection, 5 x 2 mL vials) Fludarabine Juno [JO] (fludarabine phosphate 50 mg injection, 1 vial)

Pyrimidine analogues

■ CYTARABINE

Injection

4357H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	7000 mg	15	..	*886.62	31.60	Pfizer Australia Pty Ltd [PF] (cytarabine 100 mg/5 mL injection, 5 x 5 mL vials)

■ FLUOROURACIL

Restricted benefit

Patients requiring administration of fluorouracil by intravenous infusion

Injection

4394G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	5500 mg	11	..	*111.46	31.60	Fluorouracil Accord [OC] (fluorouracil 1 g/20 mL injection, 20 mL vial) Fluorouracil Accord [OC] (fluorouracil 2.5 g/50 mL injection, 50 mL vial) Fluorouracil Accord [OC] (fluorouracil 5 g/100 mL injection, 100 mL vial) Fluorouracil Accord [OC] (fluorouracil 500 mg/10 mL injection, 10 mL vial) Fluorouracil Ebewe [SZ] (fluorouracil 1 g/20 mL injection, 20 mL vial) Fluorouracil Ebewe [SZ] (fluorouracil 5 g/100 mL injection, 100 mL vial)

▪ **FLUOROURACIL**

Restricted benefit

Patients requiring administration of fluorouracil by intravenous injection

Injection

4431F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	23	..	*94.46	31.60	Fluorouracil Accord [OC] (fluorouracil 1 g/20 mL injection, 20 mL vial) Fluorouracil Accord [OC] (fluorouracil 2.5 g/50 mL injection, 50 mL vial) Fluorouracil Accord [OC] (fluorouracil 5 g/100 mL injection, 100 mL vial) Fluorouracil Accord [OC] (fluorouracil 500 mg/10 mL injection, 10 mL vial) Fluorouracil Ebewe [SZ] (fluorouracil 1 g/20 mL injection, 20 mL vial) Fluorouracil Ebewe [SZ] (fluorouracil 5 g/100 mL injection, 100 mL vial)

▪ **GEMCITABINE**

Caution Pharmaceutical benefits containing gemcitabine may have different concentrations.

Injection

4439P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3000 mg	17	..	*151.67	31.60	DBL Gemcitabine Injection [PF] (gemcitabine 1 g/26.3 mL injection, 26.3 mL vial) DBL Gemcitabine Injection [PF] (gemcitabine 2 g/52.6 mL injection, 52.6 mL vial)

PLANT ALKALOIDS AND OTHER NATURAL PRODUCTS

Vinca alkaloids and analogues

▪ **VINBLASTINE**

Injection

4618C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	20 mg	17	..	*161.96	31.60	DBL Vinblastine [PF] (vinblastine sulfate 10 mg/10 mL injection, 5 x 10 mL vials)

▪ **VINCRIStINE**

Injection

4619D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	2 mg	7	..	*106.46	31.60	DBL Vincristine Sulfate [PF] (vincristine sulfate 1 mg/mL injection, 5 x 1 mL vials)

▪ **VINOReLBINE**

Injection

4620E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	70 mg	7	..	*150.52	31.60	Navelbine [FB] (vinorelbine 50 mg/5 mL injection, 5 mL vial) Vinorelbine Ebewe [SZ] (vinorelbine 10 mg/mL injection, 1 mL vial) Vinorelbine Ebewe [SZ] (vinorelbine 50 mg/5 mL injection, 5 mL vial)

Podophyllotoxin derivatives

▪ **ETOPOSIDE**

Injection

4428C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	440 mg	14	..	*282.12	31.60	Etopophos [LM] (etoposide phosphate 1.136 g (etoposide 1 g) injection, 1 vial) Etoposide Ebewe [SZ] (etoposide 100 mg/5 mL injection, 5 x 5 mL vials)

Taxanes

▪ **CABAZITAXEL**

Note Where the term 'novel hormonal drug' appears in this restriction, it refers to: (i) abiraterone, (ii) abiraterone and methylprednisolone, (iii) apalutamide, (iv) darolutamide, (v) enzalutamide.

Authority required (STREAMLINED)

13207

Castration resistant metastatic carcinoma of the prostate

Clinical criteria:

- The treatment must be in combination with prednisone or prednisolone, **AND**
- The condition must be resistant to treatment with docetaxel; **OR**
- Patient must have a documented intolerance necessitating permanent treatment withdrawal or a contraindication to docetaxel, **AND**
- The treatment must not be used in combination with a novel hormonal drug, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must not receive PBS-subsidised cabazitaxel if progressive disease develops while on cabazitaxel.

Injection

4376H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	55 mg	5	..	*177.29	31.60	Cabazitaxel Accord [OC] (cabazitaxel 60 mg/3 mL injection, 3 mL vial) Cabazitaxel Ever Pharma [IT] (cabazitaxel 60 mg/6 mL injection, 6 mL vial) Cabazitaxel Juno [JU] (cabazitaxel 60 mg/1.5 mL injection [1.5 mL vial] (& inert substance diluent [4.5 mL vial], 1 pack) MSN Cabazitaxel [RQ] (cabazitaxel 60 mg/1.5 mL injection [1.5 mL vial] (& inert substance diluent [4.5 mL vial], 1 pack)

▪ **DOCETAXEL**

Note Pharmaceutical benefits that have the forms docetaxel solution concentrate for I.V. infusion 80 mg in 4 mL and docetaxel solution concentrate for I.V. infusion 80 mg in 8 mL are equivalent for the purposes of substitution.

Note Pharmaceutical benefits that have the forms docetaxel solution concentrate for I.V. infusion 160 mg in 8 mL and docetaxel solution concentrate for I.V. infusion 160 mg in 16 mL are equivalent for the purposes of substitution.

Injection

10148D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	250 mg	5	..	*155.32	31.60	DBL Docetaxel Concentrated Injection [PF] (docetaxel 160 mg/16 mL injection, 16 mL vial) DBL Docetaxel Concentrated Injection [PF] (docetaxel 80 mg/8 mL injection, 8 mL vial) Docetaxel Accord [OC] (docetaxel 160 mg/8 mL injection, 8 mL vial) Docetaxel Accord [OC] (docetaxel 80 mg/4 mL injection, 4 mL vial)

▪ **NANOPARTICLE ALBUMIN-BOUND PACLITAXEL**

Authority required (STREAMLINED)

6106

Metastatic breast cancer

Authority required (STREAMLINED)

6119

HER2 positive breast cancer

Injection

4531L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	580 mg	5	..	*2148.24	31.60	Abraxane [TS] (paclitaxel (as nanoparticle albumin-bound) 100 mg injection, 1 vial)

▪ **NANOPARTICLE ALBUMIN-BOUND PACLITAXEL**

Note Special Pricing Arrangements apply.

Note Not for use as neoadjuvant or adjuvant therapy.

Authority required (STREAMLINED)

4657

Stage IV (metastatic) adenocarcinoma of the pancreas

Clinical criteria:

- The treatment must be in combination with gemcitabine, **AND**
- The condition must not have been treated previously with PBS-subsidised therapy, **AND**
- Patient must have an Eastern Cooperative Oncology Group (ECOG) performance status score of 2 or less.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Injection

10165B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	275 mg	11	..	*1118.43	31.60	Abraxane [TS] (paclitaxel (as nanoparticle albumin-bound) 100 mg injection, 1 vial)

▪ **PACLITAXEL**

Injection

4567J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	450 mg	3	..	*162.18	31.60	Paclitaxel Accord [OC] (paclitaxel 300 mg/50 mL injection, 50 mL vial) Paclitaxel Ebewe [SZ] (paclitaxel 300 mg/50 mL injection, 50 mL vial)

Topoisomerase 1 (TOP1) inhibitors

▪ **IRINOTECAN**

Note In first-line usage, effectiveness and tolerance may be improved when irinotecan is combined with an infusional 5-fluorouracil regimen.

Injection

4451G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	800 mg	11	..	*158.62	31.60	Irinotecan Accord [OC] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial) Irinotecan Accord [OC] (irinotecan hydrochloride trihydrate 500 mg/25 mL injection, 25 mL vial) Irinotecan Alphapharm [AF] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial) Irinotecan Alphapharm [AF] (irinotecan hydrochloride trihydrate 500 mg/25 mL injection, 25 mL vial) IRINOTECAN BAXTER [BX] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial) Omegapharm Irinotecan [OE] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial) Omegapharm Irinotecan [OE] (irinotecan hydrochloride trihydrate 40 mg/2 mL injection, 2 mL vial)

▪ **TOPOTECAN**

Injection

4617B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3500 mcg	17	..	*120.70	31.60	Hycamtin [SZ] (topotecan 4 mg injection, 5 vials) Topotecan Accord [OC] (topotecan 4 mg/4 mL injection, 5 x 4 mL vials)

Other plant alkaloids and natural products

▪ **TRABECTEDIN**

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14196

Advanced (unresectable and/or metastatic) leiomyosarcoma or liposarcoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have an ECOG performance status of 2 or less, **AND**
- Patient must have received prior chemotherapy treatment including an anthracycline, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The condition must be one of the following subtypes for patients with liposarcoma: (i) dedifferentiated, (ii) myxoid, (iii) round-cell, (iv) pleomorphic.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Injection

13346K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3250 mcg	3	..	*3325.50	31.60	Yondelis [ZL] (trabectedin 1 mg injection, 1 vial)

▪ TRABECTEDIN

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14197

Advanced (unresectable and/or metastatic) leiomyosarcoma or liposarcoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Authority required (STREAMLINED)

14188

Advanced (unresectable and/or metastatic) leiomyosarcoma or liposarcoma

Treatment Phase: Transitioning from non-PBS to PBS-subsidised treatment - Grandfather arrangements

Clinical criteria:

- Patient must have been receiving treatment with this drug for this condition prior to 1 August 2023, **AND**
- Patient must have had a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 2 at the time non-PBS supply was initiated, **AND**
- Patient must have received chemotherapy treatment including an anthracycline, prior to initiating non-PBS-subsidised treatment, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The condition must be one of the following subtypes for patients with liposarcoma: (i) dedifferentiated, (ii) myxoid, (iii) round-cell, (iv) pleomorphic.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Injection

13340D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3250 mcg	7	..	*3325.50	31.60	Yondelis [ZL] (trabectedin 1 mg injection, 1 vial)

CYTOTOXIC ANTIBIOTICS AND RELATED SUBSTANCES

Anthracyclines and related substances

▪ DOXORUBICIN

Injection/intravesical

4361M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	135 mg	11	..	*125.34	31.60	Adriamycin [PF] (doxorubicin hydrochloride 200 mg/100 mL injection, 100 mL vial) Adriamycin [PF] (doxorubicin hydrochloride 50 mg/25 mL injection, 25 mL vial) Doxorubicin ACC [OC] (doxorubicin hydrochloride 200 mg/100 mL injection, 100 mL vial)

▪ DOXORUBICIN HYDROCHLORIDE (AS PEGYLATED LIPOSOMAL)

Injection

4364Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	100 mg	5	..	*747.76	31.60	Caelyx [BX] (doxorubicin hydrochloride (as pegylated liposomal) 20 mg/10 mL injection, 10 mL vial) Caelyx [BX] (doxorubicin hydrochloride (as pegylated liposomal) 50 mg/25 mL injection, 25 mL vial) Liposomal Doxorubicin SUN [RA] (doxorubicin hydrochloride (as pegylated liposomal) 20 mg/10 mL injection, 10 mL vial) Liposomal Doxorubicin SUN [RA] (doxorubicin hydrochloride (as pegylated liposomal) 50 mg/25 mL injection, 25 mL vial)

▪ EPIRUBICIN

Injection/intravesical

4375G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	220 mg	5	..	*266.52	31.60	Epirubicin Accord [OC] (epirubicin hydrochloride 200 mg/100 mL injection, 100 mL vial)

▪ IDARUBICIN

Restricted benefit

Acute myelogenous leukaemia (AML)

Injection

4440Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	30 mg	5	..	*268.08	31.60	Zavedos Solution [PF] (idarubicin hydrochloride 5 mg/5 mL injection, 5 mL vial)

▪ **MITOZANTRONE**

Injection

4514N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	30 mg	5	..	*181.22	31.60	Mitozantrone Ebewe [SZ] (mitozantrone 20 mg/10 mL injection, 10 mL vial) Onkotrone [BX] (mitozantrone 25 mg/12.5 mL injection, 12.5 mL vial)

Other cytotoxic antibiotics

▪ **BLEOMYCIN**

Restricted benefit

Germ cell neoplasms

Restricted benefit

Lymphoma

Injection

4433H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	30000 iu	11	..	*168.62	31.60	DBL Bleomycin Sulfate [PF] (bleomycin sulfate 15 000 international units injection, 1 vial)

MONOCLONAL ANTIBODIES AND ANTIBODY DRUG CONJUGATES

CD20 (Clusters of Differentiation 20) inhibitors

▪ **OBINUTUZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Follicular lymphoma

Treatment Phase: Re-induction treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised obinutuzumab, **AND**
- The condition must be CD20 positive, **AND**
- The condition must be refractory to treatment with rituximab for this condition, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for re-induction treatment purposes only, **AND**
- The treatment must be in combination with bendamustine, **AND**
- The treatment must not exceed 8 doses for re-induction treatment with this drug for this condition.

The condition is considered rituximab-refractory if the patient experiences less than a partial response or progression of disease within 6 months after completion of a prior rituximab-containing regimen.

A patient may only qualify for PBS-subsidised initiation treatment once in a lifetime under:

- i) the previously untreated induction treatment restriction; or
- ii) the rituximab-refractory re-induction restriction.

Injection

11457B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	7	..	*4614.14	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ **OBINUTUZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Stage II bulky or Stage III/IV follicular lymphoma

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be CD20 positive, **AND**
- The condition must be previously untreated, **AND**
- The condition must be symptomatic, **AND**
- The treatment must be for induction treatment purposes only, **AND**
- The treatment must be in combination with chemotherapy, **AND**

- The treatment must not exceed 10 doses for induction treatment with this drug for this condition. A patient may only qualify for PBS-subsidised initiation treatment once in a lifetime under:
 - i) the previously untreated induction treatment restriction; or
 - ii) the rituximab-refractory re-induction restriction.

Injection

11458C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	9	..	*4614.14	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ OBINUTUZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Stage II bulky or Stage III/IV follicular lymphoma

Treatment Phase: Maintenance therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug under the previously untreated initial restriction, **AND**
- The condition must be CD20 positive, **AND**
- Patient must have demonstrated a partial or complete response to PBS subsidised induction treatment with this drug for this condition, **AND**
- The treatment must be maintenance therapy, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed 12 doses or 2 years duration of treatment, whichever comes first, under this restriction, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Injection

11462G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	5	..	*4614.14	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ OBINUTUZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Follicular lymphoma

Treatment Phase: Maintenance therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug under the rituximab refractory initial restriction, **AND**
- The condition must be CD20 positive, **AND**
- The condition must have been refractory to treatment with rituximab, **AND**
- Patient must have demonstrated a partial or complete response to PBS-subsidised re-induction treatment with this drug for this condition, **AND**
- The treatment must be maintenance therapy, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed 12 doses or 2 years duration of treatment, whichever comes first, under this restriction, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Injection

11468N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	5	..	*4614.14	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ OBINUTUZUMAB

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

11015

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: For combination use with venetoclax treatment cycles 1 to 6 inclusive in first-line therapy

Clinical criteria:

- The condition must be untreated, **AND**
- The treatment must be in combination with PBS-subsidised venetoclax.

Injection

12204H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	8	..	*4614.14	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ **OBINUTUZUMAB**

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

14764

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: For combination use with acalabrutinib from treatment cycles 2 to 7 inclusive in first-line therapy

Clinical criteria:

- The condition must be untreated, **AND**
- The treatment must be in combination with PBS-subsidised acalabrutinib (refer to Product Information for timing of obinutuzumab and acalabrutinib doses).

Injection

13787P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	7	..	*4614.14	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ **OBINUTUZUMAB**

Note The latest International Workshop on CLL (iwCLL) provides guidance on various aspects of management of CLL/SLL. Notably, two of these are:

- (1) when to treat versus when to monitor the patient without therapy - see 'Indications for treatment' section; and
- (2) recognising progressive disease - see 'Definition of response, relapse, and refractory disease' section.

See the following literature reference for details:

Hallek, M et al. iwCLL guidelines for diagnosis, indications for treatment, response assessment, and supportive management of CLL. **Blood** vol. 131, 25 (2018): 2745-2760.

Note Obinutuzumab is not to be used as monotherapy or in combination with anti-cancer drugs other than chlorambucil under this restriction. For use with venetoclax, refer to the separate listing for this purpose.

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

14326

Chronic lymphocytic leukaemia (CLL)

Treatment Phase: Combination use with chlorambucil only

Clinical criteria:

- The condition must be CD20 positive, **AND**
 - The condition must be previously untreated, **AND**
 - The treatment must be in combination with chlorambucil, **AND**
 - The treatment must only be prescribed for a patient with active disease in accordance with the International Workshop on CLL (iwCLL) guidance (latest version) in relation to when to prescribe drug treatment for this condition.
- Treatment must be discontinued in patients who experience disease progression whilst on this treatment.

Injection

10407R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	7	..	*4614.14	31.60	Gazyva [RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial)

▪ **RITUXIMAB**

Note Prescribing/pharmacy claiming: prescribe/claim this benefit through the Section 100 Highly Specialised Drugs Program PBS item code(s) when administered for non-oncology indications. Prescribe/claim this benefit through the Efficient Funding of Chemotherapy PBS item code(s) when administered for oncology indications.

Injection

13102N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	800 mg	11	..	*434.52	31.60	Riximyo [SZ] (rituximab 100 mg/10 mL injection, 2 x 10 mL vials) Riximyo [SZ] (rituximab 500 mg/50 mL injection, 50 mL vial) Ruxience [PF] (rituximab 100 mg/10 mL injection, 10 mL vial) Ruxience [PF] (rituximab 500 mg/50 mL injection, 50 mL vial) Truxima [EW] (rituximab 100 mg/10 mL injection, 2 x 10 mL vials) Truxima [EW] (rituximab 500 mg/50 mL injection, 50 mL vial)

CD22 (Clusters of Differentiation 22) inhibitors

■ INOTUZUMAB OZOGAMICIN

Caution Careful monitoring of patients is required due to risk of developing hepatotoxicity, including life-threatening hepatic veno-occlusive disease, and the increased risk of post-haematopoietic stem cell transplant non-relapse mortality observed in patients treated with inotuzumab.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note A complete remission is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a full recovery of peripheral blood counts with platelets of greater than 100,000 per microliter, and absolute neutrophil count (ANC) of greater than 1,000 per microliter.

Note A complete remission with partial haematological recovery is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a partial recovery of peripheral blood counts with platelets of greater than 50,000 per microliter, and absolute neutrophil count (ANC) of greater than 500 per microliter.

Note Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time when an assessment is required must cease PBS-subsidised therapy with this agent.

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Consolidation treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised induction treatment with this drug for this condition, **AND**
 - Patient must have achieved a complete remission; OR
 - Patient must have achieved a complete remission with partial haematological recovery, **AND**
 - The treatment must not be more than 5 treatment cycles under this restriction in a lifetime, **AND**
 - Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug. This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.
- The treatment must not exceed 0.5mg per m² for all doses within a treatment cycle
Treatment with this drug for this condition must not exceed 6 treatment cycles in a lifetime.

Injection

11680R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	2820 mcg	4	..	*39416.34	31.60	Besponsa [PF] (inotuzumab ozogamicin 1 mg injection, 1 vial)

■ INOTUZUMAB OZOGAMICIN

Caution Careful monitoring of patients is required due to risk of developing hepatotoxicity, including life-threatening hepatic veno-occlusive disease, and the increased risk of post-haematopoietic stem cell transplant non-relapse mortality observed in patients treated with inotuzumab.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patients are eligible to receive a loading dose for the first dose of a treatment cycle while receiving induction treatment. Two prescriptions are required, the first prescription for the loading dose at a dose no higher than 0.8mg per m², and the second prescription for two doses at a dose no higher than 0.5mg per m². Both prescriptions must be submitted with the initial application.

Note Once a patient achieves complete remission or complete remission with partial haematological recovery, a new prescription must be written under the consolidation treatment phase.

Note A complete remission is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a full recovery of peripheral blood counts with platelets of greater than 100,000 per microliter, and absolute neutrophil count (ANC) of greater than 1,000 per microliter.

Note A complete remission with partial haematological recovery is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a partial recovery of peripheral blood counts with platelets of greater than 50,000 per microliter, and absolute neutrophil count (ANC) of greater than 500 per microliter.

Note Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time when an assessment is required must cease PBS-subsidised therapy with this agent.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos.
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be relapsed or refractory B-precursor cell ALL, with an Eastern Cooperative Oncology Group (ECOG) performance status of 2 or less, **AND**
- Patient must have received intensive combination chemotherapy for initial treatment of ALL or for subsequent salvage therapy, **AND**
- Patient must not have received more than 1 line of salvage therapy, **AND**
- Patient must have previously received a tyrosine kinase inhibitor (TKI) if the condition is Philadelphia chromosome positive, **AND**
- The condition must be CD22-positive, **AND**
- The condition must have more than 5% blasts in bone marrow, **AND**
- The treatment must not be more than 3 treatment cycles under this restriction in a lifetime.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

The authority application must be made in writing and must include:

- (1) two completed authority prescription forms;
- (2) a completed Acute Lymphoblastic Leukaemia PBS Authority Application - Supporting Information Form; and
- (3) evidence that the condition is CD22-positive; and
- (4) date of most recent chemotherapy, and if this was the initial chemotherapy regimen or salvage therapy, including what line of salvage; and
- (5) a copy of the most recent bone marrow biopsy report of no more than one month old at the time of application.

The treatment must not exceed 0.8mg per m² for the first dose of a treatment cycle (Day 1), and 0.5mg per m² for subsequent doses (Days 8 and 15) within a treatment cycle.

Treatment with this drug for this condition must not exceed 6 treatment cycles in a lifetime.

Injection

11696N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3384 mcg	2	..	*52525.58	31.60	Besponsa [PF] (inotuzumab ozogamicin 1 mg injection, 1 vial)

CD38 (Clusters of Differentiation 38) inhibitors

▪ DARATUMUMAB

Note This drug is not PBS-subsidised for use in patients with multiple myeloma who have received two or more prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent, or, who are refractory to both a PI and an immunomodulatory agent, as monotherapy.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment of second-line drug therapy for weeks 10 to 24 (administered every 3 weeks)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with bortezomib and dexamethasone, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition. Progressive disease is defined as at least 1 of the following:
 - (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
 - (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
 - (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
 - (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
 - (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
 - (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
 - (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

12220E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1920 mg	4	..	*11772.42	31.60	Darzalex [JC] (daratumumab 100 mg/5 mL injection, 5 mL vial) Darzalex [JC] (daratumumab 400 mg/20 mL injection, 20 mL vial)

▪ DARATUMUMAB

Note This drug is not PBS-subsidised for use in patients with multiple myeloma who have received two or more prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent, or, who are refractory to both a PI and an immunomodulatory agent, as monotherapy.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Initial treatment as second-line drug therapy for weeks 1 to 9 (administered once weekly)

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with bortezomib and dexamethasone, **AND**
- Patient must have progressive disease after only one prior therapy (i.e. use must be as second-line drug therapy; use as third-line drug therapy or beyond is not PBS-subsidised).

Treatment criteria:

- Patient must be undergoing treatment with this drug in one of the following situations: (i) for the first time, irrespective of whether the diagnosis has been reclassified (i.e. the diagnosis has changed between multiple myeloma/amyloidosis), (ii) changing the drug's form (intravenous/subcutaneous) within the first 9 weeks of treatment for the same PBS indication.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Details of: the histological diagnosis of multiple myeloma; prior treatments including name(s) of drug(s) and date of most recent treatment cycle; the basis of the diagnosis of progressive disease or failure to respond; and which disease activity parameters will be used to assess response, must be documented in the patient's medical records.

Confirmation of eligibility for treatment with current diagnostic reports of at least one of the following must be documented in the patient's medical records:

- (a) the level of serum monoclonal protein; or
- (b) Bence-Jones proteinuria - the results of 24-hour urinary light chain M protein excretion; or
- (c) the serum level of free kappa and lambda light chains; or
- (d) bone marrow aspirate or trephine; or
- (e) if present, the size and location of lytic bone lesions (not including compression fractures); or
- (f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
- (g) if present, the level of hypercalcaemia, corrected for albumin concentration.

As these parameters must be used to determine response, results for either (a) or (b) or (c) should be documented for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) must be documented in the patient's medical records. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (current serum M protein less than 10 g per L) must be documented in the patient's medical records.

A line of therapy is defined as 1 or more cycles of a planned treatment program. This may consist of 1 or more planned cycles of single-agent therapy or combination therapy, as well as a sequence of treatments administered in a planned manner.

A new line of therapy starts when a planned course of therapy is modified to include other treatment agents (alone or in combination) as a result of disease progression, relapse, or toxicity, with the exception to this being the need to attain a sufficient response for stem cell transplantation to proceed. A new line of therapy also starts when a planned period of observation off therapy is interrupted by a need for additional treatment for the disease.

Injection

12228N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1920 mg	8	..	*11772.42	31.60	Darzalex [JC] (daratumumab 100 mg/5 mL injection, 5 mL vial) Darzalex [JC] (daratumumab 400 mg/20 mL injection, 20 mL vial)

▪ **DARATUMUMAB**

Note This drug is not PBS-subsidised for use in patients with multiple myeloma who have received two or more prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent, or, who are refractory to both a PI and an immunomodulatory agent, as monotherapy.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment of second-line drug therapy from week 25 until disease progression (administered every 4 weeks)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition. Progressive disease is defined as at least 1 of the following:
 - (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
 - (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
 - (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
 - (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
 - (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
 - (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
 - (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

12231R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1920 mg	5	..	*11772.42	31.60	Darzalex [JC] (daratumumab 100 mg/5 mL injection, 5 mL vial)
						Darzalex [JC] (daratumumab 400 mg/20 mL injection, 20 mL vial)

HER2 (Human Epidermal Growth Factor Receptor 2) inhibitors

▪ **PERTUZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) either in the primary tumour or a metastatic lesion, confirmed through a pathology report from an Approved Pathology Authority, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must not have received prior anti-HER2 therapy for this condition, **AND**
- Patient must not have received prior chemotherapy for this condition, **AND**
- The treatment must be in combination with trastuzumab and a taxane, **AND**
- The treatment must not be in combination with nab-paclitaxel, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Details (date, unique identifying number/code, or provider number) of the pathology report from an Approved Pathology Authority confirming evidence of HER2 gene amplification in the primary tumour or a metastatic lesion by in situ hybridisation (ISH) must be provided at the time of application.

The pathology report must be documented in the patient's medical records.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to seeking the initial authority approval.

Injection

10267J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	840 mg	*5926.12	31.60	Perjeta [RO] (pertuzumab 420 mg/14 mL injection, 14 mL vial)

■ PERTUZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note The criterion that limits breaks in treatment with pertuzumab under this restriction has been temporarily modified due to the current risk of COVID-19. This allows an extended break in therapy with PBS-subsidised pertuzumab in patients who are at risk of COVID-19.

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug, **AND**
- The treatment must be in combination with trastuzumab, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

The treatment must not exceed a lifetime total of one course. However, treatment breaks are permitted. A patient who has a treatment break in PBS-subsidised treatment with this drug for reasons other than disease progression is eligible to continue to receive PBS-subsidised treatment with this drug.

Where a patient has had a treatment break the length of the break is measured from the date the most recent treatment was stopped to the date of the application for further treatment.

Injection

10333W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	420 mg	3	..	*3007.37	31.60	Perjeta [RO] (pertuzumab 420 mg/14 mL injection, 14 mL vial)

■ TRASTUZUMAB

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

10296

Early HER2 positive breast cancer

Treatment Phase: Initial treatment (weekly regimen)

Clinical criteria:

- Patient must have undergone surgery (adjuvant) or be preparing for surgery (neoadjuvant), **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
- Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.

HER2 positivity must be demonstrated by in situ hybridisation (ISH).

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

Injection

4632T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	500 mg	*467.00	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

■ TRASTUZUMAB

Note Increased maximum quantity will be authorised where a patient requires a new loading dose due to a break in therapy of more than 1 week but less than 6 weeks from the last dose or a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

10213

Early HER2 positive breast cancer

Treatment Phase: Continuing treatment (weekly regimen)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
- Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.

Injection

4639E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	250 mg	9	..	*328.55	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (&) inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB**

Note Increased maximum quantity will be authorised where a patient requires a new loading dose due to a break in therapy of more than 1 week but less than 6 weeks from the last dose or a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

10294

Early HER2 positive breast cancer

Treatment Phase: Continuing treatment (3 weekly regimen)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
- Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.

Injection

4703M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	750 mg	3	..	*706.93	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (&) inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum quantity will be authorised where a patient's weight is greater than 125 kg.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required (STREAMLINED)

9353

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) either in the primary tumour or a metastatic lesion, **AND**
- The treatment must not be in combination with nab-paclitaxel, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

Injection

10391X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	*845.38	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial)

Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack)
 Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial)
 Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial)
 Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial)
 Trazimera [PF] (trastuzumab 150 mg injection, 1 vial)
 Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

■ TRASTUZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum quantity will be authorised where a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required (STREAMLINED)

9349

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Where a patient has a break in trastuzumab therapy of more than 1 week from when the last dose was due, a new loading dose may be required.

Injection

10401K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	750 mg	3	..	*706.93	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

■ TRASTUZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum quantity will be authorised where a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9573

Metastatic (Stage IV) HER2 positive adenocarcinoma of the stomach or gastro-oesophageal junction

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) positivity as demonstrated by immunohistochemistry 2+ or more in tumour material, **AND**
- Patient must have evidence of HER2 gene amplification as demonstrated by in situ hybridisation results based on more than 6 copies of HER2 in the same tumour tissue sample, **AND**
- Patient must have evidence of HER2 gene amplification as demonstrated by in situ hybridisation results based on the ratio of HER2 to chromosome 17 being more than 2 in the same tumour tissue sample, **AND**
- Patient must commence treatment in combination with platinum based chemotherapy and capecitabine; OR
- Patient must commence treatment in combination with platinum based chemotherapy and 5 fluorouracil, **AND**
- Patient must not have previously received this drug for this condition, **AND**
- Patient must not have received prior chemotherapy for this condition, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

Injection

10581X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	*845.38	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial)

Trazimera [PF] (trastuzumab 150 mg injection, 1 vial)
Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum quantity will be authorised where a patient requires a new loading dose due to a break in therapy of more than 1 week but less than 6 weeks from the last dose or a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9571

Metastatic (Stage IV) HER2 positive adenocarcinoma of the stomach or gastro-oesophageal junction

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Injection

10588G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	750 mg	3	..	*706.93	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

10293

Early HER2 positive breast cancer

Treatment Phase: Initial treatment (3 weekly regimen)

Clinical criteria:

- Patient must have undergone surgery (adjuvant) or be preparing for surgery (neoadjuvant), **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
- Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.

HER2 positivity must be demonstrated by in situ hybridisation (ISH).

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

Injection

4650R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	*845.38	31.60	Herzuma [EW] (trastuzumab 150 mg injection, 1 vial) Herzuma [EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack) Kanjinti [JU] (trastuzumab 150 mg injection, 1 vial) Kanjinti [JU] (trastuzumab 420 mg injection, 1 vial) Ogivri [SZ] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 150 mg injection, 1 vial) Trazimera [PF] (trastuzumab 60 mg injection, 1 vial)

▪ **TRASTUZUMAB DERUXTECAN**

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Clinical criteria:

- Patient must have evidence of human epidermal growth factor (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) in either the primary tumour/a metastatic lesion - establish this finding once only with the first PBS prescription, **AND**
- The condition must have progressed following treatment with at least one prior HER2 directed regimen for metastatic breast cancer; OR
- The condition must have, at the time of treatment initiation with this drug, progressed during/within 6 months following adjuvant treatment with a HER2 directed therapy, **AND**
- Patient must have, at the time of initiating treatment with this drug, a WHO performance status no higher than 1, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this PBS indication, **AND**
- The treatment must not be prescribed where any of the following is present: (i) left ventricular ejection fraction of less than 50%, (ii) symptomatic heart failure; confirm cardiac function testing for the first PBS prescription only.

Treatment criteria:

- Patient must be undergoing initial treatment with this drug - the following are true: (i) this is the first prescription for this drug, (ii) this prescription seeks no more than 3 repeat prescriptions; OR
- Patient must be undergoing continuing treatment with drug - the following are true: (i) there has been an absence of further disease progression whilst on active treatment with this drug, (ii) this prescription does not seek to re-treat after disease progression, (iii) this prescription seeks no more than 8 repeat prescriptions.

Confirm that the following information is documented/retained in the patient's medical records once only with the first PBS prescription:

- 1) Evidence of HER2 gene amplification (evidence obtained in relation to past PBS treatment is acceptable).
- 2) Details of prior HER2 directed drug regimens prescribed for the patient.
- 3) Cardiac function test results (evidence obtained in relation to past PBS treatment is acceptable).

Injection

13718B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	675 mg	8	..	*17540.04	31.60	Enhertu [AP] (trastuzumab deruxtecan 100 mg injection, 1 vial)

▪ **TRASTUZUMAB EMTANSINE**

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPoS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) either in the primary tumour or a metastatic lesion, confirmed through a pathology report from an Approved Pathology Authority, **AND**
- The condition must have progressed following treatment with pertuzumab and trastuzumab in combination; OR
- The condition must have progressed during or within 6 months of completing adjuvant therapy with trastuzumab, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

The following information must be provided by the prescriber at the time of application:

- (a) details (date, unique identifying number/code or provider number) of the pathology report from an Approved Pathology Authority confirming evidence of HER2 gene amplification in the primary tumour or a metastatic lesion by in situ hybridisation (ISH).
- (b) dates of treatment with trastuzumab and pertuzumab;
- (c) date of demonstration of progression following treatment with trastuzumab and pertuzumab; or
- (d) date of demonstration of progression and date of completion of adjuvant trastuzumab treatment.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application.

All reports must be documented in the patient's medical records.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to seeking the initial authority approval.

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for metastatic (Stage IV) HER2 positive breast cancer, **AND**

- Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

The treatment must not exceed a lifetime total of one continuous course for this PBS indication.

Injection

10282E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	450 mg	8	..	*7268.61	31.60	Kadcyla [RO] (trastuzumab emtansine 100 mg injection, 1 vial)
						Kadcyla [RO] (trastuzumab emtansine 160 mg injection, 1 vial)

▪ **TRASTUZUMAB EMTANSINE**

Note No increase in the maximum number of repeats may be authorised.

Note Increased maximum amounts can be requested where a patient's weight is greater than 125 kg.

Authority required

Early HER2 positive breast cancer

Treatment Phase: Initial adjuvant treatment

Clinical criteria:

- The treatment must be prescribed within 12 weeks after surgery, **AND**
- Patient must have, prior to commencing treatment with this drug, evidence of residual invasive cancer in the breast and/or axillary lymph nodes following completion of surgery, as demonstrated by a pathology report, **AND**
- Patient must have completed systemic neoadjuvant therapy that included trastuzumab and taxane-based chemotherapy prior to surgery, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- The treatment must not extend beyond 42 weeks (14 cycles) duration under the initial and the continuing treatment restrictions combined.

Authority applications for initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail and must include:

(a) details (date, unique identifying number/code or provider number) of the pathology report from an Approved Pathology Authority demonstrating evidence of residual invasive carcinoma in the breast and/or axillary lymph nodes following completion of surgery.

The pathology report must be documented in the patient's medical records.

If the application is submitted through HPOS form upload or mail, it must include:

- (i) A completed authority prescription form; and
- (ii) A completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Early HER2 positive breast cancer

Treatment Phase: Continuing adjuvant treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- The treatment must not extend beyond 42 weeks (14 cycles) duration under the initial and the continuing treatment restrictions combined.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Injection

11951B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	450 mg	6	..	*7268.61	31.60	Kadcyla [RO] (trastuzumab emtansine 100 mg injection, 1 vial) Kadcyla [RO] (trastuzumab emtansine 160 mg injection, 1 vial)

EGFR (Epidermal Growth Factor Receptor) inhibitors

■ CETUXIMAB

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12470

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with PBS-subsidised encorafenib for this condition.

Injection

12816M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	550 mg	11	..	*1699.77	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

■ CETUXIMAB

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12483

Metastatic colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with PBS-subsidised encorafenib for this condition.

Injection

12820R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	880 mg	*2505.36	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

■ CETUXIMAB

Note A maximum lifetime supply for this indication is limited to a maximum of 8 treatments per site and to 10 treatments per site for patients in whom radiotherapy is interrupted.

Authority required (STREAMLINED)

4788

Stage III, IVa or IVb squamous cell cancer of the larynx, oropharynx or hypopharynx

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with radiotherapy, **AND**
- Patient must be unable to tolerate cisplatin; OR
- Patient must have a contraindication to cisplatin according to the TGA-approved Product Information.

Injection

4435K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	550 mg	5	..	*1699.77	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

■ CETUXIMAB

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

4794

Stage III, IVa or IVb squamous cell cancer of the larynx, oropharynx or hypopharynx

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be for the week prior to radiotherapy, **AND**
- Patient must have a contraindication to cisplatin according to the TGA-approved Product Information.

Authority required (STREAMLINED)

4785

Stage III, IVa or IVb squamous cell cancer of the larynx, oropharynx or hypopharynx

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with radiotherapy, **AND**
- Patient must be unable to tolerate cisplatin.

Injection

4312Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	880 mg	*2505.36	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

▪ **CETUXIMAB**

Note Special Pricing Arrangements apply.

Note This drug is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Authority required (STREAMLINED)

12045

Metastatic colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have RAS wild-type metastatic colorectal cancer, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The condition must have failed to respond to first-line chemotherapy; OR
- The condition must have progressed following first-line treatment with pembrolizumab for dMMR mCRC, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on panitumumab are not eligible to receive PBS-subsidised cetuximab.

Patients who have developed intolerance to panitumumab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised cetuximab.

Authority required (STREAMLINED)

4908

Metastatic colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have RAS wild-type metastatic colorectal cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must be previously untreated, **AND**
- The treatment must be in combination with first-line chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Injection

4436L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	880 mg	*2505.36	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbitux [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

▪ **CETUXIMAB**

Note Special Pricing Arrangements apply.

Note This drug is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Note This drug is not PBS-subsidised when chemotherapy partners are switched whilst maintaining an anti-EGFR antibody backbone in the face of progressive disease.

Note The treatment must not exceed a single course of therapy with this drug for metastatic colorectal cancer in a patient's lifetime.

Authority required (STREAMLINED)

4912

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for first-line treatment of RAS wild-type metastatic colorectal cancer, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must be in combination with first-line chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Injection

10262D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	550 mg	18	..	*1699.77	31.60	Erbitux [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial)

▪ **CETUXIMAB**

Note Special Pricing Arrangements apply.

Note This drug is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Note This drug is not PBS-subsidised when chemotherapy partners are switched whilst maintaining an anti-EGFR antibody backbone in the face of progressive disease.

Note The treatment must not exceed a single course of therapy with this drug for metastatic colorectal cancer in a patient's lifetime.

Authority required (STREAMLINED)

12016

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for treatment of RAS wild-type metastatic colorectal cancer after failure of first-line chemotherapy; OR
- Patient must have received an initial authority prescription for this drug for treatment of RAS wild-type metastatic colorectal cancer after failure of treatment with first-line pembrolizumab for dMMR mCRC, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on panitumumab are not eligible to receive PBS-subsidised cetuximab.

Patients who have developed intolerance to panitumumab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised cetuximab.

Injection

4731B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	550 mg	11	..	*1699.77	31.60	Erbix [SG] (cetuximab 100 mg/20 mL injection, 20 mL vial) Erbix [SG] (cetuximab 500 mg/100 mL injection, 100 mL vial)

▪ **PANITUMUMAB**

Note This drug is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Authority required (STREAMLINED)

12066

Metastatic colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have RAS wild-type metastatic colorectal cancer, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The condition must have failed to respond to first-line chemotherapy; OR
- The condition must have progressed following first-line treatment with pembrolizumab for dMMR mCRC, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on cetuximab are not eligible to receive PBS-subsidised panitumumab.

Patients who have developed intolerance to cetuximab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised panitumumab.

Authority required (STREAMLINED)

12035

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for treatment of RAS wild-type metastatic colorectal cancer after failure of first-line chemotherapy; OR
- Patient must have received an initial authority prescription for this drug for treatment of RAS wild-type metastatic colorectal cancer after failure of treatment with first-line pembrolizumab for dMMR mCRC, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on cetuximab are not eligible to receive PBS-subsidised panitumumab.

Patients who have developed intolerance to cetuximab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised panitumumab.

Note This drug is not PBS-subsidised when chemotherapy partners are switched whilst maintaining an anti-EGFR antibody backbone in the face of progressive disease.

Note The treatment must not exceed a single course of therapy with this drug for metastatic colorectal cancer in a patient's lifetime.

Injection

10082P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	720 mg	5	..	*3691.96	31.60	Vectibix [AN] (panitumumab 100 mg/5 mL injection, 5 mL vial) Vectibix [AN] (panitumumab 400 mg/20 mL injection, 20 mL vial)

▪ **PANITUMUMAB**

Note Special Pricing Arrangements apply.

Note Panitumumab is not PBS-subsidised for use in combination with another anti-EGFR antibody or in combination with an anti-VEGF antibody.

Authority required (STREAMLINED)

5526

Metastatic colorectal cancer

Treatment Phase: Initial Treatment

Clinical criteria:

- Patient must have RAS wild-type metastatic colorectal cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must be previously untreated, **AND**
- The treatment must be in combination with first-line chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on cetuximab are not eligible to receive PBS-subsidised panitumumab.

Patients who have developed intolerance to cetuximab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised panitumumab.

Authority required (STREAMLINED)

5452

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for panitumumab for first-line treatment of RAS wild-type metastatic colorectal cancer, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must be in combination with first-line chemotherapy, **AND**
- The treatment must be the sole PBS-subsidised anti-EGFR antibody therapy for this condition.

Patients who have progressive disease on cetuximab are not eligible to receive PBS-subsidised panitumumab.

Patients who have developed intolerance to cetuximab of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised panitumumab.

Note This drug is not PBS-subsidised when chemotherapy partners are switched whilst maintaining an anti-EGFR antibody backbone in the face of progressive disease.

Note The treatment must not exceed a single course of therapy with this drug for metastatic colorectal cancer in a patient's lifetime.

Injection

10513H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	720 mg	9	..	*3691.96	31.60	Vectibix [AN] (panitumumab 100 mg/5 mL injection, 5 mL vial) Vectibix [AN] (panitumumab 400 mg/20 mL injection, 20 mL vial)

PD-1/PDL-1 (Programmed cell death protein 1/death ligand 1) inhibitors

▪ **ATEZOLIZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10297

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- Patient must have stable or responding disease.

Injection

11277M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	7	..	*6835.99	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

■ ATEZOLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10216

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing first-line treatment of metastatic disease - 3 weekly treatment regimen

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab until disease progression, unless not tolerated.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug in this line of treatment, **AND**
- Patient must have stable or responding disease.

Injection

11802E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	7	..	*6835.99	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

■ ATEZOLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10215

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Continuing treatment - 4 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have stable or responding disease.

Injection

11930X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	5	..	*9534.92	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

■ ATEZOLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10257

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing first-line treatment of metastatic disease, as monotherapy, where concomitant bevacizumab has ceased due to intolerance - 4 weekly treatment regimen

Clinical criteria:

- Patient must have experienced intolerance to combination treatment with bevacizumab, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug in this line of treatment, **AND**
- Patient must have stable or responding disease, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

12097Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	5	..	*9534.92	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

■ ATEZOLIZUMAB

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13443

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The condition must have progressed on or after prior platinum based chemotherapy; OR
- The condition must have progressed after treatment with tepotinib.

Injection

11284X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	5	..	*6835.99	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ **ATEZOLIZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10206

Extensive-stage small cell lung cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be previously untreated, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be in combination with etoposide and a platinum-based antineoplastic drug.

Injection

11926Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	3	..	*6835.99	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ **ATEZOLIZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10521

Extensive-stage small cell lung cancer

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Injection

11929W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	4	..	*6835.99	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ **ATEZOLIZUMAB**

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13446

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Initial treatment - 4 weekly treatment regimen

Clinical criteria:

- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The condition must have progressed on or after prior platinum based chemotherapy; OR
- The condition must have progressed after treatment with tepotinib.

Injection

11931Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	3	..	*9534.92	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

▪ ATEZOLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10509

Extensive-stage small cell lung cancer

Treatment Phase: Continuing treatment - 4 weekly treatment regimen

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Injection

12078Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	3	..	*9534.92	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

▪ ATEZOLIZUMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10917

Advanced (unresectable) Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Continuing treatment of hepatocellular carcinoma - 3 weekly treatment regimen

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab until disease progression, unless not tolerated.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - Patient must not have developed disease progression while being treated with this drug for this condition.
- PBS supply of this drug must be through only one of the two continuing treatment regimens at any given time

Injection

12168K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	8	..	*6835.99	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ ATEZOLIZUMAB

Note No increase in the maximum amount or number of units may be authorised.

Note Increased repeats of up to 11 may be requested for doses of 840 mg administered every 2 weeks

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10972

Advanced (unresectable) Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Continuing treatment where bevacizumab is discontinued - 4 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - Patient must not have developed disease progression while being treated with this drug for this condition.
- PBS supply of this drug must be through only one of the two continuing treatment regimens at any given time

Injection

12174R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	5	..	*9534.92	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

▪ ATEZOLIZUMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13451

Resected early stage (Stage II to IIIA) non-small cell lung cancer (NSCLC)

Treatment Phase: 1,680 mg administered once every 4 weeks, or 840 mg every 2 weeks

Population criteria:

- Patient must be both: (i) initiating treatment, (ii) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy; OR
- Patient must be continuing existing PBS-subsidised treatment with this drug; OR
- Patient must be both: (i) transitioning from existing non-PBS to PBS subsidised supply of this drug, (ii) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy at the time this drug was initiated.

Clinical criteria:

- Patient must have/have had a WHO performance status score of no greater than 1 at treatment initiation with this drug. **AND**
- The treatment must be for the purpose of adjuvant therapy following all of: (i) surgical resection, (ii) platinum-based chemotherapy, **AND**
- The condition must have/have had, at treatment commencement, an absence of each of the following gene abnormalities confirmed via tumour material sampling: (i) an activating epidermal growth factor receptor (EGFR) gene mutation, (ii) an anaplastic lymphoma kinase (ALK) gene rearrangement, **AND**
- The condition must have/have had, at treatment commencement, confirmation of programmed cell death ligand 1 (PD-L1) expression on at least 50% of tumour cells, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment that does not occur beyond the following, whichever comes first: (i) the first instance of disease progression/recurrence, (ii) 12 months in total for this condition from the first administered dose; mark any remaining repeat prescriptions with the words 'cancelled' where (i)/(ii) has occurred.

Injection

13173H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1680 mg	5	..	*9534.92	31.60	Tecentriq [RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial)

▪ **ATEZOLIZUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13442

Resected early stage (Stage II to IIIA) non-small cell lung cancer (NSCLC)

Treatment Phase: 1,200 mg administered once every 3 weeks

Population criteria:

- Patient must be both: (i) initiating treatment, (ii) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy; OR
- Patient must be continuing existing PBS-subsidised treatment with this drug; OR
- Patient must be both: (i) transitioning from existing non-PBS to PBS subsidised supply of this drug, (ii) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy at the time this drug was initiated.

Clinical criteria:

- Patient must have/have had a WHO performance status score of no greater than 1 at treatment initiation with this drug. **AND**
- The treatment must be for the purpose of adjuvant therapy following all of: (i) surgical resection, (ii) platinum-based chemotherapy, **AND**
- The condition must have/have had, at treatment commencement, an absence of each of the following gene abnormalities confirmed via tumour material sampling: (i) an activating epidermal growth factor receptor (EGFR) gene mutation, (ii) an anaplastic lymphoma kinase (ALK) gene rearrangement, **AND**
- The condition must have/have had, at treatment commencement, confirmation of programmed cell death ligand 1 (PD-L1) expression on at least 50% of tumour cells, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment that does not occur beyond the following, whichever comes first: (i) the first instance of disease progression/recurrence, (ii) 12 months in total for this condition from the first administered dose; mark any remaining repeat prescriptions with the words 'cancelled' where (i)/(ii) has occurred.

Injection

13174J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	7	..	*6835.99	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ **ATEZOLIZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13448

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment 1

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab and platinum-doublet chemotherapy.

Clinical criteria:

- The condition must be non-squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene mutation or an anaplastic lymphoma kinase (ALK) gene rearrangement in tumour material.

Authority required (STREAMLINED)**10125**

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment 2

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab and platinum-doublet chemotherapy.

Clinical criteria:

- The condition must be non-squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must have evidence of an activating epidermal growth factor receptor (EGFR) gene mutation or of an anaplastic lymphoma kinase (ALK) gene rearrangement in tumour material, **AND**
- Patient must have progressive disease following treatment with an epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI) OR an anaplastic lymphoma kinase (ALK) tyrosine kinase inhibitor (TKI), **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer.

Injection

11807K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	5	..	*6835.99	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ ATEZOLIZUMAB

Caution The safety of atezolizumab in combination with bevacizumab has not been established in patients who have incompletely treated varices, variceal bleeding within the previous 6 months or who are at high risk of bleeding. Patients should be assessed for risk of variceal bleeding prior to treatment with this combination.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10939**

Advanced (unresectable) Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Initial treatment

Treatment criteria:

- Patient must be undergoing combination treatment with bevacizumab and atezolizumab until disease progression, unless not tolerated.

Clinical criteria:

- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must not be suitable for transarterial chemoembolisation, **AND**
- Patient must have Child Pugh class A, **AND**
- The condition must be untreated with systemic therapy; OR
- Patient must have developed intolerance to a vascular endothelial growth factor (VEGF) tyrosine kinase inhibitor (TKI) of a severity necessitating permanent treatment withdrawal.

Injection

12171N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	3	..	*6835.99	31.60	Tecentriq [RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)

▪ AVELUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10023

Stage IV (metastatic) Merkel Cell Carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 10 mg per kg every 2 weeks under this restriction.

Injection

11671G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	11	..	*8232.78	31.60	Bavencio [SG] (avelumab 200 mg/10 mL injection, 10 mL vial)

▪ **AVELUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

8947

Stage IV (metastatic) Merkel Cell Carcinoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed a total of 9 doses at a maximum dose of 10 mg per kg every 2 weeks under this restriction.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

11695M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	8	..	*8232.78	31.60	Bavencio [SG] (avelumab 200 mg/10 mL injection, 10 mL vial)

▪ **AVELUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13290

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Treatment Phase: Maintenance therapy - Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

13126W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	800 mg	11	..	*5518.06	31.60	Bavencio [SG] (avelumab 200 mg/10 mL injection, 10 mL vial)

▪ **AVELUMAB**

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13313

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Treatment Phase: Maintenance therapy - Initial treatment

Clinical criteria:

- Patient must have received first-line platinum-based chemotherapy, **AND**
- Patient must not have progressive disease following first-line platinum-based chemotherapy, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

13122P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	800 mg	7	..	*5518.06	31.60	Bavencio [SG] (avelumab 200 mg/10 mL injection, 10 mL vial)

■ CEMIPILIMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

15063

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a total of 35 cycles or up to 24 months of treatment under both initial and continuing treatment restrictions, whichever comes first.

Injection

13161Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	350 mg	6	..	*7378.62	31.60	Libtayo [WM] (cemiplimab 350 mg/7 mL injection, 7 mL vial)

■ CEMIPILIMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

15094

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must not exceed a total of 7 doses under this restriction.

Injection

13169D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	350 mg	6	..	*7378.62	31.60	Libtayo [WM] (cemiplimab 350 mg/7 mL injection, 7 mL vial)

■ CEMIPILIMAB

Caution In the first few months after starting immunotherapy, a transient tumour flare may occur that may be mistaken as disease progression despite an overall positive response to treatment.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Metastatic or locally advanced cutaneous squamous cell carcinoma (CSCC)

Treatment Phase: Initial treatment covering the first 3 treatment cycles

Clinical criteria:

- The condition must be unsuitable for each of: (i) curative surgical resection, (ii) curative radiotherapy, **AND**
- Patient must have had a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

13152F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	350 mg	2	..	*7378.62	31.60	Libtayo [WM] (cemiplimab 350 mg/7 mL injection, 7 mL vial)

▪ **CEMIPLIMAB**

Caution In the first few months after starting immunotherapy, a transient tumour flare may occur that may be mistaken as disease progression despite an overall positive response to treatment.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Metastatic or locally advanced cutaneous squamous cell carcinoma (CSCC)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Patient must not be undergoing treatment with this drug as a PBS benefit where the treatment duration extends beyond the following, whichever comes first: (i) disease progression despite treatment with this drug, (ii) 24 months from treatment initiation; annotate any remaining repeat prescriptions with the word 'cancelled' where this occurs.

Injection

13159N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	350 mg	7	..	*7378.62	31.60	Libtayo [WM] (cemiplimab 350 mg/7 mL injection, 7 mL vial)

▪ **DOSTARLIMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

15163

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Initial treatment covering the first 6 treatment cycles

Clinical criteria:

- Patient must have deficient mismatch repair (dMMR) endometrial cancer, as determined by immunohistochemistry test, **AND**
- The condition must be unsuitable for at least one of the following: (i) curative surgical resection, (ii) curative radiotherapy, **AND**
- The treatment must be initiated in combination with platinum-containing chemotherapy, **AND**
- The condition must be, at treatment initiation with this drug, either: (i) untreated with systemic therapy, (ii) treated with neoadjuvant/adjuvant systemic therapy, but the cancer has recurred or progressed after more than 6 months from the last dose of systemic therapy, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation.

Injection

14122G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	500 mg	5	..	*7736.12	31.60	Jemperli [GK] (dostarlimab 500 mg/10 mL injection, 10 mL vial)

▪ **DOSTARLIMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

15205

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 36 cumulative months from the first administered dose, once in a lifetime.

Authority required (STREAMLINED)

15196

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Transitioning from non-PBS to PBS-subsidised treatment - Grandfather treatment

Clinical criteria:

- Patient must have deficient mismatch repair (dMMR) endometrial cancer, as determined by immunohistochemistry test, **AND**
- Patient must have received non-PBS-subsidised treatment with this drug for this condition prior to 1 May 2024, **AND**
- The condition must be, prior to initiation of non-PBS-subsidised treatment with this drug, unsuitable for at least one of the following: (i) curative surgical resection, (ii) curative radiotherapy, **AND**
- The condition must be, prior to initiation of non-PBS-subsidised treatment with this drug, either: (i) untreated with systemic therapy, (ii) treated with neoadjuvant/adjuvant systemic therapy, but the cancer has recurred or progressed after more than 6 months from the last dose of systemic therapy, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation, **AND**
- The treatment must be, at initiation of non-PBS-subsidised treatment with this drug, used in combination with platinum-containing chemotherapy, **AND**
- Patient must not have developed disease progression while receiving non-PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 36 cumulative months from the first administered dose, once in a lifetime.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Injection

14130Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1000 mg	3	..	*15383.62	31.60	Jemperli [GK] (dostarlimab 500 mg/10 mL injection, 10 mL vial)

■ DURVALUMAB

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10126**

Unresectable Stage III non-small cell lung cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have received platinum based chemoradiation therapy, **AND**
- The condition must not have progressed following platinum based chemoradiation therapy, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

Authority required (STREAMLINED)**12271**

Unresectable Stage III non-small cell lung cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The treatment must not exceed 12 months in total for this condition under the initial and continuing restriction combined, **AND**
- The treatment must be once in a lifetime with this drug for this condition.

Injection

11915D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1500 mg	4	..	*11417.37	31.60	Imfinzi [AP] (durvalumab 120 mg/2.4 mL injection, 2.4 mL vial) Imfinzi [AP] (durvalumab 500 mg/10 mL injection, 10 mL vial)

■ DURVALUMAB

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**14708**

Locally advanced, metastatic or recurrent biliary tract cancer (intrahepatic cholangiocarcinoma, extrahepatic cholangiocarcinoma, and gallbladder cancer)

Population criteria:

- Patient must have either of the following at treatment initiation: (i) locally advanced biliary tract cancer that is untreated with systemic anti-cancer therapy in the unresectable setting, (ii) metastatic biliary tract cancer that is untreated with systemic anti-cancer therapy in the metastatic setting.

Clinical criteria:

- Patient must have/have had a WHO performance status score of no greater than 1 at treatment initiation with this drug. **AND**
- The treatment must be/have been initiated with both: (i) gemcitabine, (ii) cisplatin (refer to Product Information of gemcitabine and cisplatin for dosing information), **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Injection

13767N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1500 mg	5	..	*11417.37	31.60	Imfinzi [AP] (durvalumab 120 mg/2.4 mL injection, 2.4 mL vial)
						Imfinzi [AP] (durvalumab 500 mg/10 mL injection, 10 mL vial)

▪ **DURVALUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10206

Extensive-stage small cell lung cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be previously untreated, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be in combination with etoposide and a platinum-based antineoplastic drug.

Injection

13775B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1500 mg	3	..	*11417.37	31.60	Imfinzi [AP] (durvalumab 120 mg/2.4 mL injection, 2.4 mL vial)
						Imfinzi [AP] (durvalumab 500 mg/10 mL injection, 10 mL vial)

▪ **DURVALUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10509

Extensive-stage small cell lung cancer

Treatment Phase: Continuing treatment - 4 weekly treatment regimen

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Injection

13780G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1500 mg	5	..	*11417.37	31.60	Imfinzi [AP] (durvalumab 120 mg/2.4 mL injection, 2.4 mL vial)
						Imfinzi [AP] (durvalumab 500 mg/10 mL injection, 10 mL vial)

▪ **NIVOLUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

11477

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Continuing treatment as second-line drug therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- Patient must have stable or responding disease. Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11153B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

9299

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11160J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note Special Pricing Arrangements apply.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9252

Recurrent or metastatic squamous cell carcinoma of the oral cavity, pharynx or larynx

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have stable or responding disease, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11411N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

9321

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Maintenance treatment

Clinical criteria:

- Patient must have previously received of up to maximum 4 doses of PBS-subsidised combined therapy with nivolumab and ipilimumab as induction for this condition, **AND**
- The treatment must be as monotherapy for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

11642R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

9298

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Authority required (STREAMLINED)

13839

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Maintenance treatment

Clinical criteria:

- Patient must have previously received of up to maximum 4 doses of PBS-subsidised combined therapy with nivolumab and ipilimumab as induction for this condition, **AND**
- The treatment must be as monotherapy for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this PBS indication.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

10745M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	11	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ **NIVOLUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

14816

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
- Patient must not have experienced disease progression whilst on adjuvant PD-1 inhibitor treatment or disease recurrence within 6 months of completion of adjuvant PD-1 inhibitor treatment if treated for resected Stage IIIB, IIIC, IIID or IV melanoma, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

10764M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	8	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ **NIVOLUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13445

Locally advanced or metastatic non-small cell lung cancer

Treatment Phase: Initial treatment as second-line drug therapy

Clinical criteria:

- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**

- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
- The condition must have progressed on or after prior platinum based chemotherapy; OR
- The condition must have progressed after treatment with tepotinib.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11158G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	8	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

9216

Recurrent or metastatic squamous cell carcinoma of the oral cavity, pharynx or larynx

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The condition must have progressed within 6 months of the last dose of prior platinum based chemotherapy, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor for this condition.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated. Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11435W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	8	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14830

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Induction treatment

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
 - Patient must have an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1, **AND**
 - The condition must not be ocular or uveal melanoma, **AND**
 - The treatment must be in combination with PBS-subsidised treatment with ipilimumab as induction for this condition.
- Induction treatment with nivolumab must not exceed a total of 4 doses at a maximum dose of 1 mg per kg every 3 weeks. Induction treatment with ipilimumab must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks.

Injection

11543M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	3	..	*2456.13	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.
Stable disease (SD) is small changes that do not meet above criteria.

Authority required (STREAMLINED)

9312

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial Treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must have progressive disease according to the Response Evaluation Criteria in Solid Tumours (RECIST) following prior treatment with a tyrosine kinase inhibitor; OR
- Patient must have developed intolerance to a tyrosine kinase inhibitor of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11150W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	8	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13433

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial combination treatment (with ipilimumab) as first-line drug therapy

Clinical criteria:

- The condition must be squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must be in combination with platinum-based chemotherapy for the first two cycles, **AND**
- The treatment must be in combination with ipilimumab.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

11468

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing combination treatment (with ipilimumab) of first-line drug therapy

Clinical criteria:

- The condition must be squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed 24 months in total, measured from the initial dose, or, must not extend beyond disease progression, whichever comes first, **AND**
- The treatment must be in combination with ipilimumab.

Injection

12323N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	360 mg	13	..	*7191.12	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note An increase of number of repeats may be authorised up to 11 if the patient is receiving a weight based dosing of 3mg/kg every 2 weeks.

Authority required (STREAMLINED)

11985

Unresectable malignant mesothelioma

Clinical criteria:

- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be in combination with PBS-subsidised ipilimumab, unless an intolerance to ipilimumab of a severity necessitating permanent treatment withdrawal of ipilimumab, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a maximum total of 24 months in a lifetime for this condition.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

12602G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	360 mg	8	..	*7191.12	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Caution In the first few months after starting immunotherapy, a transient tumour flare may occur that may be mistaken as disease progression despite an overall positive response to treatment.

Note The stated maximum amount in this listing is based on this drug's approved Product Information recommended dosing in specific cancer types - the drug may be prescribed in a quantity up to this amount, but need not be this amount for every cancer type. Refer to this drug's approved Product Information (Dose and Method of Administration or Clinical Trials sections) for the recommended dosing in the specific cancer type.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14676

Advanced or metastatic gastro-oesophageal cancers

Clinical criteria:

- Patient must have/have had, at the time of initiating treatment with this drug, a WHO performance status no higher than 1, **AND**
- Patient must be untreated (up until initiating this drug) with programmed cell death-1/ligand-1 (PD-1/PD-L1) inhibitor therapy for gastro-oesophageal cancer.

Treatment criteria:

- Patient must not be undergoing treatment with this drug as a PBS benefit where the treatment duration extends beyond the following, whichever comes first: (i) disease progression despite treatment with this drug, (ii) 24 months from treatment initiation; annotate any remaining repeat prescriptions with the word 'cancelled' where this occurs.

Population criteria:

- Patient must be in one of the three population subsets described below.

Population 1

Conditions: gastric cancer, gastro-oesophageal junction cancer, oesophageal adenocarcinoma

Concomitant therapies: chemotherapy containing at least a fluoropyrimidine drug plus a platinum drug

Line of treatment: first-line drug treatment

Additional clinical finding: HER2 negative

Population 2

Condition: oesophageal squamous cell carcinoma (can be recurrent)

Concomitant therapies: chemotherapy containing at least a fluoropyrimidine drug plus a platinum drug

Line of treatment: first-line drug treatment

Additional clinical finding: unresectable

Population 3

Condition: oesophageal squamous cell carcinoma (can be recurrent)

Line of treatment: second-line drug treatment after chemotherapy containing at least a fluoropyrimidine drug plus a platinum drug

Additional clinical finding: unresectable

Injection

13121N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	13	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Up to an additional 2 repeat prescriptions (7 in total) may be sought only where dosing is on a 2-weekly schedule in the first 16 weeks of treatment. This listing's stated number of repeat prescriptions is based on 4-weekly dosing.

Note No increase in the maximum amount or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Adjuvant treatment of stage II or III oesophageal cancer or gastro-oesophageal junction cancer

Clinical criteria:

- The condition must have histological evidence confirming a diagnosis of a least one of: (i) adenocarcinoma, (ii) squamous cell cancer; document this evidence in the patient's medical records, **AND**
- The condition must have been treated with neoadjuvant platinum-based chemoradiotherapy, **AND**
- The treatment must be for the purposes of adjuvant use following complete surgical resection that occurred within 16 weeks prior to initiating this drug, **AND**
- The condition must have evidence, through resected specimen, that residual disease meets the Tumour Nodes Metastases (TNM) staging system (as published by the Union for International Cancer Control) of either: (i) at least ypT1, (ii) at least ypN1; document this evidence in the patient's medical records, **AND**
- Patient must have/have had, at the time of initiating treatment with this drug, a WHO performance status no higher than 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment with a dosing regimen as set out in the drug's approved Australian Product Information, **AND**
- Patient must not be undergoing PBS-subsidised treatment with this drug where this prescription extends treatment beyond whichever comes first: (i) 12 months from treatment initiation, irrespective of whether initial treatment was PBS-subsidised/non-PBS-subsidised, (ii) disease recurrence despite treatment with this drug; annotate any remaining repeat prescriptions with the word 'cancelled' where this occurs.

Injection

13240W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	5	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

▪ **NIVOLUMAB**

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note A prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk score can be calculated here: <https://www.mdcalc.com/imdc-international-metastatic-ccc-database-consortium-risk-model-metastatic-renal-cell-carcinoma>.

One point is assigned for each of:

- (i) a time of diagnosis to systemic therapy of less than 1 year
- (ii) a Karnofsky Performance Status of less than 80%
- (iii) a haemoglobin less than the lower limit of normal
- (iv) a corrected calcium level greater than the upper limit of normal
- (v) a neutrophil count greater than the upper limit of normal
- (vi) a platelet count greater than the upper limit of normal

Stated normal reference ranges may vary depending on the laboratory providing the measurement. 'Normal' here refers to the individual laboratory's stated normal reference range.

Favourable IMDC risk is a score of 0.

Intermediate IMDC risk is a score of 1 to 2.

Poor IMDC risk is a score of 3 to 6.

Document any IMDC risk score assessment in the patient's medical records.

Authority required (STREAMLINED)

14001

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must not have previously been treated, **AND**
- Patient must have a prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk classification score at treatment initiation with this drug of either: (i) 1 to 2 (intermediate risk), (ii) 3 to 6 (poor risk); document the IMDC risk classification score in the patient's medical records, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must be in combination with PBS-subsidised treatment with ipilimumab as induction for this condition. Induction treatment with nivolumab must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks. The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

11636K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	360 mg	3	..	*7191.12	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ NIVOLUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Resected Stage IIIB, IIIC, IIID or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be adjuvant to complete surgical resection, **AND**
- Patient must have a WHO performance status of 1 or less, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- The treatment must commence within 12 weeks of complete resection, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy. Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required

Resected Stage IIIB, IIIC, IIID or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for adjuvant treatment following complete surgical resection, **AND**
- Patient must not have experienced disease recurrence, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy. Patients must only receive a maximum of 240 mg every two weeks or 480 mg every four weeks under a weight based or flat dosing regimen.

Injection

11900H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	480 mg	5	..	*9558.60	31.60	Opdivo [BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial) Opdivo [BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

10705

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease.

Injection

10436G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	7	..	*7736.12	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

10701

Unresectable Stage III or Stage IV malignant melanoma
Treatment Phase: Continuing treatment - 6 weekly treatment regimen

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease.

Injection

12124D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	3	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14727

Stage II or Stage III triple negative breast cancer

Clinical criteria:

- The treatment must be initiated in combination with neoadjuvant chemotherapy, **AND**
- The condition must not have progressed/recurred whilst on treatment with this drug.

Treatment criteria:

- Patient must not be undergoing treatment with this drug beyond 52 cumulative weeks under this restriction, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 7 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 4 repeat prescriptions.

Injection

13752T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	7	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

14818

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
- Patient must not have experienced disease progression whilst on adjuvant PD-1 inhibitor treatment or disease recurrence within 6 months of completion of adjuvant PD-1 inhibitor treatment if treated for resected Stage IIIB, IIIC, IIID or IV melanoma, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed a total of 6 doses under this restriction.

Injection

10493G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	5	..	*7736.12	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

14817

Unresectable Stage III or Stage IV malignant melanoma
Treatment Phase: Initial treatment - 6 weekly treatment regimen

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
- Patient must not have experienced disease progression whilst on adjuvant PD-1 inhibitor treatment or disease recurrence within 6 months of completion of adjuvant PD-1 inhibitor treatment if treated for resected Stage IIIB, IIIC, IIID or IV melanoma, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed a total of 3 doses under this restriction.

Injection

12128H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	2	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note A Combined Positive Score (CPS) is determined by:

The number of PD-L1-stained cells (tumour cells, lymphocytes, macrophages) divided by the number of all viable tumour cells (i.e. the total number of: PD-L1-positive tumour cells plus PD-L1-negative tumour cells).

Although the result of the CPS calculation can exceed 100, the maximum score is defined as CPS 100.

A minimum of 100 viable tumour cells in the PD-L1-stained slide is required for the specimen to be considered adequate for PD-L1 evaluation.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**14324**

Recurrent, unresectable or metastatic triple negative breast cancer

Clinical criteria:

- The condition must have been (up until this drug therapy) untreated in the unresectable/metastatic disease stage, **AND**
- The condition must have been (up until this drug therapy) untreated with programmed cell death-1/ligand 1 (PD-1/PD-L1) inhibitor therapy in breast cancer, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation, **AND**
- The treatment must be in combination with chemotherapy, **AND**
- The condition must have both: (i) programmed cell death ligand 1 (PD-L1) expression confirmed by a validated test, (ii) a Combined Positive Score (CPS) of at least 10 at treatment initiation.

Treatment criteria:

- Patient must be undergoing initial treatment with this drug - this is the first prescription for this drug; OR
- Patient must be undergoing continuing treatment with this drug - both the following are true: (i) the condition has not progressed on active treatment with this drug, (ii) this prescription does not extend PBS subsidy beyond 24 cumulative months from the first administered dose, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Injection

13608F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)**13726**

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have undergone an autologous stem cell transplant (ASCT) for this condition and have experienced relapsed or refractory disease post ASCT; OR
- Patient must not be suitable for ASCT for this condition and have experienced relapsed or refractory disease following at least 2 prior treatments for this condition, **AND**

- Patient must not have received prior treatment with a PD-1 (programmed cell death-1) inhibitor for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required (STREAMLINED)

13741

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

11330H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13727

Relapsed or refractory primary mediastinal B-cell lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as primary mediastinal B-cell lymphoma through histological investigation combined with at least one of: (i) positron emission tomography - computed tomography (PET-CT) scan, (ii) PET scan, (iii) CT scan, **AND**

- Patient must have been treated with rituximab-based chemotherapy for this condition, **AND**
- Patient must be experiencing relapsed/refractory disease, **AND**
- Patient must be autologous stem cell transplant (ASCT) ineligible following a single line of treatment; OR
- Patient must have undergone an autologous stem cell transplant (ASCT); OR
- Patient must have been treated with at least 2 chemotherapy treatment lines for this condition, one of which must include rituximab-based chemotherapy, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required (STREAMLINED)

13732

Relapsed or refractory primary mediastinal B-cell lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR

- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

12129J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13431

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must not exceed a total of 7 doses under this restriction.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13432

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a total of 35 cycles or up to 24 months of treatment under both initial and continuing treatment restrictions, whichever comes first.

Injection

11494Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	6	..	*7736.12	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13739

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The condition must have progressed on or after prior platinum based chemotherapy; OR
- The condition must have progressed on or within 12 months of completion of adjuvant platinum-containing chemotherapy following cystectomy for localised muscle-invasive urothelial cancer; OR
- The condition must have progressed on or within 12 months of completion of neoadjuvant platinum-containing chemotherapy prior to cystectomy for localised muscle-invasive urothelial cancer, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13736

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

11646Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13436

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment - 6 weekly treatment regimen

Clinical criteria:

- Patient must not have previously been treated for this condition in the metastatic setting; OR
- The condition must have progressed after treatment with tepotinib, **AND**
- Patient must not have received prior treatment with a programmed cell death-1 (PD-1) inhibitor or a programmed cell death ligand-1 (PD-L1) inhibitor for non-small cell lung cancer, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must not exceed a total of 4 doses under this restriction.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13437

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment - 6 weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a total of 18 cycles or up to 24 months of treatment under both initial and continuing treatment restrictions, whichever comes first.

Injection

12119W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	3	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Initial treatment - 6 weekly treatment regimen

Clinical criteria:

- The treatment must be adjuvant to complete surgical resection, **AND**
- Patient must have a WHO performance status of 1 or less, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- The treatment must commence within 12 weeks of complete resection, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Continuing treatment - 6 weekly treatment regimen

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for adjuvant treatment following complete surgical resection, **AND**
- Patient must not have experienced disease recurrence, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

Injection

12127G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	3	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

■ PEMBROLIZUMAB

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Unresectable or metastatic deficient mismatch repair (dMMR) colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be untreated for this PBS indication (i.e untreated for each of: (i) unresectable disease, (ii) metastatic disease), **AND**
- Patient must not have received prior treatment for colorectal cancer with each of: (i) a programmed cell death-1 (PD-1) inhibitor, (ii) a programmed cell death ligand-1 (PD-L1) inhibitor, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must have deficient mismatch repair (dMMR) colorectal cancer, as determined by immunohistochemistry test.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required

Unresectable or metastatic deficient mismatch repair (dMMR) colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR

- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

12615Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Authority required (STREAMLINED)

13735

Recurrent or metastatic squamous cell carcinoma of the oral cavity, pharynx or larynx

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be incurable by local therapies in the locally advanced setting, **AND**
- Patient must not have had systemic therapy for this condition in the recurrent or metastatic setting prior to initiating PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have experienced disease recurrence within 6 months of completion of systemic therapy if previously treated in the locally advanced setting, **AND**
- Patient must have had a WHO performance status of 0 or 1, **AND**
- The treatment must be either: (i) the sole PBS-subsidised therapy where the condition expresses programmed cell death ligand 1 (PD-L1) with a combined positive score (CPS) greater than or equal to 20 in the tumour sample, (ii) in combination with platinum-based chemotherapy, unless contraindicated or not tolerated.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

13731

Recurrent or metastatic squamous cell carcinoma of the oral cavity, pharynx or larynx

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

13131D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

13948

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk classification score at treatment initiation with this drug of either: (i) 1 to 2 (intermediate risk), (ii) 3 to 6 (poor risk); document the IMDC risk classification score in the patient's medical records, **AND**
- The condition must be untreated, **AND**
- Patient must have a WHO performance status of 2 or less.

Treatment criteria:

- Patient must be undergoing combination therapy consisting of: (i) pembrolizumab, (ii) lenvatinib; OR
- Patient must be undergoing monotherapy with this drug due to a contraindication/intolerance to the other drug in the combination mentioned above, requiring temporary/permanent discontinuation; document the details in the patient's medical records, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note A prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk score can be calculated here: <https://www.mdcalc.com/imdc-international-metastatic-renal-cell-carcinoma>.

One point is assigned for each of:

- (i) a time of diagnosis to systemic therapy of less than 1 year
- (ii) a Karnofsky Performance Status of less than 80%
- (iii) a haemoglobin less than the lower limit of normal
- (iv) a corrected calcium level greater than the upper limit of normal
- (v) a neutrophil count greater than the upper limit of normal
- (vi) a platelet count greater than the upper limit of normal

Stated normal reference ranges may vary depending on the laboratory providing the measurement. 'Normal' here refers to the individual laboratory's stated normal reference range.

Favourable IMDC risk is a score of 0.

Intermediate IMDC risk is a score of 1 to 2.

Poor IMDC risk is a score of 3 to 6.

Document any IMDC risk score assessment in the patient's medical records.

Authority required (STREAMLINED)

13949

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing combination therapy consisting of: (i) pembrolizumab, (ii) lenvatinib; OR
- Patient must be undergoing monotherapy with this drug due to a contraindication/intolerance to the other drug in the combination mentioned above, requiring temporary/permanent discontinuation; document the details in the patient's medical records, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Injection

13254N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ PEMBROLIZUMAB

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Initial treatment - 3 weekly treatment regimen

Clinical criteria:

- The treatment must be in addition to complete surgical resection, **AND**
- Patient must have a WHO performance status of 1 or less, **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- The treatment must commence within 12 weeks of complete resection, **AND**
- Patient must not have received more than 12 months of therapy (irrespective of whether therapy has been partly PBS-subsidised/non-PBS-subsidised).

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note Commencing treatment within 12 weeks of complete resection means either 12 weeks after resection or 12 weeks prior to resection.

Where non-PBS-subsidised supply has occurred, the total amount of PBS-subsidised supply is intended to be the balance of 18 doses less the number of non-PBS-subsidised doses.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Continuing treatment - 3 weekly treatment regimen

Treatment criteria:

- Patient must be undergoing continuing PBS-subsidised treatment commenced through an 'Initial treatment' listing.

Clinical criteria:

- Patient must not have experienced disease recurrence, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received more than 12 months of therapy (irrespective of whether therapy has been partly PBS-subsidised/non-PBS-subsidised).

Injection

12130K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	7	..	*7736.12	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ **PEMBROLIZUMAB**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14027

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have received prior treatment with platinum-based chemotherapy, **AND**
- The condition must be untreated with each of: (i) programmed cell death-1/ligand-1 (PD-1/PDL-1) inhibitor therapy, (ii) tyrosine kinase inhibitor therapy, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation.

Treatment criteria:

- Patient must be undergoing combination therapy consisting of: (i) pembrolizumab, (ii) lenvatinib; OR
- Patient must be undergoing monotherapy with this drug due to a contraindication/intolerance to the other drug in the combination mentioned above, requiring temporary/permanent discontinuation; document the details in the patient's medical records, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

14044

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing combination therapy consisting of: (i) pembrolizumab, (ii) lenvatinib; OR
- Patient must be undergoing monotherapy with this drug due to a contraindication/intolerance to the other drug in the combination mentioned above, requiring temporary/permanent discontinuation; document the details in the patient's medical records, **AND**

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions, **AND**
- Patient must not be undergoing continuing PBS-subsidised treatment where this benefit is extending treatment beyond 24 cumulative months from the first administered dose, once in a lifetime.

Authority required (STREAMLINED)

14028

Advanced, metastatic or recurrent endometrial carcinoma

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply - Grandfather arrangements

Clinical criteria:

- Patient must have received non-PBS-subsidised treatment with this drug for this condition prior to 1 June 2023, **AND**
- The treatment must be occurring in a patient where each of the following is true: (i) the patient had received prior treatment with platinum-based chemotherapy, (ii) the patient was untreated at treatment initiation with each of: (a) programmed cell death-1/ligand-1 (PD-1/PDL-1) inhibitor therapy, (b) tyrosine kinase inhibitor therapy, (iii) the patient's WHO performance status was no higher than 1 at treatment initiation, (iv) this drug is being prescribed in either: (a) a combination of pembrolizumab plus lenvatinib only, (b) as monotherapy where there was a contraindication/intolerance to the other drug in the combination - document the details in the patient's medical records, (v) disease progression has not occurred whilst on treatment, (vi) this prescription does not extend treatment beyond 24 months from the first administered dose.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Injection

13286G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

▪ PEMBROLIZUMAB

Note Patient should be treated with the recommended dose of pembrolizumab according to the TGA-approved Product Information.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14403

Advanced carcinoma of the cervix

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be at least one of (i) persistent carcinoma, (ii) recurrent carcinoma, (iii) metastatic carcinoma of the cervix, **AND**
- The condition must be unsuitable for curative treatment with either of (i) surgical resection, (ii) radiation, **AND**
- Patient must have WHO performance status no higher than 1, **AND**
- Patient must not have received prior treatment for this PBS indication.

Treatment criteria:

- Patient must be undergoing concomitant treatment with chemotherapy, containing a minimum of: (i) a platinum-based chemotherapy agent, plus (ii) paclitaxel, **AND**
- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required (STREAMLINED)

14404

Advanced carcinoma of the cervix

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a total of (i) 24 months, (ii) 35 doses (based on a 3-weekly dose regimen), (iii) 17 doses (based on a 6-weekly dose regimen) whichever comes first from the first dose of this drug regardless if it was PBS/non-PBS subsidised.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Authority required (STREAMLINED)

14405

Advanced carcinoma of the cervix

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply - Grandfather arrangements

Clinical criteria:

- Patient must be currently receiving non-PBS-subsidised treatment with this drug for this condition, with treatment having commenced prior to 1 October 2023, **AND**
- Patient must have met all other PBS eligibility criteria that a non-Grandfather patient would ordinarily be required to meet, meaning that at the time non-PBS supply was commenced, the patient: (i) had either one of (1) persistent carcinoma, (2) recurrent carcinoma, (3) metastatic carcinoma of the cervix; (ii) had a WHO performance status no higher than 1; (iii) was unsuitable for curative treatment with either of (1) surgical resection, (2) radiation; (iv) had not received prior treatment for this PBS indication; (v) was treated concomitantly with platinum-based chemotherapy agent, plus paclitaxel, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a total of (i) 24 months, (ii) 35 doses (based on a 3-weekly dose regimen), (iii) 17 doses (based on a 6-weekly dose regimen) whichever comes first from the first dose of this drug regardless if it was PBS/non-PBS subsidised.

Treatment criteria:

- Patient must be undergoing treatment with this drug administered once every 3 weeks - prescribe up to 6 repeat prescriptions; OR
- Patient must be undergoing treatment with this drug administered once every 6 weeks - prescribe up to 3 repeat prescriptions.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Injection

13635P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	400 mg	6	..	*15383.62	31.60	Keytruda [MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)

VEGF/VEGFR (Vascular Endothelial Growth Factor) inhibitors

▪ **BEVACIZUMAB**

Injection

12479T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1800 mg	7	..	*1714.34	31.60	Abevmy [SZ] (bevacizumab 100 mg/4 mL injection, 4 mL vial) Abevmy [SZ] (bevacizumab 400 mg/16 mL injection, 16 mL vial) Bevaciptin [LR] (bevacizumab 100 mg/4 mL injection, 4 mL vial) Bevaciptin [LR] (bevacizumab 400 mg/16 mL injection, 16 mL vial) Mvasi [AN] (bevacizumab 100 mg/4 mL injection, 4 mL vial) Mvasi [AN] (bevacizumab 400 mg/16 mL injection, 16 mL vial)

Other monoclonal antibodies and antibody drug conjugates

▪ **BLINATUMOMAB**

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome, neurological toxicities and reactivation of John Cunningham virus (JC) viral infection.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must be relapsed or refractory B-precursor cell ALL, with an Eastern Cooperative Oncology Group (ECOG) performance status of 2 or less, **AND**
- The condition must not be present in the central nervous system or testis, **AND**
- Patient must have previously received a tyrosine kinase inhibitor (TKI) if the condition is Philadelphia chromosome positive, **AND**
- Patient must have received intensive combination chemotherapy for initial treatment of ALL or for subsequent salvage therapy, **AND**
- Patient must not have received more than 1 line of salvage therapy, **AND**
- The condition must be one of the following: (i) untreated with this drug for measurable residual disease, (ii) treated with this drug for measurable residual disease, but the condition has not relapsed within 6 months of completing that course of treatment, **AND**
- The condition must have more than 5% blasts in bone marrow, **AND**
- The treatment must not be more than 2 treatment cycles under this restriction in a lifetime.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 9 days of the first cycle and the first 2 days of the second cycle. For all subsequent cycle starts and re-initiation (e.g. if treatment is interrupted for 4 or more hours), supervision by a health care professional or hospitalisation is recommended.

An amount of 651 microgram will be sufficient for a continuous infusion of blinatumomab over 28 days in cycle 1. An amount of 784 microgram, which may be obtained under Induction treatment - balance of supply restriction, will be sufficient for a continuous infusion of blinatumomab over 28 days in cycle 2.

Blinatumomab is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Acute Lymphoblastic Leukaemia PBS Authority Application - Supporting Information Form; and
- (3) date of most recent chemotherapy, and if this was the initial chemotherapy regimen or salvage therapy, including what line of salvage; and
- (4) if applicable, the date of completion of blinatumomab treatment for measurable residual disease and the date of the patient's subsequent relapse; and
- (5) the percentage blasts in bone marrow count that is no more than 4 weeks old at the time of application.

Injection

11118E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	651 mcg	*66317.34	31.60	Blinicyto [AN] (blinatumomab 38.5 microgram injection [1 vial] (&) inert substance solution [10 mL vial], 1 pack)

■ BLINATUMOMAB

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome, neurological toxicities and reactivation of John Cunningham virus (JC) viral infection.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Induction treatment - balance of supply

Clinical criteria:

- The condition must be relapsed or refractory B-precursor cell ALL, with an Eastern Cooperative Oncology Group (ECOG) performance status of 2 or less, **AND**
- The condition must not be present in the central nervous system or testis, **AND**
- Patient must have previously received a tyrosine kinase inhibitor (TKI) if the condition is Philadelphia chromosome positive, **AND**
- Patient must have received insufficient therapy with this agent for this condition under the Induction treatment restriction to complete a maximum of 2 treatment cycles in a lifetime.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 9 days of the first cycle and the first 2 days of the second cycle. For all subsequent cycle starts and re-initiation (e.g. if treatment is interrupted for 4 or more hours), supervision by a health care professional or hospitalisation is recommended.

An amount of 784 mcg will be sufficient for a continuous infusion of blinatumomab over 28 days in cycle 2.

Blinatumomab is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Injection

11120G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	784 mcg	*77355.46	31.60	Blinicyto [AN] (blinatumomab 38.5 microgram injection [1 vial] (&) inert substance solution [10 mL vial], 1 pack)

■ BLINATUMOMAB

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome, neurological toxicities and reactivation of John Cunningham virus (JC) viral infection.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note A complete remission is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a full recovery of peripheral blood counts with platelets of greater than 100,000 per microliter, and absolute neutrophil count (ANC) of greater than 1,000 per microliter.

Note A complete remission with partial haematological recovery is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a partial recovery of peripheral blood counts with platelets of greater than 50,000 per microliter, and absolute neutrophil count (ANC) of greater than 500 per microliter.

Note Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time when an assessment is required must cease PBS-subsidised therapy with this agent.

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Consolidation treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised induction treatment with this drug for this condition, **AND**
- Patient must have achieved a complete remission; OR
- Patient must have achieved a complete remission with partial haematological recovery, **AND**
- The treatment must not be more than 3 treatment cycles under this restriction in a lifetime, **AND**
- Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug.

Injection

11117D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	784 mcg	2	..	*77355.46	31.60	Blinicyto [AN] (blinatumomab 38.5 microgram injection [1 vial] (&) inert substance solution [10 mL vial], 1 pack)

▪ **BLINATUMOMAB**

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome, neurological toxicities and reactivation of John Cunningham virus (JC) viral infection.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note A complete remission is defined as bone marrow blasts of less than or equal to 5%, no evidence of disease and a full recovery of peripheral blood counts with platelets of greater than 100,000 per microliter, and absolute neutrophil count (ANC) of greater than 1,000 per microliter.

Authority required

Measurable residual disease of precursor B-cell acute lymphoblastic leukaemia (Pre-B-cell ALL)

Treatment Phase: Initial treatment of measurable residual disease of Pre-B-cell ALL

Treatment criteria:

- Must be treated by a physician experienced in the treatment of haematological malignancies.

Clinical criteria:

- Patient must have an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1, **AND**
- The condition must not be present in the central nervous system or testis, **AND**
- Patient must have achieved complete remission following intensive combination chemotherapy for initial treatment of acute lymphoblastic leukaemia (ALL) or for subsequent salvage therapy, **AND**
- Patient must have measurable residual disease based on measurement in bone marrow, documented after an interval of at least 2 weeks from the last course of systemic chemotherapy given as intensive combination chemotherapy treatment of ALL/as subsequent salvage therapy, whichever was the later, measured using flow cytometry/molecular methods,

AND

- The treatment must not be more than 2 treatment cycles under this restriction in a lifetime.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 days of the first cycle and the first 2 days of the second cycle.

For all subsequent cycle starts and re-initiation (e.g. if treatment is interrupted for four or more hours), supervision by a health care professional or hospitalisation is recommended.

An amount of 784 mcg will be sufficient for a continuous infusion of blinatumomab over 28 days in each cycle.

Blinatumomab is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Measurable residual disease positive Acute Lymphoblastic Leukaemia PBS Authority Application - Supporting Information Form; and
- (3) date of most recent chemotherapy, and if this was the initial chemotherapy regimen or salvage therapy; and
- (4) the percentage blasts in bone marrow count that is no more than 4 weeks old at the time of application.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Measurable residual disease of precursor B-cell acute lymphoblastic leukaemia (Pre-B-cell ALL)
Treatment Phase: Continuing treatment of previously measurable residual disease of Pre-B-cell ALL

Treatment criteria:

- Must be treated by a physician experienced in the treatment of haematological malignancies.

Clinical criteria:

- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition, **AND**
- Patient must have achieved a complete remission, **AND**
- The condition must be negative for measurable residual disease using the same method used to determine initial PBS eligibility, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition, **AND**
- The treatment must not be more than 2 treatment cycles under this restriction in a lifetime.

For all subsequent cycle starts and re-initiation (e.g. if treatment is interrupted for four or more hours), supervision by a health care professional or hospitalisation is recommended.

An amount of 784 microgram will be sufficient for a continuous infusion of blinatumomab over 28 days in each cycle.

Blinatumomab is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Injection

11850Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	784 mcg	1	..	*77355.46	31.60	Blinicyto [AN] (blinatumomab 38.5 microgram injection [1 vial] (&) inert substance solution [10 mL vial], 1 pack)

■ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

CD30 positive systemic anaplastic large cell lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be for curative intent, **AND**
- Patient must have undergone appropriate prior front-line curative intent chemotherapy, **AND**
- Patient must demonstrate relapsed or chemotherapy-refractory disease, **AND**
- Patient must have responded to PBS-subsidised treatment with this drug if previously used for initial treatment of CD30 positive peripheral T-cell lymphoma, non-cutaneous type, **AND**
- The treatment must not exceed 4 cycles under this restriction.

Applications for authorisation of initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail and must include:

- (a) details (date, unique identifying number or provider number) of a histology report showing evidence of the tumour's CD30 positivity; and
- (b) The date of initial diagnosis of systemic anaplastic large cell lymphoma; and
- (c) Dates of commencement and completion of front-line curative intent chemotherapy; and

(d) a declaration of whether the patient's disease is relapsed or refractory, and the date and means by which the patient's disease was assessed as being relapsed or refractory.

All reports must be documented in the patient's medical records.

If the application is submitted through HPOS form upload or mail, it must include:

(i) A completed authority prescription form; and

(ii) A completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

10166C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	3	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

CD30 positive systemic anaplastic large cell lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed 12 cycles under this restriction in a lifetime.

Injection

10171H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	11	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have undergone a primary autologous stem cell transplant (ASCT), **AND**
- Patient must have experienced a relapsed CD30+ Hodgkin lymphoma post ASCT; OR
- Patient must have experienced a refractory CD30+ Hodgkin lymphoma post ASCT, **AND**
- Patient must not receive more than 4 cycles of treatment under this restriction.

Applications for authorisation of initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail.

If the application is submitted through HPOS upload or mail, it must include:

(a) a completed authority prescription form; and

(b) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

11073T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	3	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ BRENTUXIMAB VEDOTIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have undergone an autologous stem cell transplant (ASCT) for this condition, **AND**
- Patient must not be suitable for ASCT for this condition; OR
- Patient must not be suitable for treatment with multi-agent chemotherapy for this condition, **AND**
- Patient must have experienced a relapsed CD30+ Hodgkin lymphoma following at least two prior treatments for this condition; OR
- Patient must have experienced a refractory CD30+ Hodgkin lymphoma following at least two prior treatments for this condition, **AND**
- Patient must not receive more than 4 cycles of treatment under this restriction.

Applications for authorisation of initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail.

If the application is submitted through HPOS upload or mail, it must include:

(a) a completed authority prescription form; and

(b) a completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

11079D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	3	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ BRENTUXIMAB VEDOTIN

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must not have undergone an autologous stem cell transplant (ASCT) for this condition, **AND**
- Patient must not be suitable for ASCT for this condition; OR
- Patient must not be suitable for treatment with multi-agent chemotherapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 12 cycles of treatment under this restriction.

The treatment must not exceed a total of 16 cycles of combined initial and continuing treatment in a lifetime.

Injection

11087M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	11	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ BRENTUXIMAB VEDOTIN

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed or Refractory Hodgkin lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have undergone a primary autologous stem cell transplant (ASCT) for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 12 cycles of treatment under this restriction.

The treatment must not exceed a total of 16 cycles of combined initial and continuing treatment in a lifetime.

Injection

11096B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	11	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

CD30 positive cutaneous T-cell lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have pathologically confirmed CD30 positive cutaneous T-cell lymphoma, **AND**
- Patient must have CD30 positivity of at least 3% of malignant cells, **AND**
- Patient must have a diagnosis of mycosis fungoides; OR
- Patient must have a diagnosis of Sezary syndrome; OR
- Patient must have a diagnosis of primary cutaneous anaplastic large cell lymphoma, **AND**
- Patient must have received prior systemic treatment for this condition, **AND**
- The condition must be relapsed or refractory, **AND**
- The treatment must not exceed 4 cycles under this restriction in a lifetime, **AND**
- The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition.

The authority application must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail and must include:

(a) details (date, unique identifying number/code or provider number) of the histopathology report from an Approved Pathology Authority demonstrating the patient has a diagnosis of either mycosis fungoides, Sezary syndrome or primary cutaneous anaplastic large cell lymphoma; and

(b) details (date, unique identifying number/code or provider number) of a histology report on the tumour sample or of a flow cytometric analysis of lymphoma cells of the blood showing CD30 positivity of at least 3% of malignant cells; and

(c) Date of commencement and completion of the most recent prior systemic treatment.

All reports must be documented in the patient's medical records.

If the application is submitted through HPOS form upload or mail, it must include:

(i) A completed authority prescription form; and

(ii) A completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

11660Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	180 mg	3	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

CD30 positive cutaneous T-cell lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - Patient must have achieved an objective response with this drug, **AND**
 - Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
 - The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this condition, **AND**
 - The treatment must not exceed 12 cycles under this restriction in a lifetime.
- An objective response is defined as the demonstration of response by clinical observation of skin lesions, or response by positron-emission tomography (PET) and/or computed tomography (CT) standard criteria.

Injection

11664X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	180 mg	11	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note This product is not PBS-subsidised for the treatment of previously untreated CD30 positive cutaneous T-cell lymphoma.

Authority required

CD30 positive peripheral T-cell lymphoma, non-cutaneous type

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have histological confirmation of CD30 expression in at least 3% of malignant cells, **AND**
- The treatment must be for first line therapy for this condition, **AND**
- The treatment must be for curative intent, **AND**
- The treatment must be in combination with cyclophosphamide, doxorubicin and prednisone, **AND**
- The treatment must not be more than 6 treatment cycles under this restriction in a lifetime.

Applications for authorisation of initial treatment must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail and must include:

(a) details (date, unique identifying number/code or provider number) of a histology report on the tumour sample from an Approved Pathology Authority showing CD30 positivity of at least 3% malignant cells; and

(b) The date of initial diagnosis of Peripheral T-cell lymphoma.

All reports must be documented in the patient's medical records.

If the application is submitted through HPOS form upload or mail, it must include:

(i) A completed authority prescription form; and

(ii) A completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

Injection

12646N	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	5	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **BRENTUXIMAB VEDOTIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Note This product is not PBS-subsidised for the treatment of previously untreated CD30 positive cutaneous T-cell lymphoma.

Authority required

CD30 positive peripheral T-cell lymphoma, non-cutaneous type

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with cyclophosphamide, doxorubicin and prednisone, **AND**
- Patient must have completed 6 initial cycles of PBS-subsidised treatment with this drug for this indication, **AND**
- Patient must have achieved at least a partial response to the 6 initial cycles of treatment with a combination of this drug and cyclophosphamide, doxorubicin and prednisone for this indication, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be more than 2 treatment cycles under this restriction in a lifetime.

Partial response is defined using Lugano Response Criteria for Non-Hodgkin Lymphoma as:

- (a) Positron emission tomography-based response: lymph nodes and extralymphatic sites - a score of 4 (uptake moderately > liver), or 5 (uptake markedly higher than liver and/or new lesions), with reduced uptake compared with baseline and residual mass(es) of any size; nonmeasured lesions - not applicable; organ enlargement - not applicable; new lesions - none; bone marrow - residual uptake higher than uptake in normal marrow but reduced compared with baseline (diffuse uptake compatible with reactive changes from chemotherapy allowed). If there are persistent focal changes in the marrow in the context of a nodal response, consideration should be given to further evaluation with MRI or biopsy or an interval scan; OR
- (b) Computed tomography-based response: lymph nodes and extralymphatic sites - greater than or equal to 50% decrease in the sum of the product of the perpendicular diameters for multiple lesions, of up to six (6) target measurable nodes and extranodal sites; non-measured lesions - absent/normal, regressed but no increase; new lesions - none; bone marrow - not applicable.

Injection

12657E	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	200 mg	1	..	*18708.62	31.60	Adcetris [TK] (brentuximab vedotin 50 mg injection, 1 vial)

▪ **ELOTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with lenalidomide and dexamethasone, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

12983H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	5	..	*5038.65	31.60	Empliciti [BQ] (elotuzumab 300 mg injection, 1 vial) Empliciti [BQ] (elotuzumab 400 mg injection, 1 vial)

▪ **ELOTUZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required

Relapsed and/or refractory multiple myeloma
Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with lenalidomide and dexamethasone, **AND**
- Patient must have progressive disease after at least one prior therapy, **AND**
- Patient must have undergone or be ineligible for a stem cell transplant, **AND**
- Patient must not have previously received this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

12989P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	9	..	*5038.65	31.60	Empliciti [BQ] (elotuzumab 300 mg injection, 1 vial) Empliciti [BQ] (elotuzumab 400 mg injection, 1 vial)

▪ **ENFORTUMAB VEDOTIN**

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14416

Locally advanced (Stage III) or metastatic (Stage IV) urothelial cancer

Clinical criteria:

- The condition must have progressed on/following both: (i) platinum-based chemotherapy, (ii) programmed cell death 1/ligand 1 (PD-1/PD-L1) inhibitor therapy; OR
- The condition must have progressed on/following platinum-based chemotherapy, whilst PD-1/PD-L1 inhibitor therapy resulted in an intolerance that required treatment cessation, **AND**
- Patient must have/have had a WHO performance status score of no greater than 1 at treatment initiation with this drug. **AND**

• The treatment must be the sole PBS-subsidised systemic anti-cancer therapy for this PBS indication.

Treatment criteria:

- Patient must be undergoing treatment with this drug for the first time; OR
- Patient must be undergoing continuing treatment with this drug, with each of the following being true: (i) all other PBS eligibility criteria in this restriction are met, (ii) disease progression is absent.

Injection

13648H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	125 mg	8	..	*6212.01	31.60	Padcev [LL] (enfortumab vedotin 20 mg injection, 1 vial) Padcev [LL] (enfortumab vedotin 30 mg injection, 1 vial)

▪ **GEMTUZUMAB OZOGAMICIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Acute Myeloid Leukaemia
Treatment Phase: Induction treatment

Clinical criteria:

- Patient must have confirmed CD33-positive AML prior to initiation of treatment, **AND**
- The condition must be de novo, **AND**

- The condition must be previously untreated at the time of initiation (except for prior essential treatment with hydroxyurea or leukapheresis for patients with hyperleukocytic AML), **AND**
 - Patient must have confirmed intermediate/favourable cytogenetic risk; OR
 - Patient must have unknown cytogenetic risk due to inconclusive test results, **AND**
 - Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score of 2 or less, **AND**
 - The condition must not be acute promyelocytic leukaemia, **AND**
 - The treatment must be in combination with standard intensive remission induction chemotherapy for this condition, which must include cytarabine and an anthracycline, **AND**
 - The treatment must not be used in combination with a tyrosine kinase inhibitor, **AND**
 - The condition must not be internal tandem duplication (ITD) or tyrosine kinase domain (TKD) FMS tyrosine kinase 3 (FLT3) mutation positive, **AND**
 - Patient must not receive more than 1 induction cycle under this restriction in a lifetime.
- This drug is not PBS-subsidised if it is prescribed to an in-patient in a public hospital setting.

Injection

12844B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	5 mg	2	..	*9255.62	31.60	Mylotarg [PF] (gemtuzumab ozogamicin 5 mg injection, 1 vial)

▪ **GEMTUZUMAB OZOGAMICIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Consolidation treatment

Clinical criteria:

- Patient must have achieved a complete remission following induction treatment with this drug for this condition, **AND**
- The treatment must be in combination with standard intensive remission consolidation chemotherapy for this condition, which must include cytarabine and an anthracycline, **AND**
- Patient must not receive more than 2 consolidation cycles under this restriction in a lifetime.

This drug is not PBS-subsidised if it is prescribed to an in-patient in a public hospital setting.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Complete remission following induction is defined as fewer than 5% blasts in a normocellular marrow and an absolute neutrophil count of more than 1.0×10^9 cells/L with a platelet count of 100×10^9 /L or more in the peripheral blood in the absence of transfusion.

Progressive disease is defined as the presence of any of the following:

- a) Leukaemic cells in the CSF;
- b) Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- c) Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- d) Extramedullary leukaemia.

Injection

12861X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	5 mg	1	..	*9255.62	31.60	Mylotarg [PF] (gemtuzumab ozogamicin 5 mg injection, 1 vial)

▪ **IPILIMUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

11478

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial combination treatment (with nivolumab) as first-line drug therapy

Clinical criteria:

- The condition must be squamous type non-small cell lung cancer (NSCLC), **AND**
- Patient must not have previously been treated for this condition in the metastatic setting, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The condition must not have evidence of an activating epidermal growth factor receptor (EGFR) gene or an anaplastic lymphoma kinase (ALK) gene rearrangement or a c-ROS proto-oncogene 1 (ROS1) gene arrangement in tumour material, **AND**
- The treatment must be in combination with platinum-based chemotherapy for the first two cycles, **AND**
- The treatment must be in combination with nivolumab.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

11391

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing combination treatment (with nivolumab) of first-line drug therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed 24 months in total, measured from the initial dose, or, must not extend beyond disease progression, whichever comes first, **AND**
- The treatment must be in combination with nivolumab.

Injection

12322M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	4	..	*16122.48	31.60	Yervoy [BQ] (ipilimumab 50 mg/10 mL injection, 10 mL vial)

■ **IPILIMUMAB**

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Authority required (STREAMLINED)

11930

Unresectable malignant mesothelioma

Clinical criteria:

- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be in combination with PBS-subsidised nivolumab for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must not exceed a maximum total of 24 months in a lifetime for this condition.

Injection

12583G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	3	..	*16122.48	31.60	Yervoy [BQ] (ipilimumab 50 mg/10 mL injection, 10 mL vial)

■ **IPILIMUMAB**

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note In the first few months after start of immunotherapy, some patients can have a transient tumour flare with subsequent disease response. When progression is suspected, this should be confirmed through a confirmatory scan, taken at least 4 weeks later.

Note A prognostic International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) survival risk score can be calculated here: <https://www.mdcalc.com/imdc-international-metastatic-renal-cell-carcinoma>.

One point is assigned for each of:

- a time of diagnosis to systemic therapy of less than 1 year
- a Karnofsky Performance Status of less than 80%
- a haemoglobin less than the lower limit of normal
- a corrected calcium level greater than the upper limit of normal
- a neutrophil count greater than the upper limit of normal
- a platelet count greater than the upper limit of normal

Stated normal reference ranges may vary depending on the laboratory providing the measurement. 'Normal' here refers to the individual laboratory's stated normal reference range.

Favourable IMDC risk is a score of 0.

Intermediate IMDC risk is a score of 1 to 2.

Poor IMDC risk is a score of 3 to 6.

Document any IMDC risk score assessment in the patient's medical records.

Authority required (STREAMLINED)

8555

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Induction treatment

Clinical criteria:

- The condition must not have previously been treated, **AND**
- The condition must be classified as intermediate to poor risk according to the International Metastatic Renal Cell Carcinoma Database Consortium (IMDC), **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must be in combination with PBS-subsidised treatment with nivolumab as induction therapy for this condition.

Induction treatment with ipilimumab must not exceed a total of 4 doses at a maximum dose of 1 mg per kg every 3 weeks. The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

11628B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	3	..	*16122.48	31.60	Yervoy [BQ] (ipilimumab 50 mg/10 mL injection, 10 mL vial)

▪ **IPILIMUMAB**

Caution Combination treatment with ipilimumab and nivolumab is associated with an increased incidence and severity of immune-related adverse reactions compared with monotherapy with these agents. Monitoring at least prior to each dose is recommended.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

6562

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Induction treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received prior treatment with ipilimumab, **AND**
- The treatment must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks. The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Note For patients who commence therapy with ipilimumab:

- Decisions concerning efficacy should await completion of the entire induction regimen (four doses) and should be made in conjunction with established criteria for immunological responses. However induction may be ceased or delayed if symptomatic progressive disease or intolerable adverse events occur and if, in the opinion of the clinician, continuation of treatment poses a risk to the patient;
- Tumour responses may occur beyond the initial 12 week induction phase and evaluation for potential later responses should be undertaken regularly for the first year.

Authority required (STREAMLINED)

6585

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Re-induction treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have progressive disease after achieving an initial objective response to the most recent course of ipilimumab treatment (induction or re-induction), **AND**
- The treatment must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks.

An initial objective response to treatment is defined as either:

- sustained stable disease of greater than or equal to 3 months duration measured from at least 2 weeks after the date of completion of the most recent course of ipilimumab; or
- a partial or complete response.

The patient's body weight must be documented in the patient's medical records at the time treatment with ipilimumab is initiated.

Authority required (STREAMLINED)

14808

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Induction treatment

Clinical criteria:

- Patient must not have received prior treatment with nivolumab plus relatlimab, ipilimumab or a PD-1 (programmed cell death-1) inhibitor for the treatment of unresectable Stage III or Stage IV malignant melanoma, **AND**
- Patient must have an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1, **AND**
- The condition must not be ocular or uveal melanoma, **AND**
- The treatment must be in combination with PBS-subsidised treatment with nivolumab as induction therapy for this condition.

Induction treatment with nivolumab must not exceed a total of 4 doses at a maximum dose of 1 mg per kg every 3 weeks.

Induction treatment with ipilimumab must not exceed a total of 4 doses at a maximum dose of 3 mg per kg every 3 weeks.

The patient's body weight must be documented in the patient's medical records at the time treatment is initiated.

Injection

2641B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	360 mg	3	..	*42845.58	31.60	Yervoy [BQ] (ipilimumab 200 mg/40 mL injection, 40 mL vial) Yervoy [BQ] (ipilimumab 50 mg/10 mL injection, 10 mL vial)

■ SACITUZUMAB GOVITECAN

Caution This medicine contains a cytotoxic component and causes chemotherapy-like toxicity, in particular, it can cause severe or life-threatening neutropenia and severe diarrhoea. For further information, refer to the Product Information.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12669

Unresectable locally advanced or metastatic triple-negative breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this PBS indication.

Injection

12945H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	13	..	*10480.75	31.60	Trodely [GI] (sacituzumab govitecan 180 mg injection, 1 vial)

■ SACITUZUMAB GOVITECAN

Caution This medicine contains a cytotoxic component and causes chemotherapy-like toxicity, in particular, it can cause severe or life-threatening neutropenia and severe diarrhoea. For further information, refer to the Product Information.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12656

Unresectable locally advanced or metastatic triple-negative breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have progressive disease following two or more prior systemic therapies, at least one of them in the locally advanced or metastatic setting, **AND**
- The condition must be inoperable, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 1 prior to treatment initiation, **AND**
- The treatment must be the sole PBS-subsidised therapy for this PBS indication.

Injection

12966K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1200 mg	7	..	*10480.75	31.60	Trodely [GI] (sacituzumab govitecan 180 mg injection, 1 vial)

OTHER ANTINEOPLASTIC AGENTS

Platinum compounds

■ CARBOPLATIN

Injection

4309T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	900 mg	5	..	*141.78	31.60	Carboplatin Accord [OC] (carboplatin 450 mg/45 mL injection, 45 mL vial)

■ CISPLATIN

Injection

4319H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	220 mg	14	..	*137.59	31.60	Cisplatin Accord [OC] (cisplatin 100 mg/100 mL injection, 100 mL vial) Cisplatin Accord [OC] (cisplatin 50 mg/50 mL injection, 50 mL vial)

▪ **OXALIPLATIN**

Injection

4542C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	300 mg	11	..	*133.52	31.60	Oxaliplatin Accord [OC] (oxaliplatin 100 mg/20 mL injection, 20 mL vial) Oxaliplatin SUN [RA] (oxaliplatin 100 mg/20 mL injection, 20 mL vial) Oxaliplatin SUN [RA] (oxaliplatin 200 mg/40 mL injection, 40 mL vial)

Proteasome inhibitors

▪ **BORTEZOMIB**

Restricted benefit

Multiple myeloma

Restricted benefit

Newly diagnosed systemic light chain amyloidosis

Treatment Phase: Administration on Days 1, 8, 15 and 22 of six treatment cycles (28 days per cycle) in total

Treatment criteria:

- Patient must be undergoing concurrent treatment with PBS-subsidised daratumumab for this PBS indication.

Injection

12227M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3000 mcg	15	..	*166.09	31.60	Bortezom [CR] (bortezomib 3.5 mg injection, 1 vial) Bortezomib Accord [OC] (bortezomib 1 mg injection, 1 vial) Bortezomib Accord [OC] (bortezomib 2.5 mg/mL injection, 1 mL vial) Bortezomib Accord [OC] (bortezomib 3.5 mg injection, 1 vial) Bortezomib Accord [OC] (bortezomib 3.5 mg/1.4 mL injection, 1.4 mL vial) Bortezomib Baxter [BX] (bortezomib 3.5 mg injection, 1 vial) BORTEZOMIB EUGIA [YG] (bortezomib 3.5 mg injection, 1 vial) Bortezomib Ever Pharma [IT] (bortezomib 2.5 mg/mL injection, 1 mL vial) Bortezomib Ever Pharma [IT] (bortezomib 3.5 mg/1.4 mL injection, 1.4 mL vial) Bortezomib Juno [JU] (bortezomib 2.5 mg injection, 1 vial) Bortezomib Juno [JU] (bortezomib 3.5 mg injection, 1 vial) Bortezomib Sandoz [SZ] (bortezomib 3.5 mg injection, 1 vial) BORTEZOMIB-TEVA [TB] (bortezomib 3.5 mg injection, 1 vial) DBL Bortezomib [PF] (bortezomib 1 mg injection, 1 vial) DBL Bortezomib [PF] (bortezomib 2.5 mg injection, 1 vial) DBL Bortezomib [PF] (bortezomib 3 mg injection, 1 vial) DBL Bortezomib [PF] (bortezomib 3.5 mg injection, 1 vial)

▪ **CARFILZOMIB**

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum amount or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12934

Multiple myeloma

Treatment Phase: Initial treatment - twice weekly treatment regimen

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must have progressive disease after at least one prior therapy, **AND**
- Patient must have undergone or be ineligible for a stem cell transplant, **AND**
- Patient must not have previously received this drug for this condition, **AND**
- Patient must not receive more than three cycles of treatment under this restriction.

Progressive disease is defined as at least 1 of the following:

- at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or

- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Authority required (STREAMLINED)

12930

Multiple myeloma

Treatment Phase: Continuing treatment - twice weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - The treatment must be in combination with dexamethasone, **AND**
 - Patient must not develop disease progression while receiving treatment with this drug for this condition, **AND**
 - Patient must not receive more than 3 cycles of treatment per continuing treatment course authorised under this restriction.
- Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

11229B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	120 mg	17	..	*2499.60	31.60	Kyprolis [AN] (carfilzomib 10 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 30 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 60 mg injection, 1 vial)

▪ **CARFILZOMIB**

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum amount or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

12694

Multiple myeloma

Treatment Phase: Initial treatment - once weekly treatment regimen

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must have progressive disease after at least one prior therapy, **AND**
- Patient must have undergone or be ineligible for a stem cell transplant, **AND**
- Patient must not have previously received this drug for this condition, **AND**
- Patient must not receive more than three cycles of treatment under this restriction.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Authority required (STREAMLINED)

12849

Multiple myeloma

Treatment Phase: Continuing treatment - once weekly treatment regimen

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must not develop disease progression while receiving treatment with this drug for this condition, **AND**
- Patient must not receive more than 3 cycles of treatment per continuing treatment course authorised under this restriction. Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

12244K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	160 mg	8	..	*3303.27	31.60	Kyprolis [AN] (carfilzomib 10 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 30 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 60 mg injection, 1 vial)

▪ **CARFILZOMIB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

14389

Relapsed and/or refractory multiple myeloma

Treatment Phase: Initial treatment for Cycles 1 to 3

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with lenalidomide and dexamethasone, **AND**
- Patient must have progressive disease after at least one prior therapy, **AND**
- Patient must not have previously received this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Provide details of the histological diagnosis of multiple myeloma, prior treatments including name(s) of drug(s) and date of the most recent treatment cycle; the basis of the diagnosis of progressive disease or failure to respond; and which disease activity parameters will be used to assess response once only through the Authority application for lenalidomide.

Authority required (STREAMLINED)

14363

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment for Cycles 3 to 12

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with lenalidomide and dexamethasone, **AND**
- Patient must not have progressive disease while receiving treatment with this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or

- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Authority required (STREAMLINED)

14364

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment for Cycles 13 onwards

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with lenalidomide and dexamethasone, **AND**
- Patient must not have progressive disease while receiving treatment with this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Injection

13638T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	60 mg	17	..	*1294.11	31.60	Kyprolis [AN] (carfilzomib 10 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 30 mg injection, 1 vial) Kyprolis [AN] (carfilzomib 60 mg injection, 1 vial)

Other antineoplastic agents

▪ **ARSENIC**

Authority required (STREAMLINED)

6018

Acute promyelocytic leukaemia

Treatment Phase: Induction and consolidation treatment

Clinical criteria:

- The condition must be characterised by the presence of the t(15:17) translocation or PML/RAR-alpha fusion gene transcript.

Injection

10691Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	18 mg	140	..	*205.54	31.60	Arsenic Trioxide Accord [OC] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials) Arsenic Trioxide-AFT [AE] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL ampoules) Arsenic Trioxide Juno [JU] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials) Phenasen [FF] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials)

▪ **ARSENIC**

Authority required (STREAMLINED)

4793

Acute promyelocytic leukaemia

Treatment Phase: Induction and consolidation treatment

Clinical criteria:

- The condition must be characterised by the presence of the t(15:17) translocation or PML/RAR-alpha fusion gene transcript, **AND**

- The condition must be relapsed, **AND**
- Patient must be arsenic naive at induction.

Authority required (STREAMLINED)

5997

Acute promyelocytic leukaemia

Clinical criteria:

- The condition must be characterised by the presence of the t(15:17) translocation or PML/RAR-alpha fusion gene transcript.

Injection

4371C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	18 mg	89	..	*205.54	31.60	Arsenic Trioxide Accord [OC] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials) Arsenic Trioxide-AFT [AE] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL ampoules) Arsenic Trioxide Juno [JU] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials) Phenasen [FF] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials)

▪ **ERIBULIN**

Note A patient who has progressive disease with eribulin is no longer eligible for PBS-subsidised eribulin.

Authority required (STREAMLINED)

4649

Locally advanced or metastatic breast cancer

Clinical criteria:

- Patient must have progressive disease, **AND**
- Patient must have failed at least two prior chemotherapeutic regimens for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Injection

10144X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3 mg	13	..	*772.62	31.60	Halaven [EI] (eribulin mesilate 1 mg/2 mL injection, 2 mL vial)

▪ **ERIBULIN**

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7258

Advanced (unresectable and/or metastatic) liposarcoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have an ECOG performance status of 2 or less, **AND**
- The condition must be dedifferentiated, myxoid, round-cell or pleomorphic subtype, **AND**
- Patient must have received prior chemotherapy treatment including an anthracycline and ifosfamide (unless contraindicated) for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Population criteria:

- Patient must be aged 18 years or older.

Authority required (STREAMLINED)

7280

Advanced (unresectable and/or metastatic) liposarcoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop progressive disease while being treated with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Population criteria:

- Patient must be aged 18 years or older.

Injection

11212D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	3 mg	7	..	*772.62	31.60	Halaven [EI] (eribulin mesilate 1 mg/2 mL injection, 2 mL vial)

▪ **TEBENTAFUSP**

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome (CRS).

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**15085**

Advanced (unresectable or metastatic) uveal melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition; OR
- Patient must have previously received inpatient treatment with this drug for this condition in the public hospital setting, **AND**
- Patient must not receive PBS-subsidised treatment with this drug for this condition if it is no longer determined to be clinically beneficial by the treating clinician.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 doses (on Days 1, 8 and 15) and for at least 16 hours after each infusion is completed. If the patient does not experience hypotension that is Grade 2 or worse (requiring medical intervention) with the third dose, subsequent doses can be administered in an appropriate outpatient/ambulatory care setting. Supervision by a health care professional is recommended for a minimum of 30 minutes following each infusion.

Injection

13823M	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	136 mcg	7	..	*35255.74	31.60	Kimtrak [WM] (tebentafusp 100 microgram/0.5 mL injection, 0.5 mL vial)

TEBENTAFUSP

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome (CRS).

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**14821**

Advanced (unresectable or metastatic) uveal melanoma

Treatment Phase: Initial treatment - day 8

Clinical criteria:

- Patient must have HLA-A*02:01-positive disease, **AND**
- Patient must have previously received PBS-subsidised initial day 1 treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 doses (on Days 1, 8 and 15) and for at least 16 hours after each infusion is completed. If the patient does not experience hypotension that is Grade 2 or worse (requiring medical intervention) with the third dose, subsequent doses can be administered in an appropriate outpatient/ambulatory care setting. Supervision by a health care professional is recommended for a minimum of 30 minutes following each infusion.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Positive HLA-A*02:01 assessment must be documented in the patient's medical records.

Injection

13819H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	30 mcg	*17672.18	31.60	Kimtrak [WM] (tebentafusp 100 microgram/0.5 mL injection, 0.5 mL vial)

TEBENTAFUSP

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome (CRS).

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**14825**

Advanced (unresectable or metastatic) uveal melanoma

Treatment Phase: Initial treatment - day 15

Clinical criteria:

- Patient must have HLA-A*02:01-positive disease, **AND**
- Patient must have previously received PBS-subsidised initial day 8 treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 doses (on Days 1, 8 and 15) and for at least 16 hours after each infusion is completed. If the patient does not experience hypotension that is Grade 2 or worse (requiring medical intervention) with the third dose, subsequent doses can be administered in an

appropriate outpatient/ambulatory care setting. Supervision by a health care professional is recommended for a minimum of 30 minutes following each infusion.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Positive HLA-A*02:01 assessment must be documented in the patient's medical records.

Injection

13831Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	68 mcg	*17672.18	31.60	Kimmtrak [WM] (tebentafusp 100 microgram/0.5 mL injection, 0.5 mL vial)

▪ **TEBENTAFUSP**

Caution Careful monitoring of patients is required due to risk of developing life-threatening Cytokine Release Syndrome (CRS).

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum amount or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Advanced (unresectable or metastatic) uveal melanoma

Treatment Phase: Initial treatment - day 1

Clinical criteria:

- Patient must have HLA-A*02:01-positive disease, **AND**
- Patient must have uveal melanoma that has been confirmed either (i) histologically, (ii) cytologically, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have received prior systemic therapy for metastatic disease.

Population criteria:

- Patient must be at least 18 years of age.

According to the TGA-approved Product Information, hospitalisation is recommended at minimum for the first 3 doses (on Days 1, 8 and 15) and for at least 16 hours after each infusion is completed. If the patient does not experience hypotension that is Grade 2 or worse (requiring medical intervention) with the third dose, subsequent doses can be administered in an appropriate outpatient/ambulatory care setting. Supervision by a health care professional is recommended for a minimum of 30 minutes following each infusion.

This drug is not PBS-subsidised if it is administered to an in-patient in a public hospital setting.

Positive HLA-A*02:01 assessment must be documented in the patient's medical records.

Injection

13832B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	20 mcg	*17672.18	31.60	Kimmtrak [WM] (tebentafusp 100 microgram/0.5 mL injection, 0.5 mL vial)

Related Pharmaceutical Benefits for Public Hospital use

ALIMENTARY TRACT AND METABOLISM.....	156
ANTIEMETICS AND ANTINAUSEANTS.....	156
ANTIEMETICS AND ANTINAUSEANTS	156
<hr/>	
ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	159
ANTINEOPLASTIC AGENTS	159
MONOCLONAL ANTIBODIES AND ANTIBODY DRUG CONJUGATES.....	159
IMMUNOSTIMULANTS	163
IMMUNOSTIMULANTS	163
<hr/>	
VARIOUS	163
ALL OTHER THERAPEUTIC PRODUCTS	163
ALL OTHER THERAPEUTIC PRODUCTS.....	163

ALIMENTARY TRACT AND METABOLISM

ANTIEMETICS AND ANTINAUSEANTS

ANTIEMETICS AND ANTINAUSEANTS

Serotonin (5HT₃) antagonists

▪ FOSNETUPITANT + PALONOSETRON

Note This medicine is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Note Various sources of information outline the emetic risk associated with cancer treatment. Examples include the National Comprehensive Cancer Network guidelines (USA), eviQ guidelines and approved Product Information of individual drugs. These examples are not a comprehensive list of which anti-cancer drugs that have moderate to high emesis risk.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Nausea and vomiting

Clinical criteria:

- The treatment must be for prevention of nausea and vomiting associated with moderate to highly emetogenic anti-cancer therapy, **AND**
- The treatment must be in combination with dexamethasone, unless contraindicated, **AND**
- Patient must be unable to swallow; OR
- Patient must be contraindicated to oral anti-emetics.

fosnetupitant 235 mg + palonosetron 250 microgram injection, 1 vial

13650K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	91.27	31.60	Akynzeo IV [JZ]

▪ GRANISETRON

Restricted benefit

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

granisetron 3 mg/3 mL injection, 3 mL ampoule

5899L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2.50	3.90	^a Granisetron-AFT [AE] ^a Kytril [IX]	^a Granisetron Kabi [PK]

granisetron 2 mg tablet, 1

5898K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	2	*16.58	17.98	Kytril [IX]

▪ NETUPITANT + PALONOSETRON

Note This medicine is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Note Various sources of information outline the emetic risk associated with cancer treatment. Examples include the National Comprehensive Cancer Network guidelines (USA), eviQ guidelines and approved Product Information of individual drugs. These examples are not a comprehensive list of which anti-cancer drugs that have moderate to high emesis risk.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

14443

Nausea and vomiting

Clinical criteria:

- The treatment must be in combination with dexamethasone, unless contraindicated, **AND**
- The treatment must be for prevention of nausea and vomiting associated with moderate to highly emetogenic anti-cancer therapy.

netupitant 300 mg + palonosetron 500 microgram capsule, 1

10714X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	‡1	5	..	81.66	31.60	Akynzeo [JZ]

▪ ONDANSETRON

Restricted benefit

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

ondansetron 4 mg/5 mL oral liquid, 50 mL

5848T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer	
	‡1	80.78	31.60	Zofran syrup 50 mL [AS]	

ondansetron 4 mg tablet, 4

5967C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3.50	4.90	^a APO-Ondansetron [TX] ^a Ondansetron-DRLA [RZ] ^a Ondansetron SZ [HX] ^a Zofran [AS]	^a APX-Ondansetron [TY] ^a Ondansetron Mylan Tablets [AF] ^a Ondansetron Tablets Viatris [AL] ^a Zotren 4 [RF]

ondansetron 8 mg tablet, 4

5968D	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	4.33	5.73	^a APO-Ondansetron [TX] ^a Ondansetron-DRLA [RZ] ^a Ondansetron SZ [HX] ^a Zofran [AS]	^a APX-Ondansetron [TY] ^a Ondansetron Mylan Tablets [AF] ^a Ondansetron Tablets Viatris [AL] ^a Zotren 8 [RF]

■ ONDANSETRON

Restricted benefit

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

ondansetron 4 mg orally disintegrating tablet, 4

5857G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3.50	4.90	^a APX-Ondansetron ODT [TY] ^a Ondansetron ODT-DRLA [RZ] ^a Zotren ODT [RF]	^a Ondansetron Mylan ODT [AF] ^a Ondansetron SZ ODT [HX]

ondansetron 8 mg orally disintegrating tablet, 4

5858H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	4.33	5.73	^a APX-Ondansetron ODT [TY] ^a Ondansetron ODT-DRLA [RZ] ^a Ondansetron SZ ODT [HX]	^a Ondansetron Mylan ODT [AF] ^a Ondansetron ODT Viatris [AL] ^a Zotren ODT [RF]

■ PALONOSETRON

Note No increase in the maximum quantity or number of units may be authorised.

Note This drug is not PBS-subsidised for administration with oral 5-HT3 antagonists.

Restricted benefit

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

palonosetron 250 microgram/5 mL injection, 5 mL vial

5853C	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	9.91	11.31	^a Aloxi [JZ] ^a PALONOSETRON Medsurge [DZ]	^a Palonosetron Dr.Reddy's [RZ]

Other antiemetics

■ APREPITANT

Note Aprepitant is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

4223

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy, **AND**

- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following agents: altretamine; carmustine; cisplatin when a single dose constitutes a cycle of chemotherapy; cyclophosphamide at a dose of 1500 mg per square metre per day or greater; dacarbazine; procarbazine when a single dose constitutes a cycle of chemotherapy; streptozocin.

No more than 1 capsule of aprepitant 165 mg will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)

4216

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat breast cancer, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone, **AND**
- Patient must be scheduled to be co-administered cyclophosphamide and an anthracycline.

No more than 1 capsule of aprepitant 165 mg will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)

6464

Nausea and vomiting

Clinical criteria:

- The condition must be associated with moderately emetogenic cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone on day 1 of a chemotherapy cycle, **AND**
- Patient must have had a prior episode of chemotherapy induced nausea or vomiting, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following intravenous chemotherapy agents: arsenic trioxide; azacitidine; cyclophosphamide at a dose of less than 1500 mg per square metre per day; cytarabine at a dose of greater than 1 g per square metre per day; dactinomycin; daunorubicin; doxorubicin; epirubicin; fotemustine; idarubicin; ifosfamide; irinotecan; melphalan; methotrexate at a dose of 250 mg to 1 g per square metre; raltitrexed.

No more than 1 capsule of aprepitant 165 mg will be authorised per cycle of cytotoxic chemotherapy.

Concomitant use of a 5HT3 antagonist should not occur with aprepitant on days 2 and 3 of any chemotherapy cycle.

Authority required (STREAMLINED)

6383

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy, **AND**
 - The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone on day 1 of a chemotherapy cycle, **AND**
 - Patient must be scheduled to be administered a chemotherapy regimen that includes either carboplatin or oxaliplatin.
- No more than 1 capsule of aprepitant 165 mg will be authorised per cycle of cytotoxic chemotherapy.

Concomitant use of a 5HT3 antagonist should not occur with aprepitant on days 2 and 3 of any chemotherapy cycle.

aprepitant 165 mg capsule, 1

2550F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	55.17	31.60	^a Aprepitant APOTEX [TX]	^a APREPITANT SCP [XC]

▪ **FOSAPREPITANT**

Note This medicine is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

6886

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following agents: altretamine; carmustine; cisplatin when a single dose constitutes a cycle of chemotherapy; cyclophosphamide at a dose of 1500 mg per square metre per day or greater; dacarbazine; procarbazine when a single dose constitutes a cycle of chemotherapy; streptozocin.

No more than 1 vial of fosaprepitant 150 mg injection will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)

6891

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat breast cancer, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone, **AND**
- Patient must be scheduled to be co-administered cyclophosphamide and an anthracycline.

No more than 1 vial of fosaprepitant 150 mg injection will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)

6887

Nausea and vomiting

Clinical criteria:

- The condition must be associated with moderately emetogenic cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone on day 1 of a chemotherapy cycle, **AND**
- Patient must have had a prior episode of chemotherapy induced nausea or vomiting, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following intravenous chemotherapy agents: arsenic trioxide; azacitidine; cyclophosphamide at a dose of less than 1500 mg per square metre per day; cytarabine at a dose of greater than 1 g per square metre per day; dactinomycin; daunorubicin; doxorubicin; epirubicin; fotemustine; idarubicin; ifosfamide; irinotecan; melphalan; methotrexate at a dose of 250 mg to 1 g per square metre; raltitrexed.

No more than 1 vial of fosaprepitant 150 mg injection will be authorised per cycle of cytotoxic chemotherapy.

Concomitant use of a 5HT3 antagonist should not occur with fosaprepitant on days 2 and 3 of any chemotherapy cycle.

Authority required (STREAMLINED)

6852

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone on day 1 of a chemotherapy cycle, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes either carboplatin or oxaliplatin.

No more than 1 vial of fosaprepitant 150 mg injection will be authorised per cycle of cytotoxic chemotherapy.

Concomitant use of a 5HT3 antagonist should not occur with fosaprepitant on days 2 and 3 of any chemotherapy cycle.

fosaprepitant 150 mg injection, 1 vial

11103J	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	69.23	31.60	^a Emend IV [MK] ^a FOSAPREPITANT MEDSURGE [DZ]	^a FOSAPREPITANT-AFT [AE] ^a FOSAPREPITANT MSN [RQ]

■ **ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS**

■ **ANTINEOPLASTIC AGENTS**

MONOCLONAL ANTIBODIES AND ANTIBODY DRUG CONJUGATES

CD38 (Clusters of Differentiation 38) inhibitors

■ **DARATUMUMAB**

Note This drug is not PBS-subsidised for use in patients with multiple myeloma who have received two or more prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent, or, who are refractory to both a PI and an immunomodulatory agent, as monotherapy.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment of second-line drug therapy from week 25 until disease progression (administered every 4 weeks)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition. Progressive disease is defined as at least 1 of the following:
 - (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
 - (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
 - (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
 - (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
 - (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
 - (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
 - (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

daratumumab 1.8 g/15 mL injection, 15 mL vial

12682L	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7010.28	31.60	Darzalex SC [JC]

▪ DARATUMUMAB

Note This drug is not PBS-subsidised for use in patients with multiple myeloma who have received two or more prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent, or, who are refractory to both a PI and an immunomodulatory agent, as monotherapy.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Continuing treatment of second-line drug therapy for weeks 10 to 24 (administered every 3 weeks)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with bortezomib and dexamethasone, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

Progressive disease is defined as at least 1 of the following:

- at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- an increase in the size or number of lytic bone lesions (not including compression fractures); or
- at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

daratumumab 1.8 g/15 mL injection, 15 mL vial

12745T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	4	..	7010.28	31.60	Darzalex SC [JC]

▪ DARATUMUMAB

Note This drug is not PBS-subsidised for use in patients with multiple myeloma who have received two or more prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent, or, who are refractory to both a PI and an immunomodulatory agent, as monotherapy.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Relapsed and/or refractory multiple myeloma

Treatment Phase: Initial treatment as second-line drug therapy for weeks 1 to 9 (administered once weekly)

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be in combination with bortezomib and dexamethasone, **AND**
- Patient must have progressive disease after only one prior therapy (i.e. use must be as second-line drug therapy; use as third-line drug therapy or beyond is not PBS-subsidised).

Treatment criteria:

- Patient must be undergoing treatment with this drug in one of the following situations: (i) for the first time, irrespective of whether the diagnosis has been reclassified (i.e. the diagnosis has changed between multiple myeloma/amyloidosis), (ii) changing the drug's form (intravenous/subcutaneous) within the first 9 weeks of treatment for the same PBS indication.

Progressive disease is defined as at least 1 of the following:

- at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase in the difference between involved free light chain and uninvolved free light chain; or
- at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or

- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein. Details of: the histological diagnosis of multiple myeloma; prior treatments including name(s) of drug(s) and date of most recent treatment cycle; the basis of the diagnosis of progressive disease or failure to respond; and which disease activity parameters will be used to assess response, must be documented in the patient's medical records.

Confirmation of eligibility for treatment with current diagnostic reports of at least one of the following must be documented in the patient's medical records:

- (a) the level of serum monoclonal protein; or
- (b) Bence-Jones proteinuria - the results of 24-hour urinary light chain M protein excretion; or
- (c) the serum level of free kappa and lambda light chains; or
- (d) bone marrow aspirate or trephine; or
- (e) if present, the size and location of lytic bone lesions (not including compression fractures); or
- (f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
- (g) if present, the level of hypercalcaemia, corrected for albumin concentration.

As these parameters must be used to determine response, results for either (a) or (b) or (c) should be documented for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) must be documented in the patient's medical records. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (current serum M protein less than 10 g per L) must be documented in the patient's medical records.

A line of therapy is defined as 1 or more cycles of a planned treatment program. This may consist of 1 or more planned cycles of single-agent therapy or combination therapy, as well as a sequence of treatments administered in a planned manner.

A new line of therapy starts when a planned course of therapy is modified to include other treatment agents (alone or in combination) as a result of disease progression, relapse, or toxicity, with the exception to this being the need to attain a sufficient response for stem cell transplantation to proceed. A new line of therapy also starts when a planned period of observation off therapy is interrupted by a need for additional treatment for the disease.

daratumumab 1.8 g/15 mL injection, 15 mL vial

12746W	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	8	..	7010.28	31.60	Darzalex SC [JC]

▪ **DARATUMUMAB**

Note The intravenously administered presentation of this drug is not PBS listed for this indication at the request of the sponsor.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 888 333. Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au Applications for authorisation under this restriction should be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/hpos)

Alternatively, applications for authority to prescribe can be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
 Services Australia
 PBS Authorities
 GPO Box 9826
 [Your capital city]

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Newly diagnosed systemic light chain amyloidosis
 Treatment Phase: Initial treatment from week 0 to week 24

Clinical criteria:

- The condition must have histological evidence consistent with a diagnosis of systemic light-chain amyloidosis, **AND**
- The condition must be untreated with drug therapy, including this drug, irrespective of whether the diagnosis has been reclassified (i.e. the diagnosis changes between multiple myeloma/amyloidosis), **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score of no higher than 2 at treatment initiation.

Treatment criteria:

- Must be treated by a haematologist (this does not exclude treatment via a multidisciplinary team, but the PBS authority application must be sought by the treating haematologist), **AND**
- Patient must be undergoing concomitant treatment limited to each of: (i) bortezomib, (ii) cyclophosphamide, (iii) dexamethasone, at certain weeks of treatment as outlined in the drug's approved Product Information.

Related

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

The authority application must be made via the Online PBS Authorities System (real time assessment), or in writing via HPOS form upload or mail, and must include:

Details of the histological evidence supporting the diagnosis of systemic light chain amyloidosis, limited to: (i) the name of pathologist/pathology provider, (ii) the site of biopsy

If the application is submitted through HPOS form upload or mail, it must include:

- (i) A completed authority prescription form; and
- (ii) A completed authority application form relevant to the indication and treatment phase (the latest version is located on the website specified in the Administrative Advice).

daratumumab 1.8 g/15 mL injection, 15 mL vial

13201T	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	15	..	7010.28	31.60	Darzalex SC [JC]

▪ DARATUMUMAB

Note The intravenously administered presentation of this drug is not PBS listed for this indication at the request of the sponsor.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Newly diagnosed systemic light chain amyloidosis

Treatment Phase: Continuing treatment from week 25 onwards (administered once every four weeks)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Must be treated by a haematologist (this does not exclude treatment via a multidisciplinary team, but the PBS authority application must be sought by the treating haematologist), **AND**
- Patient must be undergoing continuing treatment that does not extend treatment duration beyond whichever comes first: (i) disease progression, (ii) 96 cumulative weeks from the first administered dose, once in a lifetime.

daratumumab 1.8 g/15 mL injection, 15 mL vial

13203X	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7010.28	31.60	Darzalex SC [JC]

HER2 (Human Epidermal Growth Factor Receptor 2) inhibitors

▪ TRASTUZUMAB

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

10212

Early HER2 positive breast cancer

Treatment Phase: 3 weekly treatment regimen

Clinical criteria:

- Patient must have undergone surgery (adjuvant) or be preparing for surgery (neoadjuvant), **AND**
 - The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
 - Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
 - Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.
- Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

trastuzumab 600 mg/5 mL injection, 5 mL vial

10743K	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1470.22	31.60	Herceptin SC [RO]

▪ TRASTUZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required (STREAMLINED)

9353

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) either in the primary tumour or a metastatic lesion, **AND**
 - The treatment must not be in combination with nab-paclitaxel, **AND**
 - The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.
- Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

trastuzumab 600 mg/5 mL injection, 5 mL vial

10811B	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	1470.22	31.60	Herceptin SC [RO]

▪ **TRASTUZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. Monday to Friday).

Authority required (STREAMLINED)

9462

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

trastuzumab 600 mg/5 mL injection, 5 mL vial

10817H	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1470.22	31.60	Herceptin SC [RO]

▪ **IMMUNOSTIMULANTS**

IMMUNOSTIMULANTS

Other immunostimulants

▪ **MYCOBACTERIUM BOVIS BCG DANISH STRAIN**

Restricted benefit

Primary and relapsing superficial urothelial carcinoma of the bladder

Mycobacterium bovis BCG Danish strain 30 mg injection, 4 vials

12925G	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	±3	1	..	*1317.00	31.60	VesiCulture [LM]

▪ **MYCOBACTERIUM BOVIS BCG TICE STRAIN**

Restricted benefit

Primary and relapsing superficial urothelial carcinoma of the bladder

Mycobacterium bovis BCG Tice strain 500 million CFU injection, 3 vials

5902P	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	399.49	31.60	OncoTICE [MK]

▪ **VARIOUS**

▪ **ALL OTHER THERAPEUTIC PRODUCTS**

ALL OTHER THERAPEUTIC PRODUCTS

Detoxifying agents for antineoplastic treatment

▪ **FOLINIC ACID**

folinic acid 100 mg/10 mL injection, 10 x 10 mL ampoules

1904F	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	43.80	31.60	Leucovorin Calcium (Pfizer Australia Pty Ltd) [PF]

folinic acid 50 mg/5 mL injection, 10 x 5 mL ampoules

1899Y	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	38.90	31.60	Leucovorin Calcium (Pfizer Australia Pty Ltd) [PF]

VARIOUS

▪ FOLINIC ACID

Restricted benefit

Megaloblastic anaemias

Clinical criteria:

- The condition must be a result of folic acid deficiency from the use of folic acid antagonists.

folinic acid 15 mg tablet, 10

5904R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	76.00	31.60	Leucovorin Calcium (Hospira Pty Limited) [PF]

▪ MESNA

Restricted benefit

Urothelial toxicity

Treatment Phase: Prophylaxis or reduction of toxicity

Clinical criteria:

- The treatment must be adjunctive therapy to ifosfamide or high dose cyclophosphamide.

mesna 1 g/10 mL injection, 15 x 10 mL ampoules

5961R	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	117.16	31.60	Uromitexan [BX]

mesna 400 mg/4 mL injection, 15 x 4 mL ampoules

5960Q	Max. Amount	No. of Rpts	Premium \$	DPMA \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	51.74	31.60	Uromitexan [BX]

Index of Manufacturers' Code

Code	Manufacturer
AE	AFT Pharmaceuticals (AU) Pty Ltd
AF	Alphapharm Pty Ltd
AL	Alphapharm Pty Ltd
AN	Amgen Australia Pty Limited
AP	AstraZeneca Pty Ltd
AS	Aspen Pharmacare Australia Pty Limited
BQ	Bristol-Myers Squibb Australia Pty Ltd
BX	Baxter Healthcare Pty Limited
CR	Pharmacor Pty Limited
DZ	Medsurge Healthcare Pty Ltd
EI	Eisai Australia Pty Ltd
EW	Celltrion Healthcare Australia Pty Ltd
FB	Pierre Fabre Australia Pty Ltd
FF	Phebra Pty Ltd
GI	Gilead Sciences Pty Limited
GK	GlaxoSmithKline Australia Pty Ltd
HX	Sandoz Pty Ltd
IT	InterPharma Pty Ltd
IX	Clinect Pty Ltd
JC	Janssen-Cilag Pty Ltd
JO	Juno Pharmaceuticals Pty Ltd
JU	Juno Pharmaceuticals Pty Ltd
JZ	Juniper Biologics Pty Ltd
LL	Astellas Pharma Australia Pty Ltd
LM	Link Medical Products Pty Ltd
LR	Cipla Australia Pty Ltd
MF	Mundipharma Pty Limited
MK	Merck Sharp & Dohme (Australia) Pty Ltd
OC	Accord Healthcare Pty. Ltd.
OD	Accord Healthcare Pty. Ltd.
OE	Omegapharm Pty Ltd
PF	Pfizer Australia Pty Ltd
PK	Fresenius Kabi Australia Pty Limited
RA	Sun Pharma ANZ Pty Ltd
RF	Arrow Pharma Pty Ltd
RO	Roche Products Pty Ltd
RQ	Reach Pharmaceuticals Pty Ltd
RZ	Dr Reddy's Laboratories (Australia) Pty Ltd
SG	Merck Healthcare Pty Ltd
SZ	Sandoz Pty Ltd
TB	Teva Pharma Australia Pty Ltd
TK	Takeda Pharmaceuticals Australia Pty. Ltd.
TS	Specialised Therapeutics Australia Pty Ltd
TX	Apotex Pty Ltd
TY	Apotex Pty Ltd
WM	MEDISON PHARMA AUSTRALIA PTY LIMITED
XC	Southern Cross Pharma Pty Ltd
YG	EUGIA PHARMA (AUSTRALIA) PTY LTD
ZL	Specialised Therapeutics Pharma Pty Ltd

Generic/Proprietary Index

Abevmy[<i>SZ</i>] (bevacizumab 100 mg/4 mL injection, 4 mL vial)	59, 133	Bortezomib Baxter[<i>BX</i>] (bortezomib 3.5 mg injection, 1 vial)	73, 147
Abevmy[<i>SZ</i>] (bevacizumab 400 mg/16 mL injection, 16 mL vial)	59, 133	BORTEZOMIB EUGIA[<i>YG</i>] (bortezomib 3.5 mg injection, 1 vial)	73, 147
Abraxane[<i>TS</i>] (paclitaxel (as nanoparticle albumin-bound) 100 mg injection, 1 vial).....	12, 13, 86, 87	Bortezomib Ever Pharma[<i>IT</i>] (bortezomib 2.5 mg/mL injection, 1 mL vial)	73, 147
Adcetris[<i>TK</i>] (brentuximab vedotin 50 mg injection, 1 vial)	63, 64, 65, 66, 67, 137, 138, 139, 140, 141	Bortezomib Ever Pharma[<i>IT</i>] (bortezomib 3.5 mg/1.4 mL injection, 1.4 mL vial)	73, 147
Adriamycin[<i>PF</i>] (doxorubicin hydrochloride 200 mg/100 mL injection, 100 mL vial).....	14, 88	Bortezomib Juno[<i>JU</i>] (bortezomib 2.5 mg injection, 1 vial)	73, 147
Adriamycin[<i>PF</i>] (doxorubicin hydrochloride 50 mg/25 mL injection, 25 mL vial)	14, 88	Bortezomib Juno[<i>JU</i>] (bortezomib 3.5 mg injection, 1 vial)	73, 147
Akynzeo IV[<i>JZ</i>]	156	Bortezomib Sandoz[<i>SZ</i>] (bortezomib 3.5 mg injection, 1 vial)	73, 147
Akynzeo[<i>JZ</i>].....	156	BORTEZOMIB-TEVA[<i>TB</i>] (bortezomib 3.5 mg injection, 1 vial)	73, 147
Aloxi[<i>JZ</i>].....	157	BRENTUXIMAB VEDOTIN.....	62, 63, 64, 65, 66, 136, 137, 138, 139, 140
APO-Ondansetron[<i>TX</i>]	157	CABAZITAXEL	12, 86
APREPITANT	157	Cabazitaxel Accord[<i>OC</i>] (cabazitaxel 60 mg/3 mL injection, 3 mL vial)	12, 86
Aprepitant APOTEX[<i>TX</i>].....	158	Cabazitaxel Ever Pharma[<i>IT</i>] (cabazitaxel 60 mg/6 mL injection, 6 mL vial)	12, 86
APREPITANT SCP (XC).....	158	Cabazitaxel Juno[<i>JU</i>] (cabazitaxel 60 mg/1.5 mL injection [1.5 mL vial] (&) inert substance diluent [4.5 mL vial], 1 pack)	12, 86
APX-Ondansetron (TY).....	157	Caelyx[<i>BX</i>] (doxorubicin hydrochloride (as pegylated liposomal) 20 mg/10 mL injection, 10 mL vial)	14, 88
APX-Ondansetron ODT(TY)	157	Caelyx[<i>BX</i>] (doxorubicin hydrochloride (as pegylated liposomal) 50 mg/25 mL injection, 25 mL vial)	14, 88
ARSENIC	76, 150	CARBOPLATIN	72, 146
Arsenic Trioxide Accord[<i>OC</i>] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials).....	76, 77, 150, 151	Carboplatin Accord[<i>OC</i>] (carboplatin 450 mg/45 mL injection, 45 mL vial)	72, 146
Arsenic Trioxide Juno[<i>JU</i>] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials).....	76, 77, 150, 151	CARFILZOMIB	73, 74, 75, 147, 148, 149
Arsenic Trioxide-AFT[<i>AE</i>] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL ampoules).....	76, 77, 150, 151	CEMIPLIMAB	38, 39, 112, 113
ATEZOLIZUMAB31, 32, 33, 34, 35, 36, 105, 106, 107, 108, 109, 110		CETUXIMAB	28, 29, 30, 102, 103, 104
AVELUMAB	36, 37, 110, 111	CISPLATIN	72, 146
Bavencio[<i>SG</i>] (avelumab 200 mg/10 mL injection, 10 mL vial)	37, 38, 111, 112	Cisplatin Accord[<i>OC</i>] (cisplatin 100 mg/100 mL injection, 100 mL vial)	72, 146
BENDAMUSTINE	8, 82	Cisplatin Accord[<i>OC</i>] (cisplatin 50 mg/50 mL injection, 50 mL vial)	72, 146
Bendamustine Juno[<i>JU</i>] (bendamustine hydrochloride 100 mg injection, 1 vial)	8, 82	CLADRIBINE	
Bendamustine Juno[<i>JU</i>] (bendamustine hydrochloride 25 mg injection, 1 vial)	8, 82	.Chemotherapy items for Private Hospital use	10
Bendamustine Sandoz[<i>SZ</i>] (bendamustine hydrochloride 100 mg injection, 1 vial).....	8, 82	.Chemotherapy items for Public Hospital use	84
Bendamustine Sandoz[<i>SZ</i>] (bendamustine hydrochloride 25 mg injection, 1 vial).....	8, 82	CYCLOPHOSPHAMIDE.....	9, 83
Bendamustine Viatris[<i>AF</i>] (bendamustine hydrochloride 100 mg injection, 1 vial).....	8, 82	CYCLOPHOSPHAMIDE-REACH[<i>RQ</i>] (cyclophosphamide 1 g injection, 1 vial)	9, 83
Bendamustine Viatris[<i>AF</i>] (bendamustine hydrochloride 25 mg injection, 1 vial)	8, 82	CYCLOPHOSPHAMIDE-REACH[<i>RQ</i>] (cyclophosphamide 500 mg injection, 1 vial)	9, 83
Besponsa[<i>PF</i>] (inotuzumab ozogamicin 1 mg injection, 1 vial)	18, 19, 92, 93	CYTARABINE	10, 84
Bevaciptin[<i>LR</i>] (bevacizumab 100 mg/4 mL injection, 4 mL vial)	59, 133	DARATUMUMAB	19, 20, 93, 94, 95, 159, 160, 161, 162
Bevaciptin[<i>LR</i>] (bevacizumab 400 mg/16 mL injection, 16 mL vial).....	59, 133	Darzalex SC[<i>JC</i>]	160, 161, 162
BEVACIZUMAB	59, 133	Darzalex[<i>JC</i>] (daratumumab 100 mg/5 mL injection, 5 mL vial)	19, 20, 21, 93, 94, 95
BLEOMYCIN.....	15, 89	Darzalex[<i>JC</i>] (daratumumab 400 mg/20 mL injection, 20 mL vial)	19, 20, 21, 93, 94, 95
BLINATUMOMAB	59, 60, 61, 133, 134, 135	DBL Bleomycin Sulfate[<i>PF</i>] (bleomycin sulfate 15 000 international units injection, 1 vial)	15, 89
Blinicyto[<i>AN</i>] (blinatumomab 38.5 microgram injection [1 vial] (&) inert substance solution [10 mL vial], 1 pack) 60, 61, 62, 134, 135, 136		DBL Bortezomib[<i>PF</i>] (bortezomib 1 mg injection, 1 vial)	73, 147
Bortezom[<i>CR</i>] (bortezomib 3.5 mg injection, 1 vial)	73, 147	DBL Bortezomib[<i>PF</i>] (bortezomib 2.5 mg injection, 1 vial).....	73, 147
BORTEZOMIB	73, 147	DBL Bortezomib[<i>PF</i>] (bortezomib 3 mg injection, 1 vial)	73, 147
Bortezomib Accord[<i>OC</i>] (bortezomib 1 mg injection, 1 vial)	73, 147	DBL Bortezomib[<i>PF</i>] (bortezomib 3.5 mg injection, 1 vial).....	73, 147
Bortezomib Accord[<i>OC</i>] (bortezomib 2.5 mg/mL injection, 1 mL vial).....	73, 147	DBL Docetaxel Concentrated Injection[<i>PF</i>] (docetaxel 160 mg/16 mL injection, 16 mL vial)	12, 86
Bortezomib Accord[<i>OC</i>] (bortezomib 3.5 mg injection, 1 vial)	73, 147		
Bortezomib Accord[<i>OC</i>] (bortezomib 3.5 mg/1.4 mL injection, 1.4 mL vial)	73, 147		

DBL Docetaxel Concentrated Injection[PF] (docetaxel 80 mg/8 mL injection, 8 mL vial).....	12, 86	Fluorouracil Ebewe[SZ] (fluorouracil 5 g/100 mL injection, 100 mL vial).....	11, 85
DBL Gemcitabine Injection[PF] (gemcitabine 1 g/26.3 mL injection, 26.3 mL vial).....	11, 85	FOLINIC ACID.....	163, 164
DBL Gemcitabine Injection[PF] (gemcitabine 2 g/52.6 mL injection, 52.6 mL vial).....	11, 85	Folotyn[MF] (pralatrexate 20 mg/mL injection, 1 mL vial) 10,	84
DBL Methotrexate[PF] (methotrexate 1 g/10 mL injection, 10 mL vial).....	9, 83	FOSAPREPITANT.....	158
DBL Methotrexate[PF] (methotrexate 5 mg/2 mL injection, 5 x 2 mL vials).....	9, 83	FOSAPREPITANT MEDSURGE(DZ).....	159
DBL Methotrexate[PF] (methotrexate 50 mg/2 mL injection, 5 x 2 mL vials).....	9, 83	FOSAPREPITANT MSN (RQ).....	159
DBL Methotrexate[PF] (methotrexate 500 mg/20 mL injection, 20 mL vial).....	9, 83	FOSAPREPITANT-AFT (AE).....	159
DBL Vinblastine[PF] (vinblastine sulfate 10 mg/10 mL injection, 5 x 10 mL vials).....	11, 85	FOSNETUPITANT + PALONOSETRON.....	156
DBL Vincristine Sulfate[PF] (vincristine sulfate 1 mg/mL injection, 5 x 1 mL vials).....	11, 85	Gazyva[RO] (obinutuzumab 1 g/40 mL injection, 40 mL vial).....	15, 16, 17, 89, 90, 91
DOCETAXEL.....	12, 86	GEMCITABINE.....	11, 85
Docetaxel Accord[OC] (docetaxel 160 mg/8 mL injection, 8 mL vial).....	12, 86	GEMTUZUMAB OZOGAMICIN.....	68, 69, 142, 143
Docetaxel Accord[OC] (docetaxel 80 mg/4 mL injection, 4 mL vial).....	12, 86	GRANISETRON.....	156
DOSTARLIMAB.....	39, 113	Granisetron Kabi (PK).....	156
DOXORUBICIN.....	14, 88	Granisetron-AFT(AE).....	156
Doxorubicin ACC[OC] (doxorubicin hydrochloride 200 mg/100 mL injection, 100 mL vial).....	14, 88	Halaven[EI] (eribulin mesilate 1 mg/2 mL injection, 2 mL vial).....	77, 151
DOXORUBICIN HYDROCHLORIDE (AS PEGYLATED LIPOSOMAL).....	14, 88	Herceptin SC(RO).....	162, 163
DURVALUMAB.....	40, 41, 114, 115	Herzuma[EW] (trastuzumab 150 mg injection, 1 vial) 22, 23,	24, 25, 96, 97, 98, 99
ELOTUZUMAB.....	67, 141	Herzuma[EW] (trastuzumab 440 mg injection [1 vial] (& inert substance diluent [20 mL vial], 1 pack)....	22, 23, 24, 25, 96, 97, 98, 99
Emend IV(MK).....	159	Holoxan[BX] (ifosfamide 1 g injection, 1 vial).....	9, 83
Empliciti[BQ] (elotuzumab 300 mg injection, 1 vial) ..	67, 68, 141, 142	Holoxan[BX] (ifosfamide 2 g injection, 1 vial).....	9, 83
Empliciti[BQ] (elotuzumab 400 mg injection, 1 vial) ..	67, 68, 141, 142	Hycamtin[SZ] (topotecan 4 mg injection, 5 vials).....	13, 87
Endoxan[BX] (cyclophosphamide 1 g injection, 1 vial) 9,	83	IDARUBICIN.....	14, 88
Endoxan[BX] (cyclophosphamide 2 g injection, 1 vial) 9,	83	IFOSFAMIDE.....	9, 83
ENFORTUMAB VEDOTIN.....	68, 142	Imfinzi[AP] (durvalumab 120 mg/2.4 mL injection, 2.4 mL vial).....	40, 41, 114, 115
Enhertu[AP] (trastuzumab deruxtecan 100 mg injection, 1 vial).....	26, 100	Imfinzi[AP] (durvalumab 500 mg/10 mL injection, 10 mL vial).....	40, 41, 114, 115
EPIRUBICIN.....	14, 88	INOTUZUMAB OZOGAMICIN.....	18, 92
Epirubicin Accord[OC] (epirubicin hydrochloride 200 mg/100 mL injection, 100 mL vial).....	14, 88	IPILIMUMAB.....	69, 70, 71, 143, 144, 145
Erbitux[SG] (cetuximab 100 mg/20 mL injection, 20 mL vial).....	28, 29, 30, 102, 103, 104	IRINOTECAN.....	13, 87
Erbitux[SG] (cetuximab 500 mg/100 mL injection, 100 mL vial).....	28, 29, 30, 102, 103, 104	Irinotecan Accord[OC] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial).....	13, 87
ERIBULIN.....	77, 151	Irinotecan Accord[OC] (irinotecan hydrochloride trihydrate 500 mg/25 mL injection, 25 mL vial).....	13, 87
Etopophos[LM] (etoposide phosphate 1.136 g (etoposide 1 g) injection, 1 vial).....	12, 86	Irinotecan Alphapharm[AF] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial).....	13, 87
ETOPOSIDE.....	12, 86	Irinotecan Alphapharm[AF] (irinotecan hydrochloride trihydrate 500 mg/25 mL injection, 25 mL vial).....	13, 87
Etoposide Ebewe[SZ] (etoposide 100 mg/5 mL injection, 5 x 5 mL vials).....	12, 86	IRINOTECAN BAXTER[BX] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial).....	13, 87
FLUDARABINE.....	10, 84	Jemperli[GK] (dostarlimab 500 mg/10 mL injection, 10 mL vial).....	39, 40, 113, 114
Fludarabine Ebewe[SZ] (fludarabine phosphate 50 mg/2 mL injection, 5 x 2 mL vials).....	10, 84	Kadcyla[RO] (trastuzumab emtansine 100 mg injection, 1 vial).....	27, 28, 101, 102
Fludarabine Juno[JO] (fludarabine phosphate 50 mg injection, 1 vial).....	10, 84	Kadcyla[RO] (trastuzumab emtansine 160 mg injection, 1 vial).....	27, 28, 101, 102
FLUOROURACIL.....	10, 11, 84, 85	Kanjinti[JU] (trastuzumab 150 mg injection, 1 vial)....	22, 23, 24, 25, 96, 97, 98, 99
Fluorouracil Accord[OC] (fluorouracil 1 g/20 mL injection, 20 mL vial).....	11, 85	Kanjinti[JU] (trastuzumab 420 mg injection, 1 vial)....	22, 23, 24, 25, 96, 97, 98, 99
Fluorouracil Accord[OC] (fluorouracil 2.5 g/50 mL injection, 50 mL vial).....	11, 85	Keytruda[MK] (pembrolizumab 100 mg/4 mL injection, 4 mL vial)48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 122,	123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133
Fluorouracil Accord[OC] (fluorouracil 5 g/100 mL injection, 100 mL vial).....	11, 85	Kimmtrak[WM] (tebentafusp 100 microgram/0.5 mL injection, 0.5 mL vial).....	78, 79, 152, 153
Fluorouracil Accord[OC] (fluorouracil 500 mg/10 mL injection, 10 mL vial).....	11, 85	Kyprolis[AN] (carfilzomib 10 mg injection, 1 vial) .74, 75, 76,	148, 149, 150
Fluorouracil Ebewe[SZ] (fluorouracil 1 g/20 mL injection, 20 mL vial).....	11, 85	Kyprolis[AN] (carfilzomib 30 mg injection, 1 vial) .74, 75, 76,	148, 149, 150
		Kyprolis[AN] (carfilzomib 60 mg injection, 1 vial) .74, 75, 76,	148, 149, 150
		Kytril(IX).....	156
		Leucovorin Calcium (Hospira Pty Limited)(PF).....	164

Leucovorin Calcium (Pfizer Australia Pty Ltd)(PF).....	163
Leustatin[IX] (cladribine 10 mg/10 mL injection, 10 mL vial)	
.....	10, 84
Libtayo[WM] (cemiplimab 350 mg/7 mL injection, 7 mL vial)	
.....	38, 39, 112, 113
Liposomal Doxorubicin SUN[RA] (doxorubicin hydrochloride (as pegylated liposomal) 20 mg/10 mL injection, 10 mL vial)	14, 88
Liposomal Doxorubicin SUN[RA] (doxorubicin hydrochloride (as pegylated liposomal) 50 mg/25 mL injection, 25 mL vial)	14, 88
Litak[AF] (cladribine 10 mg/5 mL injection, 5 mL vial)	10, 84
MESNA.....	164
METHOTREXATE	
. Chemotherapy items for Private Hospital use.....	9
. Chemotherapy items for Public Hospital use.....	83
Methotrexate Accord[OD] (methotrexate 1 g/10 mL injection, 10 mL vial)	9, 83
Methotrexate Ebewe[SZ] (methotrexate 5 g/50 mL injection, 50 mL vial)	9, 83
MITOZANTRONE.....	15, 89
Mitozantrone Ebewe[SZ] (mitozantrone 20 mg/10 mL injection, 10 mL vial)	15, 89
MSN Cabazitaxel[RQ] (cabazitaxel 60 mg/1.5 mL injection [1.5 mL vial] (& inert substance diluent [4.5 mL vial], 1 pack)	12, 86
Mvasi[AN] (bevacizumab 100 mg/4 mL injection, 4 mL vial)	59, 133
Mvasi[AN] (bevacizumab 400 mg/16 mL injection, 16 mL vial)	59, 133
MYCOBACTERIUM BOVIS BCG DANISH STRAIN.....	163
MYCOBACTERIUM BOVIS BCG TICE STRAIN.....	163
Mylotarg[PF] (gemtuzumab ozogamicin 5 mg injection, 1 vial)	69, 143
NANOPARTICLE ALBUMIN-BOUND PACLITAXEL	12, 86
Navelbine[FB] (vinorelbine 50 mg/5 mL injection, 5 mL vial)	11, 85
NETUPITANT + PALONOSETRON.....	156
NIVOLUMAB..	41, 42, 43, 44, 45, 46, 47, 48, 115, 116, 117, 118, 119, 120, 121, 122
OBINUTUZUMAB.....	15, 16, 17, 89, 90, 91
Ogivri[SZ] (trastuzumab 150 mg injection, 1 vial)	22, 23, 24, 25, 96, 97, 98, 99
Omegapharm Irinotecan[OE] (irinotecan hydrochloride trihydrate 100 mg/5 mL injection, 5 mL vial)	13, 87
Omegapharm Irinotecan[OE] (irinotecan hydrochloride trihydrate 40 mg/2 mL injection, 2 mL vial)	13, 87
OncoTICE(MK).....	163
ONDANSETRON.....	156, 157
Ondansetron Mylan ODT (AF).....	157
Ondansetron Mylan Tablets (AF).....	157
Ondansetron ODT Viatris (AL).....	157
Ondansetron ODT-DRLA(RZ).....	157
Ondansetron SZ ODT (HX).....	157
Ondansetron SZ ODT(HX).....	157
Ondansetron SZ(HX).....	157
Ondansetron Tablets Viatris (AL).....	157
Ondansetron-DRLA(RZ).....	157
Onkotrone[BX] (mitozantrone 25 mg/12.5 mL injection, 12.5 mL vial)	15, 89
Opdivo[BQ] (nivolumab 100 mg/10 mL injection, 10 mL vial)	42, 43, 44, 45, 46, 47, 48, 116, 117, 118, 119, 120, 121, 122
Opdivo[BQ] (nivolumab 40 mg/4 mL injection, 4 mL vial)	42, 43, 44, 45, 46, 47, 48, 116, 117, 118, 119, 120, 121, 122
OXALIPLATIN.....	72, 147
Oxaliplatin Accord[OC] (oxaliplatin 100 mg/20 mL injection, 20 mL vial)	72, 147
Oxaliplatin SUN[RA] (oxaliplatin 100 mg/20 mL injection, 20 mL vial)	72, 147
Oxaliplatin SUN[RA] (oxaliplatin 200 mg/40 mL injection, 40 mL vial)	72, 147
PACLITAXEL.....	13, 87
Paclitaxel Accord[OC] (paclitaxel 300 mg/50 mL injection, 50 mL vial)	13, 87
Paclitaxel Ebewe[SZ] (paclitaxel 300 mg/50 mL injection, 50 mL vial)	13, 87
Padcev[LL] (enfortumab vedotin 20 mg injection, 1 vial)	68, 142
Padcev[LL] (enfortumab vedotin 30 mg injection, 1 vial)	68, 142
PALONOSETRON.....	157
Palonosetron Dr.Reddy's (RZ).....	157
PALONOSETRON Medsurge(DZ).....	157
PANITUMUMAB.....	30, 31, 104, 105
PEMBROLIZUMAB ...	48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132
PEMETREXED.....	9, 83
Pemetrexed Accord[OD] (pemetrexed 1 g injection, 1 vial)	9, 83
Pemetrexed Accord[OD] (pemetrexed 100 mg injection, 1 vial)	9, 83
Pemetrexed Accord[OD] (pemetrexed 500 mg injection, 1 vial)	9, 83
Pemetrexed APOTEX[TX] (pemetrexed 500 mg injection, 1 vial)	9, 83
Pemetrexed Ever Pharma[IT] (pemetrexed 1 g/40 mL injection, 40 mL vial)	9, 83
Pemetrexed Ever Pharma[IT] (pemetrexed 100 mg/4 mL injection, 4 mL vial)	9, 83
Pemetrexed Ever Pharma[IT] (pemetrexed 500 mg/20 mL injection, 20 mL vial)	9, 83
Pemetrexed SUN[RA] (pemetrexed 1 g injection, 1 vial)	9, 83
Pemetrexed SUN[RA] (pemetrexed 100 mg injection, 1 vial)	10, 84
Pemetrexed SUN[RA] (pemetrexed 500 mg injection, 1 vial)	10, 84
Perjeta[RO] (pertuzumab 420 mg/14 mL injection, 14 mL vial)	21, 22, 95, 96
PERTUZUMAB.....	21, 22, 95, 96
Pfizer Australia Pty Ltd[PF] (cytarabine 100 mg/5 mL injection, 5 x 5 mL vials)	
. Chemotherapy items for Private Hospital use.....	10
. Chemotherapy items for Public Hospital use.....	84
Phenasen[FF] (arsenic trioxide 10 mg/10 mL injection, 10 x 10 mL vials)	76, 77, 150, 151
PRALATREXATE.....	10, 84
RALTITREXED.....	10, 84
Ribomustin[JC] (bendamustine hydrochloride 100 mg injection, 1 vial)	8, 82
Ribomustin[JC] (bendamustine hydrochloride 25 mg injection, 1 vial)	8, 82
RITUXIMAB	
. Chemotherapy items for Private Hospital use.....	17
. Chemotherapy items for Public Hospital use.....	91
Riximyo[SZ] (rituximab 100 mg/10 mL injection, 2 x 10 mL vials)	17, 91
Riximyo[SZ] (rituximab 500 mg/50 mL injection, 50 mL vial)	17, 91
Ruxience[PF] (rituximab 100 mg/10 mL injection, 10 mL vial)	
. Chemotherapy items for Private Hospital use.....	17
. Chemotherapy items for Public Hospital use.....	91
Ruxience[PF] (rituximab 500 mg/50 mL injection, 50 mL vial)	
. Chemotherapy items for Private Hospital use.....	17
. Chemotherapy items for Public Hospital use.....	91
SACITUZUMAB GOVITECAN.....	72, 146

TEBENTAFUSP.....	77, 78, 79, 151, 152, 153
Tecentriq[RO] (atezolizumab 1.2 g/20 mL injection, 20 mL vial)	32, 33, 34, 35, 36, 106, 107, 108, 109, 110
Tecentriq[RO] (atezolizumab 840 mg/14 mL injection, 14 mL vial).....	32, 34, 35, 106, 108, 109
Tomudex[PF] (raltitrexed 2 mg injection, 1 vial).....	10, 84
TOPOTECAN	13, 87
Topotecan Accord[OC] (topotecan 4 mg/4 mL injection, 5 x 4 mL vials).....	13, 87
TRABECTEDIN	13, 14, 87, 88
TRASTUZUMAB.....	22, 23, 24, 25, 96, 97, 98, 99, 162, 163
TRASTUZUMAB DERUXTECAN	25, 99
TRASTUZUMAB EMTANSINE	26, 27, 100, 101
Trazimera[PF] (trastuzumab 150 mg injection, 1 vial).....	22, 23, 24, 25, 96, 97, 98, 99
Trazimera[PF] (trastuzumab 60 mg injection, 1 vial).....	22, 23, 24, 25, 96, 97, 98, 99
Trodelvy[GJ] (sacituzumab govitecan 180 mg injection, 1 vial)	72, 146
Truxima[EW] (rituximab 100 mg/10 mL injection, 2 x 10 mL vials).....	17, 91
Truxima[EW] (rituximab 500 mg/50 mL injection, 50 mL vial)	17, 91
Uromitexan(BX)	164
Vectibix[AN] (panitumumab 100 mg/5 mL injection, 5 mL vial)	31, 105
Vectibix[AN] (panitumumab 400 mg/20 mL injection, 20 mL vial)	31, 105
VesiCulture(LM)	163
VINBLASTINE	11, 85
VINCRIStINE.....	11, 85
VINOReLBINE	11, 85
Vinorelbine Ebewe[SZ] (vinorelbine 10 mg/mL injection, 1 mL vial)	11, 85
Vinorelbine Ebewe[SZ] (vinorelbine 50 mg/5 mL injection, 5 mL vial)	11, 85
Yervoy[BQ] (ipilimumab 200 mg/40 mL injection, 40 mL vial)	71, 146
Yervoy[BQ] (ipilimumab 50 mg/10 mL injection, 10 mL vial)	70, 71, 144, 145, 146
Yondelis[ZL] (trabectedin 1 mg injection, 1 vial) ..	13, 14, 87, 88
Zavedos Solution[PF] (idarubicin hydrochloride 5 mg/5 mL injection, 5 mL vial)	15, 89
Zofran syrup 50 mL(AS)	157
Zofran(AS).....	157
Zotren 4 (RF)	157
Zotren 8 (RF).....	157
Zotren ODT (RF)	157
Zotren ODT(RF)	157